1

Introduction

Established in 1947, Brookhaven National Laboratory is a multi-program national laboratory managed for the U.S. Department of Energy by Brookhaven Science Associates (BSA), a partnership formed by Stony Brook University and Battelle Memorial Institute. BSA has been managing and operating the Laboratory under a performance-based contract with DOE since 1998. From 1947 to 1998, BNL was operated by Associated Universities Incorporated. Prior to 1947, the site operated as Camp Upton, a U.S. Army training camp, which was active from 1917 to 1920 during World War I and from 1940 to 1946 during World War II.

BNL is one of 10 national Laboratories under DOE's Office of Science, which provides most of the Laboratory's research dollars and direction. BNL has a history of outstanding scientific achievements. For over 60 years, Laboratory researchers have successfully worked to envision, construct, and operate large and innovative scientific facilities in pursuit of research advances in many fields. Programs in place at BNL emphasize continual improvement in environmental, safety, security, and health performance.

1.1 LABORATORY MISSION

BNL's broad mission is to carry out basic and applied research in long-term programs in a safe and environmentally sound manner with the cooperation, support, and involvement of its scientific and local communities. The fundamental elements of the Laboratory's role in support of DOE's strategic missions in energy resources, environmental quality, and national security are:

- To conceive, construct, and operate complex, leading-edge, user-oriented research facilities.
- To develop advanced technologies that address national needs and initiate their transfer to other organizations and to the commercial sector.
- To disseminate technical knowledge to educate future generations of scientists and engineers.
- To maintain technical currency in the nation's workforce and encourage scientific awareness in the general public.

BNL's Environmental, Safety, Security, and Health (ESSH) Policy is the Laboratory's

commitment to continual improvement in ESSH performance. Under this policy, the Laboratory's goals are to protect the environment, conserve resources, and prevent pollution; maintain a safe workplace by planning work and performing it safely; provide security for people, property, information, computing systems, and facilities; protect human health within our boundaries and in the surrounding community; achieve and maintain compliance with applicable ESSH requirements; maintain an open, proactive, and constructive relationship with employees, neighbors, regulators, DOE, and other stakeholders; and continually improve ESSH performance.

BNL was the first DOE Office of Science National Laboratory to be registered under the prestigious International ISO 14001 environmental management standard in 2001. In addition, in December 2006, BNL was the first DOE Laboratory to achieve full registration under the Occupational Health and Safety Assessment Series (OHSAS) 18001 Standard. These programs are described in detail in Chapter 2 of this report. Registration to these standards was maintained throughout 2008.

1.2 HISTORY

BNL was founded in 1947 by the Atomic Energy Commission (AEC), a predecessor to the present DOE. AEC provided the initial funding for BNL's research into peaceful uses of the atom. The objective was to promote basic research in the physical, chemical, biological, and engineering aspects of the atomic sciences. The result was the creation of a regional laboratory to design, construct, and operate large scientific machines that individual institutions could not afford to develop on their own.

Although BNL no longer operates any research reactors, the Laboratory's first major scientific facility was the Brookhaven Graphite Research Reactor (BGRR), which was the first reactor to be constructed in the United States following World War II. The reactor's primary mission was to produce neutrons for scientific experimentation in the fields of medicine, biology, chemistry, physics, and nuclear technology. The BGRR operated from 1950 to 1968 and is currently being decommissioned and will be dismantled. The BGRR will undergo long-term routine inspection and surveillance when decommissioning is complete.

The BGRR's research capacity was replaced and surpassed in 1965 by the High Flux Beam Reactor (HFBR). The HFBR was used solely for scientific research and provided neutrons for experiments in materials science, chemistry, biology, and physics. For more than 30 years, the HFBR was one of the premier neutron beam reactors in the world. In 1997, workers discovered that a leak in the HFBR spent fuel storage pool had been releasing tritium to the groundwater (see Chapter 7 for further details). In November 1999, the Secretary of Energy decided that the HFBR would be permanently shut down. In 2008, actions continued to prepare the HFBR for permanent decontamination and dismantling (D&D). With input from the community, a final Record of Decision was approved outlining the remedy for the D&D project. To date, completed actions include the removal and disposal of HFBR fuel and primary coolant, shipment of

equipment for reuse at other facilities, cleanup and transfer of the Cold Neutron Facility for reuse, dismantling of many ancillary buildings in the HFBR complex, and removal and disposal of the reactor control rod blades and beam plugs. Near-term actions include the dismantling of the remaining ancillary buildings, removal of contaminated underground utilities and piping, and preparation of the confinement building for safe storage. An associated action conducted in 2008 was the cleanup of the Waste Loading Area, which was previously used for staging cleanup-derived wastes prior to shipping the materials for disposal via railway.

Medical research at BNL began in 1950 with the opening of one of the first hospitals devoted to nuclear medicine. It was followed by the Medical Research Center in 1958 and the Brookhaven Medical Research Reactor (BMRR) in 1959. The BMRR was the first nuclear reactor in the nation to be constructed specifically for medical research. Due to a reduction of research funding, the BMRR was shut down in December 2000. All spent fuel from the BMRR has been removed and transported off site, and the facility is currently in a "cold" shutdown mode as a radiological facility.

The Brookhaven Linac Isotope Producer (BLIP) was built in 1973. It creates radioactive forms of ordinary chemical elements that can be used alone or incorporated into radiotracers for use in nuclear medicine research or for clinical diagnosis and treatment. BNL's Center for Translational Neuroimaging (CTN) uses brainimaging tools, including positron emission tomography (PET) and magnetic resonance imaging (MRI) equipment, to research causes of, and treatments for, brain diseases such as drug addiction, appetite disorders, attention deficit disorder, and neurodegenerative disease. The development of PET and MRI also has helped facilitate the development of new drugs for physicians worldwide to treat patients for cancer and heart disease.

High-energy particle physics research at BNL began in 1952 with the Cosmotron, the first particle accelerator to achieve billion-electron-volt energies. Work at the Cosmotron resulted in a Noble Prize in 1957. After 14 years of service,

the Cosmotron ceased operation and was dismantled due to design limitations that restricted the energies it could achieve. The Alternating Gradient Synchrotron (AGS), a much larger particle accelerator, became operational in 1960. The AGS allowed scientists to accelerate protons to energies that yielded many discoveries of new particles and phenomena, for which BNL researchers were awarded three Nobel Prizes in physics. The AGS receives protons from BNL's linear accelerator (Linac), designed and built in the late 1960s as a major upgrade to the AGS complex. The Linac's purpose is to provide accelerated protons for use at AGS facilities and BLIP. The AGS booster, constructed in 1991, further enhanced the capabilities of the AGS, enabling it to accelerate protons and heavy ions to even higher energies. The Tandem Van de Graaff accelerator began operating in 1970 and is the starting point of the chain of accelerators that provide ions of gold, other heavy metals, and protons for experiments at the Relativistic Heavy Ion Collider (RHIC).

RHIC began operation in 2000. Inside this two-ringed particle accelerator, two beams of gold ions, heavy metals, or protons circulating at nearly the speed of light, collide head-on, releasing large amounts of energy. RHIC is used to study what the universe may have looked like in the first few moments after its creation, offering insights into the fundamental forces and properties of matter. Planned upgrades to RHIC will expand the facility's research capabilities. The first upgrade, RHIC II, will increase the collider's collision rate and improve the sensitivity of the large detectors it uses. Another planned upgrade, the eRHIC, will add a highenergy electron ring to create the world's only electron-heavy ion collider, which physicists expect will probe a new form of matter.

The NASA Space Radiation Laboratory (NSRL) became operational in 2003. It is jointly managed by DOE's Office of Science and NASA's Johnson Space Center. The NSRL uses heavy ions extracted from the AGS booster to produce beams of radiation similar to radiation that would be encountered by astronauts on long missions. Studies are conducted to assess risks and test protective measures. The NSRL

is one of the few facilities in the world that can simulate the harsh cosmic and solar radiation environment found in space.

The National Synchrotron Light Source (NSLS) uses a linear accelerator and booster synchrotron to guide charged particles in orbit inside two electron storage rings for use in a wide range of physical and biological experiments. The NSLS produces beams of very intense light in the x-ray, ultraviolet, and infrared spectra, allowing scientists to study the structure of proteins, investigate the properties of new materials, and understand the fate of chemicals in the environment. Although the current NSLS has been continually updated since its commissioning in 1982, today the practical limits of its performance have been reached. To continue advances in these fields, the NSLS-II has been conceived as the next generation synchrotron light source. To help meet the critical scientific challenges of our energy future, this new stateof-the-art, medium-energy electron storage ring synchrotron will provide x-rays more than 10,000 times brighter than the current NSLS and will focus on research at the nanoscale. The NSLS-II will enable scientists to focus on some of the nation's most important scientific challenges at the nanoscale level, including clean, affordable energy, molecular electronics, and high-temperature superconductors. Construction of the new facility officially started in October 2008.

The Laboratory's Research Support Building (RSB) was completed in 2006, and provides administrative and support functions in a single location for employees and visiting scientists. The RSB has been awarded the Leadership in Energy and Environmental Design (LEED) Silver certification from the U.S. Green Building Council. The award is based on five categories: sustainability, water efficiency, energy and atmosphere, materials and resources, and indoor environmental quality.

Construction of the Center for Functional Nanomaterials (CFN) began in 2005 and was completed in May 2007. The CFN provides state-of-the-art capabilities for the fabrication and study of nanoscale materials, with an emphasis on atomic-level tailoring to achieve de-



sired properties and functions. Nanoscience has the potential to bring about and accelerate new technologies in energy distribution, drug delivery, sensors, and industrial processes. The CFN is a science-based user facility, used for developing strong scientific programs while offering broad access to its capabilities and collaboration through an active user program. It is one of five Nanoscale Science Research Centers funded by DOE's Office of Science and supports the Laboratory's goal of leadership in the development of advanced materials and processes for energy applications. Like the RSB, the CFN has also been awarded LEED Silver certification.

In addition, groundbreaking is expected in 2009 for the new Interdisciplinary Science Building (ISB), an energy-efficient and environmentally sustainable building that will provide labs, offices, and support functions to bring together a broad spectrum of researchers in a single location to foster energy research.

In 2008, the Laboratory was a proposed host-site for the construction of the largest solar energy project in New York State. Under this project, the Long Island Power Authority sought proposals for the construction of up to 50 megawatts of solar-generated power. This project, if awarded, will help the U.S. become less reliant on foreign energy sources and help meet renewable energy goals. In addition, a solar photovoltaic research and development facility may be developed for research, education, and community outreach to foster increased knowledge in the use of solar energy. The outcome of this proposal is expected to be finalized in 2009.

Past operations and research at the BNL site dating back to the early 1940s when it was Camp Upton have resulted in localized environmental contamination. As a result, the Laboratory was added to the federal Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) National Priorities List of contaminated sites in 1989. One of 27 sites on Long Island identified for priority cleanup, BNL has made significant progress toward improving environmental operations and remediating past contamination. DOE will continue to fund cleanup projects until the Laboratory is restored and removed from the National Priori-

ties List. Major accomplishments in cleanup activities at BNL are discussed further throughout this report.

1.3 RESEARCH AND DISCOVERIES

BNL conducts research in nuclear and highenergy physics; the physics and chemistry of materials; nanoscience; energy and environmental research; national security and nonproliferation; neurosciences and medical imaging; structural biology; and computational sciences. BNL's world-class research facilities are also available to university, industrial, and government personnel.

To date, six Nobel Prizes have been awarded for discoveries made wholly or partly at BNL. Some significant discoveries and developments made at the Laboratory include L-dopa, used to treat Parkinson's disease; the first synthesis of human insulin; the use of x-rays and neutrons to study biological specimens; the radionuclide thallium-201, used in millions of cardiac stress tests each year; the radionuclide technetium-99, also used to diagnose heart disease; x-ray angiography for noninvasive cardiac imaging; research on solar neutrinos and how they change form as they move through space; magnetically-levitated (magley) trains; energy technologies studies; and researching pollutioneating bacteria.

Examples of current research at the Laboratory include the investigation of new nanostructures and nanoparticles; the development of high-temperature superconductors; novel states of matter being revealed at RHIC; medical imaging techniques to investigate the brain mechanisms underlying drug addiction, psychiatric disorders, and metabolism; new methods of understanding the earth's climate; and research into how infections begin.

1.4 FACILITIES AND OPERATIONS

Most of the Laboratory's principal facilities are located near the center of the site. The developed area is approximately 1,650 acres:

- 500 acres originally developed by the Army (as part of Camp Upton) and still used for offices and other operational buildings
- 200 acres occupied by large, specialized



research facilities

- 550 acres used for outlying facilities, such as the Sewage Treatment Plant, research agricultural fields, housing facilities, and fire breaks
- 400 acres of roads, parking lots, and connecting areas

The balance of the site, approximately 3,600 acres, is mostly wooded and represents the native pine barrens ecosystem.

The major scientific facilities at BNL are briefly described in Figure 1-1. Additional facilities, shown in Figure 1-2 and briefly described below, support BNL's science and technology mission by providing basic utility and environmental services.


- Central Chilled Water Plant. This plant provides chilled water sitewide for air conditioning and process refrigeration via underground piping. The plant has a large refrigeration capacity and reduces the need for local refrigeration plants and air conditioning.
- Central Steam Facility (CSF). This facility provides high-pressure steam for facility and process heating sitewide. Either natural gas or fuel oil can be used to produce the steam, which is conveyed to other facilities through underground piping. Condensate is collected and returned to the CSF for reuse, to conserve water and energy.
- Fire Station. The Fire Station houses six response vehicles. The BNL Fire Rescue Group provides on-site fire suppression, emergency medical services, hazardous material response, salvage, and property protection.
- Major Petroleum Facility (MPF). This facility provides reserve fuel for the CSF during times of peak operation. With a total capacity of 2.3 million gallons, the MPF primarily stores No. 6 fuel oil. The 1997 conversion of CSF boilers to burn natural gas as well as oil has significantly reduced the Laboratory's reliance on oil as a sole fuel source when other fuels are more economical.
- Sewage Treatment Plant (STP). This plant treats sanitary and certain process wastewa-

- ter from BNL facilities prior to discharge into the Peconic River, similar to the operations of a municipal sewage treatment plant. The plant has a design capacity of 3 million gallons per day. Effluent is monitored and controlled under a permit issued by the New York State Department of Environmental Conservation (NYSDEC).
- Waste Concentration Facility (WCF). This facility was previously used for the receipt, processing, and volume reduction of aqueous radioactive waste. At present, the WCF houses equipment and auxiliary systems required for operation of the liquid lowlevel radioactive waste storage and pump systems.
- Waste Management Facility (WMF). This facility is a state-of-the-art complex for managing the wastes generated from BNL's research and operations activities. The facility was built with advanced environmental protection systems and features, and began operation in December 1997.
- *Water Treatment Plant (WTP)*. The potable water treatment plant has a capacity of 5 million gallons per day. Potable water is obtained from six on-site wells. Three wells located along the western boundary of the site are treated at the WTP with a lime-softening process to remove naturally occurring iron and by the addition of sodium hypochlorite for bacterial control. The plant is also equipped with dual air-stripping towers to ensure that volatile organic compounds (VOCs) are at or below New York State drinking water standards. Three wells located along the eastern section of the developed site are treated by the addition of sodium hydroxide to increase the pH of the water to make it less corrosive and by the addition of sodium hypochlorite to control bacteria. BNL's potable water met all drinking water standards in 2008.

1.5 LOCATION, LOCAL POPULATION, AND LOCAL ECONOMY

BNL is located on Long Island, 60 miles east of New York City. The Laboratory's 5,265-acre site is near Long Island's geographic center and





1. Relativistic Heavy Ion Collider (RHIC)

The RHIC is a world-class scientific research facility. The RHIC accelerator drives two intersecting beams of gold ions, other heavy metal ions, and protons head on to form subatomic collisions. What physicists learn from these collisions may help us understand more about why the physical world works the way it does, from the smallest subatomic particles, to the largest stars. Current RHIC experiments include the Solenoidal Tracker at RHIC (STAR), a detector used to track particles produced by ion collisions; the PHENIX detector, used to record different particles emerging from collisions; the Broad Range Hadron Magnetic Spectrometer (BRAHMS), used to study particles as they pass through detectors; and PHOBOS, a detector designed to examine and analyze a very large number of unselected gold ion collisions.

2. Alternating Gradient Synchrotron (AGS)

The AGS is a particle accelerator used to propel protons and heavy ions, such as gold and iron, to high energies for physics research. The Linear Accelerator (Linac) serves as a proton injector for the AGS Booster.

3. AGS Booster

The AGS Booster is a circular accelerator used for physics research and radiobiology studies. It receives either a proton beam from the Linac or heavy ions from the Tandem Van de Graaff and accelerates these before injecting them into the AGS ring for further acceleration. The Booster also serves as the energetic heavy ion source for the NASA Space Radiation Laboratory, which is used to simulate the harsh cosmic and solar radiation environment found in space.

4. Linear Accelerator (Linac) and Brookhaven Linac Isotope Producer (BLIP)

The Linac provides beams of polarized protons for the AGS and RHIC. The excess beam capacity is used to produce radioisotopes for research and medical imaging at the BLIP. The BLIP is one of the nation's key production facilities for radioisotopes, which are crucial to clinical nuclear medicine. The BLIP also supports research on new diagnostic and therapeutic radiopharmaceuticals.

Heavy Ion Transfer Line (HITL)

The HITL connects the Tandem Van de Graaff and the AGS Booster. This interconnection enables the transport of ions of intermediate mass to the AGS Booster, where they are accelerated before injection into the AGS. The ions are then extracted and sent to the AGS experimental area for physics research.

6. Radiation Therapy Facility (RTF)

Part of the Medical Research Center, the RTF is a high-energy dual x-ray mode linear accelerator used for radiation therapy for cancer patients. This accelerator delivers therapeutically useful beams of x-rays and electrons for conventional and advanced medical radiotherapy techniques.

7. Brookhaven Medical Research Reactor (BMRR)

The BMRR was the world's first nuclear reactor built exclusively for medical research and therapy. It produced neutrons in an optimal energy range for experimental treatment of a type of brain cancer known as glioblastoma

multiforme. The BMRR was shut down in December 2000 due to a reduction in medical research fundina.

Scanning Transmission Electron Microscope (STEM)

The STEM facility includes two microscopes, STEM 1 and STEM 3, used for biological research. Both devices allow scientists to see the intricate details of living things, from bacteria to human tissue. The images provide a picture and data that are used in Mass Analysis.

9. Center for Functional Nanomaterials (CFN)

The CFN provides state-of-the-art capabilities for the fabrication and study of nanoscale materials, with an emphasis on atomic-level tailoring to achieve desired properties and functions. The CFN is a science-based user facility, simultaneously developing strong scientific programs while offering broad access to its capabilities and collaboration through an active user program. The overarching scientific theme of the CFN is the development and understanding of nanoscale materials that address the Nations' challenges in energy security, consistent with the Department of Energy mission.

National Synchrotron Light Source (NSLS)

The NSLS uses a linear accelerator and booster synchrotron as an injection system for two electron storage rings that provide intense light spanning the electromagnetic spectrum from the infrared through x-rays. The properties of this light and the 80 specially designed experimental stations, called beamlines, allow scientists to perform a large variety of experiments.

11. High Flux Beam Reactor (HFBR)

The HFBR was one of the premier neutron physics research facilities in the world. Neutron beams produced at the HFBR were used to investigate the molecular structure of materials, which aided in pharmaceutical design and materials development and expanded the knowledge base of physics, chemistry, and biology. The HFBR was permanently shut down in November 1999 and is currently being decommissioned under the Environmental Restoration Program.


12. Tandem Van de Graaff and Cyclotron

These accelerators are used in medium energy physics investigations and for producing special nuclides. The Tandem Van de Graff accelerators are used to bombard materials with ions for manufacturing and testing purposes, and to supply RHIC with heavy ions. The cyclotrons, operated by the Chemistry Department, are used for the production of radiotracers for use in Positron Emission Tomography and Magnetic Resonance Imaging studies.

13. Brookhaven Graphite Research Reactor (BGRR)

The BGRR was the first reactor to be constructed in the United States following World War II. It was used for scientific exploration in the fields of medicine, biology, chemistry, physics, and nuclear engineering. The BGRR is currently being decommissioned under the Environmental Restoration Program.





is part of the Town of Brookhaven, the largest township (both in area and population) in Suffolk County. The Laboratory is one of the five largest, high-technology employers on Long Island, with approximately 2,800 employees that include scientists, engineers, technicians, and administrative personnel. More than 75 percent of BNL employees live and shop in Suffolk County. In addition, BNL annually hosts an estimated 4,000 visiting scientists, more than 30 percent of whom are from New York State universities and businesses. The visiting scientists and sometimes their families, as well as visiting students, reside in apartments and dormitories on site or in nearby communities.

An independent Suffolk County Planning

Commission report concluded that BNL's spending for operations, procurement, payroll, construction, medical benefits, and technology transfer spreads throughout Long Island's economy, making BNL vital to the local economic health, as well as to New York State (Kamer 2006). In 2008, BNL purchased more than \$32.5 million worth of supplies and services from Long Island businesses. Approximately \$5.8 million was spent on 508 purchases in Nassau County, and \$26.6 million was spent on 3,001 purchases in Suffolk County. BNL's total annual budget in 2008 was approximately \$530.9 million, of which approximately 60.7 percent was spent on employee salaries, wages, and fringe benefits.

1.6 GEOLOGY AND HYDROLOGY

BNL is situated on the western rim of the shallow Peconic River watershed. The marshy areas in the northern and eastern sections of the site are part of the headwaters of the Peconic River. Depending on the height of the water table relative to the base of the riverbed, the Peconic River both recharges to, and receives water from, the underlying upper glacial aquifer. In times of sustained drought, the river water recharges to the groundwater; with normal to above-normal precipitation, the river receives water from the aquifer.

In general, the terrain of the BNL site is gently rolling, with elevations varying between 44 and 120 feet above mean sea level. Depth to groundwater from the land surface ranges from 5 feet near the Peconic River to about 80 feet in the higher elevations of the central and western portions of the site. Studies of Long Island hydrology and geology in the vicinity of the Laboratory indicate that the uppermost Pleistocene deposits, composed of highly permeable glacial sands and gravel, are between 120 and 250 feet thick (Warren et al. 1968, Scorca et al. 1999). Water penetrates these deposits readily, and there is little direct runoff into surface streams unless precipitation is intense. The sandy deposits store large quantities of water in the Upper Glacial aquifer. On average, about half of the annual precipitation is lost to the atmosphere through evapotranspiration and the other half percolates through the soil to recharge the groundwater (Koppelman 1978).

The Long Island Regional Planning Board and Suffolk County have identified the Laboratory site as overlying a deep-flow recharge zone for Long Island groundwater (Koppelman 1978). Precipitation and surface water that recharge within this zone have the potential to replenish the Magothy and Lloyd aquifer systems lying below the Upper Glacial aquifer. It has been estimated that up to two-fifths of the recharge from rainfall moves into the deeper aquifers. The extent to which groundwater on site contributes to deep-flow recharge has been confirmed through the use of an extensive network of shallow and deep wells installed at BNL and surrounding areas (Geraghty & Miller

1996). This groundwater system is the primary source of drinking water for both on- and offsite private and public supply wells and has been designated a sole source aquifer system by the Environmental Protection Agency.

During 2008, the Laboratory used approximately 1.15 million gallons of groundwater per day to meet potable water needs and heating and cooling requirements. Approximately 75 percent of the water pumped from BNL supply wells is returned to the aguifer through on-site recharge basins and permitted discharges to the Peconic River. Under normal hydrologic conditions, most of the water discharged to the river recharges to the Upper Glacial aquifer before leaving the site. Human consumption, evaporation (cooling tower and wind losses), and sewer line losses account for the remaining 25 percent. An additional 3.4 million gallons of groundwater were pumped each day from remediation wells. This water is treated to remove contaminants and is then returned to the aquifer by way of recharge basins or injection wells.

Groundwater flow directions across the BNL site are influenced by natural drainage systems: eastward along the Peconic River, southeast toward the Forge River, and south toward the Carmans River (Figure 1-3). Pumping from on-site supply wells affects the direction and speed of groundwater flow, especially in the central, developed areas of the site. The main groundwater divide on Long Island is aligned generally east-west and lies approximately one-half mile north of the Laboratory. Groundwater north of the divide flows northward and ultimately discharges to the Long Island Sound. Groundwater south of the divide flows east and south, discharging to the Peconic River, Peconic Bay, south shore streams, Great South Bay, and Atlantic Ocean. The regional groundwater flow system is discussed in greater detail in Stratigraphy and Hydrologic Conditions at the Brookhaven National Laboratory and Vicinity (Scorca et al. 1999). In most areas at BNL, the horizontal velocity of groundwater is approximately 0.75 to 1.2 feet per day (Geraghty & Miller 1996). In general, this means that groundwater travels for approximately 20 to 22






Figure 1-3. BNL Groundwater Flow Map.

years as it moves from the central, developed area of the site to the Laboratory's southern boundary.


1.7 CLIMATE

The Meteorological Group at BNL has been recording weather data on site since 1949. The Laboratory is broadly influenced by continental and maritime weather systems. Locally, the Long Island Sound, Atlantic Ocean, and associated bays influence wind directions and humidity and provide a moderating influence on extreme summer and winter temperatures. The prevailing ground-level winds at BNL are from the southwest during the summer, from the northwest during the winter, and about equally from those two directions during the spring and fall (Nagle 1975, 1978). Figure 1-4 shows the 2008 annual wind rose for BNL, which depicts

the annual frequency distribution of wind speed and direction, measured at an on-site meteorological tower at heights of 33 feet (10 meters) and 300 feet (88 meters) above land surface.


The average monthly temperature in the area for 2008 was 51.97 degrees Fahrenheit (°F). The average yearly temperature for the area was 50.23°F. Figures 1-5 and 1-6 show the 2008 monthly mean temperatures and the historical annual mean temperatures, respectively.

The total annual precipitation in 2008 was 51.64 inches. Figures 1-7 and 1-8 show the 2008 monthly and the 60-year annual precipitation data. Snowfall for the 2007–2008 winter season was 11.5 inches, well below the 31.2 inches of average yearly snowfall for Long Island, and about 10 times less than the record high snowfall of 90.8 inches, set in the 1995–1996 snow season.



frequency; that is, how often the wind came from that direction. The wind direction was measured at heights of 10 and 88 meters. This diagram indicates that the predominant wind direction was from the south southwest at the 10-m level and west northwest at the 88-m level. Due to instrument failure, the 2008 Wind Rose data is limited to January through August 2008.

Figure 1-4. BNL Wind Rose (2008).




June Figure 1-6. BNL Annual Mean Temperature Trend (60 Years).

July

Sep

Oct

Dec






Figure 1-8. BNL 2008 Annual Precipitation Trend (60 Years).

Jan

Feb

Mar

Apr

May

1.8 NATURAL RESOURCES

The Laboratory is located in the oak/chestnut forest region of the Coastal Plain and constitutes about 5 percent of the 100,000-acre New York State-designated region on Long Island known as the Central Pine Barrens. The section of the Peconic River running through BNL is designated as "scenic" under the New York State Wild, Scenic, and Recreational River System Act of 1972. Due to the general topography and porous soil, the land is very well drained and there is little surface runoff or open standing water. However, depressions form numerous small, pocket wetlands with standing water on a seasonal basis (vernal pools), and there are six regulated wetlands on site. Thus, a mosaic of wet and dry areas correlates with variations in topography and depth to the water table.

Vegetation on site is in various stages of succession, which reflects a history of disturbances to the area. For example, when Camp Upton was constructed in 1917, the site was entirely cleared of its native pines and oaks. Although portions of the site were replanted in the 1930s, portions were cleared again in 1940 when Camp Upton was reactivated by the U.S. Army. Other past disturbances include fire, local flooding, and draining. Current operations minimize disturbances to the more natural areas of the site.

More than 230 plant species have been identified at the Laboratory, including two species that are threatened in New York State and two that are classified as rare. Fifteen animal species identified on site include a number that are protected in New York State, as well as species common to mixed hardwood forests and open grassland habitats. At least 85 species of birds have been observed nesting on site, and more than 200 transitory bird species have been documented visiting the site. (BNL is located within the Atlantic Flyway, with scrub/shrub habitats that offer food and rest to migratory songbirds.) Permanently flooded retention basins and other watercourses support amphibians and aquatic reptiles. Thirteen amphibian and 12 reptile species have been identified at BNL. Recent ecological studies have confirmed 26 breeding sites for the New York State endangered eastern tiger salamander in ponds and recharge basins. Ten

species of fish have been identified as endemic to the site, including the banded sunfish and the swamp darter, both of which are threatened in New York State. Two types of butterflies that are protected in New York State are believed to breed on site due to the presence of their preferred habitat and host plants, and a New York State threatened damselfly was found on site in 2005. To eliminate or minimize any negative effects that Laboratory operations might cause to these species, precautions are in place to protect the on-site habitats and natural resources.

In November 2000, DOE established the Upton Ecological and Research Reserve at BNL. The 530-acre Upton Reserve (10 percent of the Laboratory's property) is on the eastern portion of the site, in the Core Preservation Area of the Central Pine Barrens. The Upton Reserve creates a unique ecosystem of forests and wetlands that provides habitats for plants, mammals, birds, reptiles, and amphibians. From 2000 to 2004, funding provided by DOE under an Inter-Agency Agreement between DOE and the U.S. Fish & Wildlife Services was used to conduct resource management programs for the conservation, enhancement, and restoration of wildlife and habitat in the reserve. In 2005, management was transitioned to the Foundation for Ecological Research in the Northeast (FERN). Management of the Upton Reserve falls within the scope of BNL's Natural Resource Management Plan, and the area will continue to be managed for its key ecological values and as an area for ecological research. Additional information regarding the Upton Reserve and the Laboratory's natural resources can be found in Chapter 6 of this report.

1.9 CULTURAL RESOURCES

The Laboratory is responsible for ensuring compliance with historic preservation requirements. BNL's Cultural Resource Management Plan was developed to identify, assess, and document the Laboratory's historic and cultural resources. These resources include World War I trenches; Civilian Conservation Corps features; World War II buildings; and historic structures, programs, and discoveries associated with highenergy physics, research reactors, and other

science conducted at BNL. The Laboratory currently has three facilities classified as eligible for listing on the National Register of Historic Places: the Brookhaven Graphite Research Reactor complex, the High Flux Beam Reactor complex, and the World War I training trenches associated with Camp Upton. Further information can be found in Chapter 6.

REFERENCES AND BIBLIOGRAPHY

BNL. 2003. Natural Resource Management Plan for Brookhaven National Laboratory. Brookhaven National Laboratory, Upton, NY.

BNL. 2005. Cultural Resource Management Plan for Brookhaven National Laboratory. Brookhaven National Laboratory, Upton, NY.

DOE Order 231.1.A. 2003. *Environment, Safety and Health Reporting*. U.S. Department of Energy, Washington, DC.

Geraghty and Miller, Inc. 1996. Regional Groundwater Model, Brookhaven National Laboratory, Upton, New York. A Report to Brookhaven National Laboratory. November 1996. Kamer, Pearl M. 2006. *The Economic Impact of Brookhaven National Laboratory on the New York State.*Suffolk County Planning Commission, Suffolk County Department of Planning. October 2005.

Koppelman, L.E. 1978. The Long Island Comprehensive Waste Treatment Management Plan (Long Island 208 Study), Vol. I and II. Long Island Regional Planning Board, Hauppauge, NY. July 1978.

Nagle, C.M. 1975. *Climatology of Brookhaven National Laboratory:* 1949–1973. BNL-50466. Brookhaven National Laboratory, Upton, NY. November 1975.

Nagle, C.M. 1978. *Climatology of Brookhaven National Laboratory:* 1974–1977. BNL-50857. Brookhaven National Laboratory, Upton, NY. May 1978.

NYCRR. Title 27. *Wild, Scenic, and Recreational River Systems Act.* Article 15 and subsequent updates. New York State Department of Environmental Conservation, Albany, NY.

Scorca, M.P., W.R. Dorsch, and D.E. Paquette. 1999. Stratigraphy and Hydrologic Conditions at the Brookhaven National Laboratory and Vicinity, Suffolk County, New York, 1994–97. U.S. Geological Survey Water Resources Investigations Report 99-4086. 55 pp.

Warren, M.A., W. deLaguna, and N.J. Lusczynski.1968. Hydrology of Brookhaven National Laboratory and Vicinity, Suffolk County, New York. U.S. Geological Survey Bulletin, 1156-C.



