Polarised DIS Experiment

Lecture II

Gerhard Mallot



Plan

- QCD analysis of $g_1(x,Q^2)$ and PDFs
- status of g_2
- semi-inclusive scattering and flavour decomposition
- transversity h_1 and Collins asymmetry
- the hunt for ΔG
  - *high-*p_T hadron pairs
  - *open charm* production
- summary

Status of g_1



- Wealth of data g_1 data for p, n and d
- Data taken at different Q^2
- Only weak Q^2-dependence in overlap region
- Interesting data from JLAB at large x for neutron: $g_1^n > 0$



Q^2 Evolution



splitting functions

non-singlet quark distribution

$$\Delta q^{\rm ns}(x,Q^2) = \sum_{i=1}^{n_f}\left(\frac{e_i^2}{\langle e^2\rangle} - 1\right)\Delta q_i(x,Q^2)$$

singlet quark distribution

$$\Delta\Sigma(x,Q^2) = \sum^{n_f}\Delta q(x,Q^2)$$

$$\frac{\rm d}{{\rm d}\ln Q^2}\Delta q^{\rm ns} = \frac{\alpha_s}{2\pi}\Delta P_{qq}^{\rm ns}\otimes\Delta q^{\rm ns}$$

$$\frac{\rm d}{{\rm d}\ln Q^2}\begin{pmatrix}\Delta\Sigma\\ \Delta g\end{pmatrix} = \frac{\alpha_s}{2\pi}\begin{pmatrix}\Delta P_{qq}^{\rm s} & 2n_f\Delta P_{qg}^{\rm s}\\ \Delta P_{gq}^{\rm s} & \Delta P_{gg}^{\rm s}\end{pmatrix}\otimes\begin{pmatrix}\Delta\Sigma\\ \Delta g\end{pmatrix}$$

DGLAP evolution

Q^2 Evolution

- non-singlet decouples from gluon evolution, moments Q^2-independent, like $\Delta u - \Delta d = g_a$
- Evolution of singlet and gluon coupled, moments $\Delta\Sigma$ and ΔG evolve with Q^2
- In principle ΔG can be determined from the Q^2 evolution of $g_1(x,Q^2)$
- Need reasonable range Q^2 at fixed x
- We have some Q^2 range, but not much, important large Q^2 at small x

$g_1(x,Q^2)$



Looks quite nice, but...

$g_1(x,Q^2)$ $F_2(x,Q^2)$


NLO
g_1^p
x=0.008 (×2048)
x=0.015 (×1024)
x=0.025 (×512)
x=0.035 (×256)
x=0.05 (×128)
x=0.08 (×64)
x=0.125 (×32)
x=0.175 (×16)
x=0.25 (×8)
x=0.35 (×4)
x=0.5 (×2)
x=0.75
E155
E143
SMC
EMC
HERMES
Q^2 (GeV^2)


$F_2+c_i(x)$
SLAC
NMC
BCDMS
H1 96-97 preliminary
H1 94-97 e$^+$p
NLO QCD Fit
$c_i(x)=0.6\cdot(i(x)-0.4)$
Q^2 /GeV^2

NLO QCD Fits

- choose scheme, usually $\overline{\text{MS}}$
- choose start value for evolution, Q^2_0
- choose parametrisations for

$$\Delta\Sigma,\ \Delta G,\ \Delta q^{ns}(x,Q^2_0)$$

- fit parameters of these parametrisations using the DLGAP equations (NLO)
- many groups, an example...
  - AAC03 Kumano DIS2004

AAC03 Fit



error of AAC00

valence well fixed
gluon undetermined
sea quarks negative

AAC03 with zero gluon at Q^2_0

χ^2/d.o.f. = 0.915



error with free gluon

Comparsion of PDF Fits



BB Fit



slightly positive gluon pol. at Q^2=5 GeV^2

unpolarised gluon

Blümlein & Böttcher

First moments

$Q^2 = 1\,\mathrm{GeV}^2$

	Δg	$\Delta\Sigma$
AAC03	0.499 ± 1.268	0.213 ± 0.138
GRSV01	0.420	0.204
LSS	0.680	0.210
BB	1.026	0.138

Kumano DIS2004

- GRSV01 [Phys. Rev. D63 (2001) 094005]
- LSS01 [Eur.Phys.J. C23 (2002) 479]
- BB02 [Nucl. Phys. B636 (2002) 225]

g_2

$$
\begin{aligned}
g_2(x,Q^2) &= g_2^{\mathrm{WW}} + \bar{g}_2(x,Q^2) \\
g_2^{\mathrm{WW}}(x,Q^2) &= -g_1(x,Q^2) + \int_x^1 \frac{g_1(y,Q^2)}{y}\,\mathrm{d}y
\end{aligned}
$$

- twist-3 term $\bar{g}_2$, matrix element d_2

$$
d_2 = 3\int_0^1 x^2\,\bar{g}_2(x,Q^2)\,\mathrm{d}x
$$

g_2

- twist-2 term: g_2^{WW}
- bag model calculations



Neutron g_2 from JLAB



Fragmentation Function $D_f^h(z,Q^2)$



D_q^h from quark q into hadron h

$z = \frac{E_h}{\nu}$ energy fraction carried by h



$$D_u^{\pi^+} \underset{\text{CC}}{=} D_{\bar{u}}^{\pi^-} \underset{\text{IS}}{=} D_{\bar{d}}^{\pi^+} \underset{\text{CC}}{=} D_d^{\pi^-}$$

$$D_d^{\pi^+} = D_{\bar{d}}^{\pi^-} = D_{\bar{u}}^{\pi^+} = D_u^{\pi^-}$$

$$\frac{1}{\sigma_0}\frac{d\sigma^h}{dz} = \frac{\sum_f e_f^2\, q_f(x,Q^2)\cdot D_f^h(z,Q^2)}{\sum_f e_f^2\, q_f(x,Q^2)}$$

Semi-inclusive DIS



final hadron "remembers" flavour of initially struck!

$$A_1^h(x,Q^2) = \frac{\int \mathrm{d}z \sum_f e_f^2 \Delta q_f(x,Q^2) \cdot D_f^h(z,Q^2)}{\int \mathrm{d}z \sum_f e_f^2 \; q_f(x,Q^2) \cdot D_f^h(z,Q^2)}$$

Flavour separated polarisation



Asymmetries can in LO be related to Δq by

$$\vec{\mathcal{A}} = \mathcal{P}\vec{\mathcal{Q}}$$

where

$$\vec{\mathcal{A}} = (A^h_{1,t}, \ldots)$$
$$\vec{\mathcal{Q}} = (\Delta q_f, \ldots)$$

$$\mathcal{P}^h_f = \frac{e_f^2 q_f(x) \int dz D^h_f}{\sum_i e_i^2 q_i(x) \int dz D^h_i(z)}$$

more data



SMC and COMPASS have more small *x* data.
Will help to clarify Δ s

Projection for Δs (2002 data)



Transversity h_1





Collins asymmetries 2002



Deuteron

Hermes Collins asymmetry



Δ G from high-p_T hadron pairs

- Contributions: a) LO b) QCD Compton c) PGF



$$A_{\mathrm{LL}}^{\ell N} \simeq \langle \hat{a}_{\mathrm{LL}}^{\gamma g \to q g} \rangle \frac{\Delta q}{q} + \langle \hat{a}_{\mathrm{LL}}^{\gamma g \to q \bar{q}} \rangle \frac{\Delta g}{g}$$

Hermes high-p_T pairs





Hermes high-p_T pairs

- Hermes conclusion:
  $\Delta g/g = 0.41 \pm 0.18(\text{stat}) \pm 0.03(\text{syst exp})$
  $\langle x_g \rangle = 0.17;\ \langle p_T^2 \rangle = 2.1\ \text{GeV}^2$

$\int_{0.06}^{0.028} \frac{\Delta g}{g} g(x) \mathrm{d}x \approx 0.6$



h^+h^-



$\pi^+\pi^-$

K^+K^-



SMC high-p_T anaylysis

- Hermes: Q^2 not reconstructed for high-p_T
- SMC: cut on $Q^2 > 1$ GeV^2, theoretical description more sound, limited statistics



Selection	$\Delta G/G \pm \delta(\Delta G/G)_{stat}$	$\langle x_g \rangle_{genPGF}$
$\sum p_T^2 > 2.5$ GeV^2	-0.07 ± 0.40	0.09
NN >0.26	-0.20 ± 0.28	0.07

COMPASS high-p_T hadron pairs



2002 data only, all Q^2, *75000 events*

- High p_T cut
  - $p_{T1}, p_{T2} > 0.7$ GeV/c
  - $p_{T1}^2 + p_{T2}^2 > 2.5$ GeV^2/c^2
- Vector mesons are removed
  - $m(h_1 h_2) > 1.5$ GeV/c^2
- Products of the target fragmentation are removed
  - $x_F > 0.1$
  - $z > 0.1$

$$\left(\frac{A_\parallel}{D}\right)^{\mu d \to hh} = -0.065 \pm 0.036(stat.) \pm 0.01(syst.)$$

Open charm at COMPASS

- Photon-gluon fusion: 1.2 D^0 per PGF $c\bar{c}$ event



$$D^0 \Rightarrow K^- \pi^+$$ BR 4 %

or: $$D^{*+} \Rightarrow D^0 \pi_s^+$$

$$\Rightarrow K^- \pi^+ \pi_s$$

D^0 from D^* about 20 %

Open charm, cuts

- most of 2002 data, prel. RICH and tracking
- $z_D > 0.2$ (background reduction)
- $|\cos(\theta^*)| < 0.85$ (background reduction)
- $10 < p_K < 35$ GeV/c (Rich πK sep.)
- define:

$$\Delta M_{K\pi\pi} = M_{K\pi\pi_s} - (M_{K\pi} + M_{\pi_s})$$

$$M(D^*) - [M(D^0) + M(\pi)]$$

$D^{*+} \rightarrow D^0 \pi_s^+$ tagging

$$D^* \rightarrow D^0 \pi_s$$
$$\rightarrow K\pi\pi_s \quad (+\,cc)$$

$$M_{K\pi} - M_{D^0}$$

$$\Delta M_{K\pi\pi} = M_{K\pi\pi_s} - (M_{K\pi} + M_{\pi_s})$$



Choose:
$3.1 < \Delta M_{K\pi\pi} < 9.1$ MeV

$D^{*+} \rightarrow D^0 \pi_s^+$ tagging





Projected precision for Δ G/G

- 2002 – 2004 data:
- from high-p_T: *(all Q^2)* $\delta(\Delta G/G) = 0.05$
- from high-p_T: *($Q^2 > 1$)* $\delta(\Delta G/G) = 0.16$
- from open charm: $\delta(\Delta G/G) = 0.24$

No systematic error included yet
(not too big, apart from possible theoretical uncertainties)

Summary

- Lots to come from:
  - Hermes
  - COMPASS
  - RHIC

- It seems SPIN goes down very far ...



"This could be the discovery of the century. Depending, of course, on how far down it goes."