

# Sensitivity to New Physics in Parity Violating Asymmetries at RHIC

Jean-Marc Virey<sup>1</sup>

Centre de Physique Théorique

CNRS Luminy, Case 907

13288 Marseille cedex 09, France

We have presented the sensitivity of the RHIC Spin experiment to some new physics contributions, from the analysis of the Parity Violating (PV) asymmetry  $A_{LL}^{PV}$  ( $\equiv d\sigma^{--} - d\sigma^{++}/d\sigma^{--} + d\sigma^{++}$ ) defined for jet production ( $d\sigma \equiv d\sigma_{1-jet}$ ).

In the first part, we have recalled the sensitivity to the presence of some new quark-quark Contact Interactions (CI). Such CI could represent some effects of quark compositeness. At RHIC, with the conventional parameters of the experiment, we will be sensitive to some energy scales of  $\Lambda \sim 3-4 \text{ TeV}$ , if PV is maximal. It appears that the RHIC is perfectly competitive with the TEVATRON for the discovery of CI, however, we have to note that the integrated luminosity is a key parameter for the polarized analysis. The study of  $A_{LL}^{PV}$  could give some unique informations on the chiral structure of the new interaction.

The second part was devoted to a review of the different theoretical motivations for the presence of a "leptophobic"  $Z'$  of relatively low mass, that we briefly report now :

1) Leptophobia corresponds to some new interactions belonging to the "pure" quark sector. 2) Leptophobic  $Z'$  appears naturally in several models derived from string theory. Within these models, the absence of  $Z'$  leptonic couplings appears in several ways : from the classification of the fermions in the different representations of the gauge group, or from kinetic mixing between two abelian gauge groups which produces a mixing between the charges of the fermions under these two  $U(1)$  factors. 3) Some theoretically consistent leptophobic  $Z'$  bosons have been constructed also for non-supersymmetric models. 4) In the framework of supersymmetric models with an additional abelian gauge factor  $U(1)'$ , it has been shown that the  $Z'$  boson could appear with a relatively low mass ( $M_Z \leq M_{Z'} \leq 1 \text{ TeV}$ ) and with a mixing angle with the  $Z$  close to zero. In addition, these classes of models have several appealing features for soft supersymmetry breaking and for cosmology. 4) The present experimental constraints on these models are rather weak. Several classes of models are absolutely unconstrained. 5) All these theoretically motivated models exhibit, in general, some PV couplings to up and down quarks, which could induce some interesting effects on  $A_{LL}^{PV}$  measured in  $p-p$  or/and  $n-n$  collisions at RHIC. The main results are presented in the following transparencies.

The last section concerned the measurement of  $A_{LL}^{PV}$  in  $p-n$  collisions, which could detect the presence of a new charged boson  $W'$  unconstrained by present experimental data.

---

<sup>1</sup>email: virey@cpt.univ-mrs.fr

2) Properties for up /  $A_{LL}^{PV}$  in p-p collisions

In general, the lepto-phobic  $Z'$  violates  
Parity in the up quark sector!

- "Flipped SU(5)": Maximal PV!

$$C_L^u' = \pm \frac{1}{2} \quad C_R^u' = 0 \quad (\Rightarrow \text{Left})$$

- "S-kinetic":  $C_R^u' = -2 C_L^u' = \frac{1}{3}$  ( $\Rightarrow$  Right)

- The constraint of all the "classes" of models is:


$$C_L^u' \neq C_R^u'$$

i.e.: axial couplings ( $C_L^u' = -C_R^u'$ ) authorized  $\Rightarrow$  No effect  
on  $A_{LL}^{PV}$ !

Except this case, for  $M_{Z'} \lesssim 300 - 400$  GeV,  
these models could be detected at RHIC  
from  $A_{LL}^{PV}$  in p-p collisions!

error bars:  $\sqrt{s} = 500 \text{ GeV}$   
 $L = 800 \text{ pb}^{-1}$   
 $\Delta E_T = 10 \text{ GeV}$   
 $\Delta y = \pm$

$$\frac{\Delta F_{\text{systematic}}}{F} = 10\%$$



### 3°) Properties for down / $F_{LL}^{PV}$ in n-n collisions

Isospin symmetry  $\Rightarrow F_{LL}^{PV}$  (n-n) rest d

What could we expect from these  $Z'$ ?

$\rightarrow$  Models with Parity Conservation (for d)

( $\Rightarrow$  No effect on  $F_{LL}^{PV}$  (n-n))

- "flipped  $SU(5)$ " and "g-kinetic" are PC :  $C_L^d = \pm C_R^d$

- PC for d is a general property of the minimal 2 doublets Higgs model

$$h_u Q u^c H_u + h_d Q d^c H_d + h_e L e^c H_d \in \frac{\mathbb{Z}_2}{W}$$

$$\text{Leptophobia} \equiv Q'(L) = 0 = Q'(e^c)$$

$$\Rightarrow Q'(H_d) = 0 \Rightarrow Q'(Q) + Q'(d^c) = 0$$

$$\Delta \left\{ \begin{array}{l} Q'(f): \text{charge} \\ \text{of } f \text{ under } U(1) \\ C_L^d = Q'(f) \\ C_R^d = -Q'(f^c) \end{array} \right.$$

$$\Rightarrow C_L^d = C_R^d \quad \text{i.e. Vectorial couplings}$$

→ Non-minimal Higgs Models

- 3 Higgs doublets (Georgi-Glashow)  
Agashe et al.
- 2 Higgs doublets + d mass terms (in  $W$ ) from higher dimensional operators: (Cvetic et al)  
Natural in string models (Faraggi: PLB 377 (96) 143)

They do not impose PC for d, conversely

we often have:  $C_L^{d'} \neq C_R^{d'}$

(i.e. PV for d except for the axial case)

- Interesting model: "Minor Z" (Caraveglio & Ross  
PLB 346 (95) 159)

$$\frac{C_L^q}{C_R^q} = \frac{C_R^{q'}}{C_L^{q'}} \Rightarrow \begin{cases} C_R^u \approx -2 C_L^u \\ C_R^d \approx -5 C_L^d \end{cases}$$

We expect strong PV for d in this model!

No curves for  $F_{LL}^{PV}(n,n)$ , sorry! Available soon.)

- Conclusions:
- $F_{LL}^{PV}(n,n)$  is sensitive to NP if PV/a
  - It could constraint the scalar sector of the new theory!

## 4) Experimental Constraints

298

### • Direct:

Come from  $\nabla_{\text{set}}^{\text{unpol}}$

JF2 : fabrics for  $K = \frac{g_{Z'}}{g_Z} = 1$ :  $100 < M_{Z'} < 250$  GeV

DØ : " "  $365 < M_{Z'} < 615$

CDF : no-constraints

→ limits obtained for SM-like  $Z'$  (i.e. same coupling)

→ limits highly model dependent

→ Disappear for  $K \lesssim 0.95$

(Re:  $K_{\text{GUT}} \sim 0.7$  !)

need  
 $\left\{ \begin{array}{l} K \leq 0.9 \text{ for } b\phi \\ \text{high masses, bonus} \\ \text{inevitable for RHIC} \end{array} \right.$

" $g$ -kinetic" absolutely unconstrained (and favorable)

→ Large uncertainties (exp. + tho.) for  $\nabla_{\text{set}}^{\text{unpol}}$  !

### • Indirect

From  $Z$ -pole (LEP/sec) + low energy PV  $\rightarrow \Theta_{Z-Z'} \lesssim \frac{10^{-2}}{10^{-3}}$

In leptophobic  $Z'$  models, in general,  $\Theta_{Z-Z'}$  is naturally  $\sim 0$   
 Conclusion:

Low mass leptophobic  $Z'$  are not in contradiction with experiments

### III $W'$ at RHIC

(P.Tanii & S.M. PLB 404 (37) 307)

$A_{LL}^{PV}$  is sensitive to a  $W'$  in p-n collisions

(result from Isospin symmetry)

→ ST effects: pp: Z-g ~ 70% W-g ~ 30%

p-n: Z-g ~ 2% W-g ~ 98% !!

→  $W_R$  from LRM

Limits on  $M_{W_R}$  are highly model dependent.

Relevant parameters: (Langacker & Umasankar PRD 40 (23))

$$V_{CKM}^R, \gamma_R, K = \frac{g_R}{g_L}, \xi, \omega \text{ (CP phases)} \quad \theta_{W_L - W_R}$$

Case the weakest constraint:  $V_{CKM}^R \approx 1$ ,  $\gamma_R$  heavy Dirac  
(natural! less (!))

$A_{LL}^{PV}$  in p-n collisions is sensitive

for this model if  $M_{W'_R} \lesssim 400$  GeV

( $K = \alpha \beta$ ))