Parity-violating Asymmetries in $W^{\pm}$ Production with STAR

L.C. Bland
Dept. of Physics, Indiana University, Bloomington, IN 47405 USA

In addition to its crucial role in accurately determining the polarized gluon structure function for the proton via measurements of the longitudinal spin correlation coefficient (A_{LL}) in $pp \rightarrow \gamma$ + jet reactions at $\sqrt{s}$ = 200 GeV, the proposed end-cap electromagnetic calorimeter (EEMC) upgrade to STAR also allows important determinations of flavor-specific quark and antiquark polarizations through measurements of the $e^{\pm}(\nu)$ decay of $W^{\pm}$ bosons produced in collisions of longitudinally polarized protons at $\sqrt{s}$ = 500 GeV. The ability to relate the measured parity-violating single-spin asymmetry (A_L) directly to the quark or antiquark polarization is greatest when the e^+ from W^+ decays are detected at large pseudorapidity, $1 < \eta < 2$, corresponding to the acceptance of the proposed EEMC.

Fig. 1 shows the expected η distribution of $e^{\pm}$ from $W^{\pm}$ decays for pp collisions at $\sqrt{s}$ = 500 GeV as generated by PYTHIA [1]. Simple expressions for the $W^{\pm}$ production cross section from ref. [2] provide substantial insight. In particular, when the $W^{\pm}$ are produced at large rapidity, y_W, then in leading-order, we expect the most asymmetric collisions in the initial state corresponding to the partonic momentum fractions, $x_1 >> x_2$. For such asymmetric collisions, we expect (Fig. 2) a direct relationship between the measured A_L and the $u(d)$ and $\bar{d}(\bar{u})$ polarizations.

To test this idea, a simulation model for the $\vec{p}p \rightarrow W^{\pm}X \rightarrow e^{\pm}(\nu)X$ reaction at $\sqrt{s}$ = 500 GeV was developed. Events were generated for $W^{\pm}$ production, excluding the proton polarization, using PYTHIA. The partonic kinematics for the event were then used to evaluate the expected A_L for each of the two beam longitudinally polarized beams (assuming P_{beam} = 0.7) using the expressions from ref. [2] and polarized structure functions from ref. [3]. The detection of e^+ from W^+ decays was modeled by including only the acceptance of the STAR TPC and barrel and end-cap EMC. The events were subjected to particle identification cuts to distinguish $e^{\pm}$ from the most important background arising from the prolific production of hadrons ($h^{\pm} \equiv$ protons(anti-protons), $K^{\pm}$ and $\pi^{\pm}$) with large transverse momentum, p_T > 10 GeV/c. Other backgrounds such as Z^0 decay, with only a single $e^{\pm}$ detected and single electrons from the Dalitz decay of high-p_T π^0 have been considered and found to be negligible. Clean distinction between $e^{\pm}$ and $h^{\pm}$ relies on (1) *isolation cuts* applied to the candidate $e^{\pm}$, (2) the much smaller response of the EMC to incident $h^{\pm}$ relative to $e^{\pm}$, and (3) rejecting events that have an accompanying jet with p_T > 5 GeV/c in the azimuthal angle range, $|\pi - (\phi_{jet} - \phi_h)| < 1$ radian. The first two of these cuts have been discussed before [4]; the third cut further reduces the hadronic background by an additional order of magnitude. The end result is that STAR can cleanly detect $e^{\pm}$ from $W^{\pm}$ decays.

Fig. 3 shows the expected A_L for the four independent quantities that can be measured (asymmetries from either beam longitudinally polarized, and detection of either e^+ or e^-). To examine if the measured asymmetries are directly related to quark and antiquark polarizations, the identities of the partons involved in the $W^{\pm}$ production were examined. Figs. 4 and 5 show the x distribution of the initial-state partons, available from PYTHIA for two different ranges of $e^{\pm}$ pseudorapidity. It's clear that only when the e^+ are detected at large η that A_L can be directly related to $u(d)$ and $\bar{d}(\bar{u})$ polarizations.

The next step to take with the simulation model is to attempt the reconstruction of the initial-state $u(d)$ and $\bar{d}(\bar{u})$ momentum fractions (x) assuming collinear collisions. As well, a more accurate simulation of the response of the STAR detector is essential to quantitatively establish the accuracy of the determination of the $u(d)$ and $\bar{d}(\bar{u})$ polarizations.

[1] PYTHIA 5.7 / JETSET 7.4 T. Sjöstrand, Comp. Phys. Comm. **82** (1994) 74.
[2] C. Bourrely, J. Soffer, F.M. Renard and P. Taxil, Phys. Rept. **177** (1989) 319.
[3] T. Gehrmann and W.J. Stirling, Phys. Rev. **D53** (1996) 6100.
[4] A.A. Derevschikov, V.L. Rykov, K.E. Shestermanov and A. Yokosawa in *Proc. 11th Intl. Symp. on High Energy Spin Physics*, Bloomington, 1994, eds. Kenneth J. Heller and Sandra L. Smith, AIP Conf. Proc. No. 343 (1995) p. 472.



$pp \to W^{\pm} X \to e^{\pm}(\nu) X$

$800\ pb^{-1}$

$\sqrt{s} = 500\ GeV$

(PYTHIA)

$$\frac{d\sigma}{dy}(W^+) = G_F \frac{\sqrt{2}}{3} \pi \tau \left[\underset{d}{u(x_a, M_W^2)}\ \underset{\bar{u}}{\bar{d}(x_b, M_W^2)} + (a \leftrightarrow b) \right]$$

W^-

$$\tau = \frac{M_W^2}{s_{pp}} \qquad x_1 = \sqrt{\tau}\, e^{-y_W} \qquad x_2 = \sqrt{\tau}\, e^{+y_W}$$

If $y_W \sim y_e$ then endcap $\Rightarrow x_2 \gg x_1$

$\Rightarrow$ Interpret A_L in terms of $\frac{\Delta q}{q}$ & $\frac{\Delta \bar{q}}{\bar{q}}$?

Fig. 1

Bourrely & Soffer



$u\bar{d} \to W^+$

$d\bar{u} \to W^-$

A_L measurements give sensitivity to

$$\Delta q = q_\uparrow - q_\downarrow$$

$$\Delta \bar{q} = \bar{q}_\uparrow - \bar{q}_\downarrow$$

$\{W^+\}$ $\{W^-\}$

"a" polarized

$$\frac{\Delta u(x_a)\,\bar{d}(x_b) - \Delta\bar{d}(x_a)\,\cancel{u(x_b)}}{u(x_a)\,\bar{d}(x_b) + \bar{d}(x_a)\,\cancel{u(x_b)}} \quad \text{small}$$

$$\xrightarrow[x_a \gg x_b]{} \frac{\Delta u(x_a)}{u(x_a)}$$

$$\frac{\Delta d(x_a)\,\bar{u}(x_b) - \Delta\bar{u}(x_a)\,\cancel{d(x_b)}}{d(x_a)\,\bar{u}(x_b) + \bar{u}(x_a)\,\cancel{d(x_b)}} \quad \text{small}$$

$$\xrightarrow[x_a \gg x_b]{} \frac{\Delta d(x_a)}{d(x_a)}$$

"b" polarized

$$\frac{u(x_a)\,\Delta\bar{d}(x_b) - \bar{d}(x_a)\,\Delta u(x_b)}{u(x_a)\,\bar{d}(x_b) + \bar{d}(x_a)\,\cancel{u(x_b)}} \quad \text{small}$$

$$\xrightarrow[x_a \gg x_b]{} \frac{\Delta\bar{d}(x_b)}{\bar{d}(x_b)} - \frac{\bar{d}(x_a)\,\Delta u(x_b)}{u(x_a)\,\bar{d}(x_b)}$$

$$\frac{d(x_a)\,\Delta\bar{u}(x_b) - \bar{u}(x_a)\,\Delta d(x_b)}{d(x_a)\,\bar{u}(x_b) + \bar{u}(x_a)\,\cancel{d(x_b)}} \quad \text{small}$$

$$\xrightarrow[x_a \gg x_b]{} \frac{\Delta\bar{u}(x_b)}{\bar{u}(x_b)} - \frac{\bar{u}(x_a)\,\Delta d(x_b)}{d(x_a)\,\bar{u}(x_b)}$$

Fig. 2

$\vec{p}p \to W^{\pm}X \to e^{\pm}(\nu)X$ $\sqrt{s} = 500$ GeV

$800\ pb^{-1}$ $P_{beam} = 0.7$

beam a polarized | beam b polarized

98/04/29 06.57



Do A_L values determine $\frac{\Delta q}{q}$, $\frac{\Delta \bar{q}}{\bar{q}}$?

→ Look in different η ranges

Fig. 3

$W^+ \to e^+ \nu$

beam a | beam b

98/04/25 21.58



parton composition of beam

$\langle A_L \rangle \sim +0.3$ $\quad x \quad$ $\langle A_L \rangle \sim +0.2$

$$-1 < \eta_{e^+} < 0$$

$\Rightarrow A_L$ not directly related to $\frac{\Delta u}{u}$ or $\frac{\Delta \bar{d}}{\bar{d}}$

Fig. 4

$W^- \to e^- \bar{\nu}$

98/04/25 21.58

beam a | beam b



parton composition of beam

$\langle A_L \rangle \sim -0.3$ x $\langle A_L \rangle \sim 0$

(endcap EMC)

$1 < \eta_{e^-} < 2$

$\Rightarrow$ A_L is directly related to $\frac{\Delta d}{d}$ (beam a) & $\frac{\Delta \bar{u}}{\bar{u}}$ (beam b)

Fig. 5