RHIC
Workshop
'98

PROBING THE NUCLEON'S TRANSVERSITY VIA TWO-PION PRODUCTION AT RHIC

Jian Tang

MIT

Collaborators: R. L. Jaffe, Xuemin Jin

The result

$$\vec{P}_\perp p \longrightarrow \pi^+\pi^- X$$





At partonic level



observable: $\vec{\pi}^+ \times \vec{\pi}^- \cdot \vec{S}_\perp$

Nucleon's polarization

$\propto \cos\phi$

Pion momentum

collins angle

The result

$$\vec{P}_\perp p \longrightarrow \pi^+\pi^- X$$





At partonic level



Observable: $\vec{\pi}^+ \times \vec{\pi}^- \cdot \vec{S}_\perp$

Nucleon's polarization

$\propto \cos\phi$

Pion momentum

collins angle

An asymmetry

$$A_{\perp T} \equiv \frac{d\sigma_\uparrow - d\sigma_\downarrow}{d\sigma_\uparrow + d\sigma_\downarrow}$$

$$= -\frac{\sqrt{6}}{4}\pi \overbrace{\sin\delta_0 \sin\delta_1 \sin(\delta_0 - \delta_1)}^{\text{FSI phases}} \overbrace{\cos\phi}^{\text{Collins angle} \propto \vec{\pi}^+ \times \vec{\pi}^- \cdot \vec{S}_\perp}$$

$$\times \underbrace{\frac{\delta\hat{\sigma}^{qq}_{qq}}{\hat{\sigma}^{qq}_{qq}}}_{\text{Hard process asymmetry } \hat{a}_{\perp T} \text{ for } qq \to qq, \text{ which is dominant here.}} \frac{\sum_a \delta q^a \, \delta\hat{q}^a_I}{\sum_a q^a \left(\sin^2\delta_0 \, \hat{q}^a_0 + \sin^2\delta_1 \, \hat{q}^a_1\right)}$$

δq^a : Transversity

$\delta\hat{q}^a_I$: unknown interference fragmentation function

q^a : unpolarized quark distribution

$\hat{q}^a_0$: "σ" fragmentation function

$\hat{q}^a_1$: "ρ" fragmentation "

Suppressed dependences

$\delta q \to \delta q(x, Q^2)$, $\quad \delta\hat{q}_I \to \delta\hat{q}_I(z, Q^2, m^2)$

$q \to q(x, Q^2)$, $\quad \hat{q}_0, \hat{q}_1 \to \hat{q}_0(z, Q^2), \hat{q}_1(z, Q^2)$

$\delta_{0,1} \to \delta_{0,1}(m^2)$

<u>Advantages</u>:

- Twist 2
- Abundant pions
- Crucial FSI are known (δ_0 δ_1)

An asymmetry

$$A_{\perp T} \equiv \frac{d\sigma_\uparrow - d\sigma_\downarrow}{d\sigma_\uparrow + d\sigma_\downarrow}$$

$$= -\frac{\sqrt{6}}{4}\pi \overbrace{\sin\delta_0 \sin\delta_1 \sin(\delta_0 - \delta_1)}^{\text{FSI phases}} \overbrace{\cos\phi}^{\text{Collins angle} \propto \vec{\pi}^+ \times \vec{\pi}^- \cdot \vec{S}_\perp}$$

$$\times \underbrace{\frac{\delta\hat{\sigma}^{qq}_{qq}}{\hat{\sigma}^{qq}_{qq}}}_{\text{Hard process asymmetry } \hat{a}_{\perp T} \text{ for } qq \to qq, \text{ which is dominant here.}} \frac{\sum_a \delta q^a \, \delta\hat{q}^a_I}{\sum_a q^a \left(\sin^2\delta_0 \, \hat{q}^a_0 + \sin^2\delta_1 \, \hat{q}^a_1\right)}$$

δq^a : Transversity

$\delta\hat{q}^a_I$: unknown interference fragmentation function

q^a : unpolarized quark distribution

$\hat{q}^a_0$: "σ" fragmentation function

$\hat{q}^a_1$: "ρ" fragmentation "

Suppressed dependences

$\delta q \to \delta q(x, Q^2)$, $\quad \delta\hat{q}_I \to \delta\hat{q}_I(z, Q^2, m^2)$

$q \to q(x, Q^2)$, $\quad \hat{q}_0, \hat{q}_1 \to \hat{q}_0(z, Q^2), \hat{q}_1(z, Q^2)$

$\delta_{0,1} \to \delta_{0,1}(m^2)$

<u>Advantages</u>:

- Twist 2
- Abundant pions
- Crucial FSI are known (δ_0 / δ_1)

Parton process $ab \to cd$	Spin Average Cross Section$-\hat{\sigma}_{ab}^{cd}$	Transversity Dependent Cross Section$-\delta\hat{\sigma}_{ab}^{cd}$
$qg \to qg$	$\frac{\hat{s}^2+\hat{u}^2}{\hat{t}^2} - \frac{4}{9}\frac{\hat{s}^2+\hat{u}^2}{\hat{s}\hat{u}}$	$\frac{\hat{s}\hat{u}}{\hat{t}^2} - \frac{4}{9}$
$\bar{q}g \to \bar{q}g$	$\frac{\hat{s}^2+\hat{u}^2}{\hat{t}^2} - \frac{4}{9}\frac{\hat{s}^2+\hat{u}^2}{\hat{s}\hat{u}}$	$\frac{\hat{s}\hat{u}}{\hat{t}^2} - \frac{4}{9}$
$qq \to qq$	$\frac{4}{9}\left(\frac{\hat{s}^2+\hat{u}^2}{\hat{t}^2} + \frac{\hat{s}^2+\hat{t}^2}{\hat{u}^2}\right) - \frac{8}{27}\frac{\hat{s}^2}{\hat{u}\hat{t}}$	$\frac{4}{27}\frac{\hat{s}}{\hat{t}} - \frac{4}{9}\frac{\hat{s}\hat{u}}{\hat{t}^2}$
$qq' \to qq'$	$\frac{4}{9}\frac{\hat{s}^2+\hat{u}^2}{\hat{t}^2}$	$-\frac{4}{9}\frac{\hat{s}\hat{u}}{\hat{t}^2}$
$q\bar{q} \to q\bar{q}$	$\frac{4}{9}\frac{\hat{s}^2+\hat{u}^2}{\hat{t}^2} + \frac{\hat{u}^2+\hat{t}^2}{\hat{s}^2} - \frac{8}{27}\frac{\hat{u}^2}{\hat{s}\hat{t}}$	$\frac{8}{27}\frac{\hat{u}}{\hat{t}} - \frac{4}{9}\frac{\hat{s}\hat{u}}{\hat{t}^2}$
$q\bar{q}' \to q\bar{q}'$	$\frac{4}{9}\frac{\hat{s}^2+\hat{u}^2}{\hat{t}^2}$	$-\frac{4}{9}\frac{\hat{s}\hat{u}}{\hat{t}^2}$

Where Each Entries multiplies factor $\pi\alpha_s^2/\hat{s}^2$

For example, $qg \to qg$



The dominance of $gq \to gq$ over other processes:

- From Hard cross section

$$\hat{a}_{2T}(gq \to gq) \sim -\frac{2}{5}$$
$$\hat{a}_{LT}(gq \to gq) \sim \frac{2}{11} \qquad \text{at } \theta_{cm} = \frac{\pi}{2}$$

- From distribution

gluon dominates at small x



For example, $qg \to qg$



The dominance of $gg \to gg$ over other processes:

- From Hard cross section

$$\hat{a}_{LT}(gg \to gg) \sim -\frac{2}{5}$$
$$\hat{a}_{LT}(gg \to gg) \sim \frac{2}{11} \qquad \text{at } \theta_{cm} = \frac{\pi}{2}$$

- From distribution

gluon dominates at small x

– *Figure of merit*



Data from P. Estabrooks and A. D. Martin, Nucl. Phys. **B79**, 301 (1974)