

Natural and Cultural Resources

The Brookhaven National Laboratory (BNL) Natural Resource Management Program is designed to protect and manage flora and fauna and the ecosystems in which they exist. The Laboratory's natural resource management strategy is based on understanding the site's resources and maintaining compliance with applicable regulations. The goals of the program include protecting and monitoring the ecosystem, conducting research, and communicating with personnel and the public on ecological issues. BNL focuses on protecting both Federal and New York State threatened and endangered species on site, as well as continuing the Laboratory's leadership role within the greater Long Island Central Pine Barrens ecosystem. Monitoring to determine whether current or historical activities are affecting natural resources is also part of the program. In 2018, deer, vegetation, and soil sampling results were consistent with previous years' results.

The overriding goal of the Cultural Resource Management Program is to ensure that proper stewardship of BNL historic resources is established and maintained. Additional goals of the program include maintaining compliance with various historic preservation and archeological laws and regulations and ensuring the availability of identified resources for research and interpretation.

6.1 NATURAL RESOURCE MANAGEMENT PROGRAM

The Natural Resource Management Program at BNL promotes stewardship of the natural resources found at the Laboratory and integrates natural resource management and protection with BNL's scientific mission. The Natural Resource Management Plan (NRMP) describes the program strategy, elements, and planned activities for managing the various natural resources found on site. The NRMP is updated every five years with the most recent update completed in 2016 (BNL 2016).

6.1.1 Identification and Mapping

An understanding of an environmental baseline is the foundation of natural resource management planning. BNL uses digital global positioning systems (GPS) and geographic information systems (GIS) to clearly relate various "layers" of geographic information (e.g., vegetation types, soil condition, habitat, forest

health, etc.). This is done to gain insight into interrelationships between the biotic systems and physical conditions at the Laboratory.

In 2014, the southern pine beetle (SPB) was discovered at several locations on Long Island, including BNL. Mapping and tracking this native forest pest that rapidly colonizes and spreads through dense stands of pitch pines began in Spring 2015. The Laboratory has continued to work with the New York State Department of Environmental Conservation (NYSDEC) and the U.S. Forest Service to map and track infestations on the Laboratory site. The efforts combine aerial surveys along with ground truthing surveys and mapping. The results of this effort are maintained within the GIS to track impacts to the forest.

Mapping associated with tracking impacts from the operation of the Long Island Solar Farm (LISF) at BNL continues to be entered into the GIS as a tool to analyze changes to wildlife populations and vegetation. In 2018,

Table 6-1. Federal and New York State Threatened, Endangered, Exploitably Vulnerable, and Species of Special Concern at BNL.

Federal and New York State Threatened & Endangered Species, Species of Special Concern, & Species of Greatest Conservation Need			
Common Name	Scientific Name	State Status	BNL Status
Insects			
Comet damer	<i>Anax longipes</i>	SGCN	Confirmed
Rusty patched bumble bee	<i>Bombus affinis</i>	FE	Possible
Frosted elfin	<i>Callophrys iris</i>	T	Likely
New England bluet	<i>Enallagma laterale</i>	SGCN	Likely
Little bluet	<i>Enallagma minusculum</i>	T	Likely
Scarlet bluet	<i>Enallagma pictum</i>	T	Likely
Pine Barrens bluet	<i>Enallagma recurvatum</i>	T	Confirmed
Mottled duskywing	<i>Erynnis martialis</i>	SC	Likely
Persius duskywing	<i>Erynnis persius persius</i>	E	Likely
Pine barrens zanclognatha	<i>Zanclognatha martha</i>	SGCN	Confirmed
Black-bordered lemon moth	<i>Marimatha nigrofimbria</i>	SGCN	Confirmed
Fish			
Banded sunfish	<i>Enneacanthus obesus</i>	T	Confirmed
Swamp darter	<i>Etheostoma fusiforme</i>	T	Confirmed
Amphibians			
Marbled salamander	<i>Ambystoma opacum</i>	SC	Confirmed
Eastern tiger salamander	<i>Ambystoma tigrinum tigrinum</i>	E	Confirmed
Fowler's toad	<i>Bufo fowleri</i>	SGCN	Confirmed
Four-toed salamander	<i>Hemidactylium scutatum</i>	SGCN	Confirmed
Eastern spadefoot toad	<i>Scaphiopus holbrookii</i>	SC	Confirmed
Reptiles			
Worm snake	<i>Carphophis amoenus</i>	SC	Confirmed
Snapping turtle	<i>Chelydra serpentina</i>	SGCN	Confirmed
Spotted turtle	<i>Clemmys guttata</i>	SC	Confirmed
Northern black racer	<i>Coluber constrictor</i>	SGCN	Confirmed
Eastern hognose snake	<i>Heterodon platyrhinos</i>	SC	Confirmed
Stinkpot turtle	<i>Sternotherus odoratus</i>	SGCN	Confirmed
Eastern box turtle	<i>Terrapene carolina</i>	SC	Confirmed
Eastern ribbon snake	<i>Thamnophis sauritus</i>	SGCN	Confirmed
Birds (nesting, transient, or potentially present)			
Cooper's hawk	<i>Accipiter cooperii</i>	SC	Confirmed
Sharp-shinned hawk	<i>Accipiter striatus</i>	SC	Confirmed
Grasshopper sparrow	<i>Ammodramus savannarum</i>	SC	Confirmed
Great egret	<i>Ardea alba</i>	SGCN	Confirmed
Whip-poor-will	<i>Caprimulgus vociferus</i>	SC	Confirmed
Northern harrier	<i>Circus cyaneus</i>	T	Confirmed
Black-billed cuckoo	<i>Coccyzus erythrophthalmus</i>	SGCN	Confirmed
Northern bobwhite	<i>Colinus virginianus</i>	SGCN	Confirmed
Prairie warbler	<i>Dendroica discolor</i>	SGCN	Confirmed
Horned lark	<i>Eremophila alpestris</i>	SC	Confirmed
Wood thrush	<i>Hylocichla mustelina</i>	SGCN	Confirmed
Red-headed woodpecker	<i>Melanerpes erythrocephalus</i>	SC	Confirmed
Osprey	<i>Pandion haliaetus</i>	SC	Confirmed
Scarlet tanager	<i>Piranga olivacea</i>	SGCN	Confirmed
Glossy ibis	<i>Plegadis falcinellus</i>	SGCN	Confirmed
Brown thrasher	<i>Toxostoma rufum</i>	SGCN	Confirmed
Blue-winged warbler	<i>Vermivora pinus</i>	SGCN	Confirmed

continued on next page

natural resource personnel and interns continued to look at use of the LISF site by wildlife, pollinators, changes in bird use, and changes in vegetation.

A wide variety of vegetation, birds, reptiles, amphibians, and mammals inhabit the site. Through implementation of the NRMP, endangered and threatened species, as well as species of special concern, have been identified as having been resident at BNL during the past 30 years or are expected to be present on site (see Table 6-1). The only New York State endangered animal species confirmed as currently inhabiting Laboratory property is the eastern tiger salamander (*Ambystoma t. tigrinum*). Endangered plants that have been confirmed on the BNL site include Engelmann spikerush (*Eleocharis engelmannii*), fireweed (*Erectites heiracifolia* var. *megalocarpa*), Ipecac spurge (*Euphorbia ipecacuanhae*), dwarf huckleberry (*Gaylussacia bigeloviana*), and whorled loosestrife (*Lysimachia quadrifoli*). Three other New York State endangered species have been identified at BNL in the past or are possibly present including: the Persius duskywing butterfly (*Erynnis p. persius*) and crested fringed orchid (*Platanthera cristata*).

Seven threatened species in New York State have been positively identified on site and three other species are considered likely to be present. Threatened species include: two fish (banded sunfish [*Enneacanthus obesus*] and swamp darter [*Etheostoma fusiforme*]); three plants (stiff-leaved goldenrod [*Oligoneuron rigida*], stargrass [*Aletris*

Table 6-1. Federal and New York State Threatened, Endangered, Exploitably Vulnerable, and Species of Special Concern at BNL (concluded).

Federal and New York State Threatened & Endangered Species, Species of Special Concern, & Species of Greatest Conservation Need			
Common Name	Scientific Name	State Status	BNL Status
Mammals			
Northern long-eared bat	<i>Myotis septentrionalis</i>	FT	Confirmed
Plants			
Small-flowered false foxglove	<i>Agalinis paupercula</i>	R	Confirmed
Stargrass	<i>Aletris farinosa</i>	T	Confirmed
Butterfly weed	<i>Asclepias tuberosa</i> ssp. <i>interior</i>	V	Confirmed
Spotted wintergreen	<i>Chimaphila maculata</i>	V	Confirmed
Flowering dogwood	<i>Cornus florida</i>	V	Confirmed
Pink lady's slipper	<i>Cypripedium acaule</i>	V	Confirmed
Ground pine	<i>Dendrolycopodium obscurum</i>	V	Confirmed
Round-leaved sundew	<i>Drosera rotundifolia</i> var. <i>rotundifolia</i>	V	Confirmed
Marginal wood fern	<i>Dryopteris marginalis</i>	V	Confirmed
Engelman spikerush	<i>Eleocharis engelmannii</i>	E	Confirmed
Fireweed	<i>Erectites heiracifolia</i> var. <i>megalocarpa</i>	E	Confirmed
Ipecac spurge	<i>Euphorbia ipecacuanhae</i>	E	Confirmed
Eastern showy aster	<i>Eurybia spectabilis</i>	T	Confirmed
Dwarf huckleberry	<i>Gaylussacia bigeloviana</i>	E	Confirmed
Winterberry	<i>Ilex verticillata</i>	V	Confirmed
Sheep laurel	<i>Kalmia angustifolia</i>	V	Confirmed
Narrow-leaved bush clover	<i>Lespedeza angustifolia</i>	R	Confirmed
Wild lupine	<i>Lupinus perennis</i>	R	Confirmed
Whorled loosestrife	<i>Lysimachia quadrifolia</i>	E	Confirmed
Bayberry	<i>Myrica pensylvanica</i>	V	Confirmed
Stiff-leaved goldenrod	<i>Oligoneuron rigida</i>	T	Confirmed
Cinnamon fern	<i>Osmunda cinnamomea</i>	V	Confirmed
Clayton's fern	<i>Osmunda claytoniana</i>	V	Confirmed
Royal fern	<i>Osmunda regalis</i>	V	Confirmed
Crested fringed orchid	<i>Plantanthera cristata</i>	E	Likely
Green fringed orchid	<i>Platanthera lacera</i>	V	Confirmed
Prostate knotweed	<i>Polygonum aviculare</i> ssp. <i>buxiforme</i>	E	Possible
Bracken fern	<i>Pteridium alquilinum</i> var. <i>pseudocaudatum</i>	E	Possible
Swamp azalea	<i>Rhododendron viscosum</i>	V	Confirmed
Long-beaked bald-rush	<i>Rhynchospora scirpoides</i>	R	Confirmed
New York fern	<i>Thelypteris novaboracensis</i>	V	Confirmed
Marsh fern	<i>Thelypteris palustris</i> var. <i>pubescens</i>	V	Confirmed
Possum haw	<i>Viburnum nudum</i> var. <i>nudum</i>	E	Possible
Virginia chain-fern	<i>Woodwardia virginica</i>	V	Confirmed

Notes:
 information based on 6 NYCRR Part 182, 6 NYCRR Part 193, and BNL survey data.
 E = endangered
 FE = federally endangered
 FT = federally threatened
 R = rare
 SC = species of special concern
 SGCN = species of greatest conservation need
 T = threatened
 V = exploitably vulnerable

farinosa], and eastern showy aster [*Eurybia spectabilis*]); the northern harrier (*Circus cyaneus*) is periodically seen in the fall. Insects listed as threatened include a damselfly Pine Barrens bluet (*Enallagma recurvatum*) which was confirmed at one of the many coastal plain ponds located on site. Two other damselflies, the little bluet (*Enallagma minisculum*) and the scarlet bluet (*Enallagma pictum*), are likely to be present at one or more of the ponds on site. The frosted elfin (*Callophrys irus*), a butterfly, has been historically present on site due to its preferred habitat and host plant, wild lupine (*Lupinus perennis*).

A number of other species that are listed as rare, of special concern, or exploitably vulnerable by New York State either currently inhabit the site, visit during migration, or have been identified historically.

BNL has one federally threatened species, the northern long-eared bat (*Myotis septentrionalis*) that is found within the forests of the Lab; and the federally endangered rusty-patch bumble bee (*Bombus affinis*) was likely to have been in the area historically. A single bee tentatively identified as *B. affinis* was identified in 2016, but no photo or specimen was kept. Subsequent searches in 2017 and 2018 did not yield evidence for its presence.

6.1.2 Habitat Protection and Enhancement

BNL has administrative processes in place to protect on-site habitats and natural resources. Activities to eliminate

or minimize negative effects on endangered, threatened, or sensitive species are either incorporated into Laboratory procedures or into specific program or project plans. Human access to critical habitats, when necessary, is limited, and habitats are enhanced to improve survival or increase populations. Routine activities, such as road maintenance, are not performed until the planned activities have been evaluated and determined to be unlikely to affect habitat.

6.1.2.1 Salamander Protection Efforts

Many safeguards are in place to protect eastern tiger salamander breeding areas. BNL staff must review any project planned near eastern tiger salamander habitats, and every effort is made to minimize impacts. A map of the breeding areas is reviewed when new projects are proposed. The map is updated as new information concerning the salamanders is generated through research and monitoring. The current map incorporates buffer areas around tiger salamander habitats of 1,000 feet based on guidance from NYSDEC. Other efforts to protect this state-endangered species include determining when adult salamanders are migrating toward breeding locations, when metamorphosis has been completed, and when juveniles are migrating after metamorphosis. During these times, construction and maintenance activities near their habitats are postponed or closely monitored.

Water quality testing is conducted as part of the routine monitoring of recharge basins, as discussed in Chapter 5. In cooperation with NYSDEC, habitat surveys have been routinely conducted since 1999. Biologists conducting egg mass and larval surveys have confirmed that 26 on-site ponds are used by eastern tiger salamanders. In 2018, surveys confirmed the presence of salamanders in seven of the 26 ponds; early winter surveys in late 2018 documented a total of 119 egg masses in two ponds and four adult salamanders in two other ponds. Long Island's drought finally concluded at the end of 2017 and ponds held water for most of the year in 2018.

6.1.2.2 Banded Sunfish

Banded sunfish protection efforts include

observing whether adequate water is present within areas currently identified as sunfish habitat, ensuring that vegetation in their habitat is not disturbed, and evaluating all activities taking place in ponds and the Peconic River on site for potential impacts on these habitats. Population estimates are periodically conducted within these waters to determine their current health. During the last population survey in 2011, approximately 6,400 banded sunfish were counted. Drought conditions that lasted from 2015 through early 2017 likely resulted in the extirpation of the banded sunfish from the BNL site. The single known habitat held water throughout 2018 and could likely sustain sunfish. Regionally, NYSDEC determined that only a few populations of banded sunfish survived the drought and is evaluating the need for restoration efforts.

6.1.2.3 Migratory Birds

A total of 216 species of birds have been identified at BNL since 1948; at least 85 species are known to nest on site. Some of these nesting birds have shown declines in their populations nationwide over the past 30 years. The Laboratory conducts routine monitoring of songbirds along seven permanent bird survey routes in various habitats on site.

In 2018, monthly surveys were conducted starting at the end of April and extending through the end of August. These surveys identified 67 bird species, compared to the 72 species identified in 2017 and 77 species in 2016. A total of 134 bird species have been identified in surveys in the past 19 years; 59 of these species were present in each of the past 19 years. Variations in the number and species identified during each survey may reflect the time of observation, variations in weather patterns between years, and possible changes in the environment.

The three most diverse transects near on-site wetlands are by the LISF and the Peconic River. The four transects passing through the various forest types on site (e.g., white pine, moist pine barrens, and dry pine barrens) showed a less diverse bird community. Bird survey data are stored in an electronic database for future reference and study. No known data on the

effects of a large, utility-scale solar array such as the LISF are known within scientific literature. To assess the effects of the LISF on local bird populations, the collection of migratory bird data in both the Biology Field and Solar Farm transects is important. The LISF vegetation and the way it is managed may play a key role as habitat for migratory birds. One species, the indigo bunting (*Passerina cyanea*), was absent along the Biology Field transect in 2011 during construction, but was heard along the Solar Farm transect in 2012, returned to the Biology Field transect in 2013, and has been present on both transects since 2014. This temporary absence is thought to be due to disturbance from construction activities while building the LISF.

The eastern bluebird (*Sialia sialis*) has been identified as a declining species of migratory birds in North America. This is due to loss of habitat and nest site competition from European starlings (*Sturnus vulgaris*) and house sparrows (*Passer domesticus*). BNL's NRMP includes habitat enhancement for the eastern bluebird. Since 2000, the Laboratory has installed more than 60 nest boxes around open grassland areas on site to enhance their population. Although many of these boxes were removed from service in 2010 in preparation for the construction of the LISF, the LISF created nearly 200 acres of suitable habitat for the eastern blue bird. Forty new boxes were installed around the northern most portions of the LISF in 2012 and are routinely used by bluebirds, house wrens, and tree swallows.

Migratory birds occasionally cause safety and health concerns, particularly Canada geese (*Branta canadensis*) and several species of migratory birds that occasionally nest on buildings or in construction areas on site. In 2006, it was determined that the resident Canada goose population at BNL reached large enough numbers that could result in health and safety issues. Beginning in 2007, under a permit from U.S. Fish and Wildlife Service, the Laboratory began managing the resident goose population by limiting the number of eggs that could hatch. Sixty-five nests were treated during 2018 to reduce the number of goslings. This was an increase of 20

nests over those treated in 2017. The increase was due to geese that hatched in 2014 and continued to reach sexual maturity. Had the egg oiling not occurred, the goose population would have increased by up to 287 birds. During 2018, approximately 20 goslings were produced, with about 30 percent surviving due to predation. By the end of 2018, the resident goose population was estimated at just over 100 birds due to population reduction resulting from hunting in the local area.

6.1.2.4 Bald Eagle

The bald eagle (*Haliaeetus leucocephalus*) has been increasing in population locally on Long Island with eight known nest sites on the island. During 2018, bald eagles were sighted numerous times in the area of the Sewage Treatment Plant (STP); throughout the fall of 2018, a pair of sub-adult eagles (one of which is pictured on the cover of this year's Site Environmental Report) was routinely observed on the BNL site. As the eagle population increases on Long Island, the potential for them to nest on the BNL site will increase as well.

6.1.2.5 Northern Long-eared Bat

As discussed in Section 6.1.1, the northern long-eared bat was added to the list of federally threatened species in 2015. BNL began planning for the eventual listing and put in place actions to minimize the likelihood of impacting this species. The two most likely activities that could impact this bat on the BNL site are building demolitions and prescribed fires. Inspections for the presence of bats are conducted in multiple ways prior to a building demolition. During spring, summer, and fall, ultrasonic acoustic monitoring is conducted around buildings scheduled for demolition to determine if there is bat activity. Regardless of the outcome of acoustic monitoring, a final internal inspection of the buildings is conducted approximately 24 hours prior to demolition to verify the absence of bats. For growing season prescribed fire, acoustic monitoring may be done within the burn unit to determine if there is bat activity. If positive results occur, surveys of the entire burn unit are completed

to identify potential roost trees and appropriate protections are put into place to ensure that bats are not impacted by fire. In 2018, only one building was demolished, and there was no impact to bats. Surveys ahead of the single prescribed fire in 2018 identified no roost trees.

6.1.3 Population Management

In addition to controlling resident Canada goose populations described above, the Laboratory also monitors or manages other populations, including species of interest, to ensure that they are sustained and to control invasive species.

6.1.3.1 Wild Turkey

The forested areas of BNL provide good nesting and foraging habitat for wild turkey (*Meleagris gallapavo*). In 2018, the on-site population continued to be approximately 500 birds due to successful nesting. Each year, NYSDEC manages a hunting period during the week of Thanksgiving, and a youth-only hunt in May for several areas across Long Island, which typically results in approximately 100 birds taken.

6.1.3.2 White-Tailed Deer

BNL consistently updates information on the resident population of white-tailed deer (*Odocoileus virginianus*). As there are no natural predators on site and hunting is not permitted at the Laboratory, there are no significant pressures on the population to migrate beyond their typical home range of approximately one square mile. Normally, a population density of ten to 30 deer per square mile is considered an optimum sustainable level for a given area. This would equate to approximately 80 to 250 deer inhabiting the BNL property under optimal circumstances. This was the approximate density in 1966, when BNL reported an estimate of 267 deer on site (Dwyer 1966). The Laboratory has been conducting routine population surveys of the white-tailed deer since 2000. The fall 2018 estimate provided a range of 350-450 animals.

Deer overpopulation can affect animal and human health (e.g., animal starvation, Lyme disease from deer ticks, and collision injuries

to both humans and animals), species diversity (e.g., songbird species reduction due to selective grazing and destruction of habitat by deer), and property damage (e.g., collision damage to autos and browsing damage to ornamental plantings). Deer-related collisions on site increased in 2018 compared to recent years, potentially an indication of increased population since the 2017 deer harvest, or it may reflect the need for additional employee education as a reminder to abide by speed limits and pay attention to wildlife.

High deer populations are a regional problem, and the Laboratory is just one area on Long Island with such an issue. Multiple east end towns are now managing deer populations either through culls (aka deer harvests), hunting, or sterilization programs. In 2008, BNL began developing a deer management plan which included an option to reduce the population through culling. The planning effort included engagement of Laboratory employees and guests in discussions concerning the need and methods for deer management. In 2012, an Environmental Assessment (EA) under the National Environmental Policy Act (NEPA) was completed and sent to New York State for comment. The Final EA was completed in the spring of 2013. Additionally, under BNL's permit for deployment of the 4-Poster tick management system issued by NYSDEC, the Laboratory is required to implement a deer management program. In February 2015, 300 animals were taken, effectively reducing the population to approximately 530 animals. Furthermore, as many as 100 additional animals did not survive the harsh winter conditions which resulted in snow cover lasting more than 30 consecutive days. A second population reduction occurred in March 2017, with 202 animals being removed, bringing the herd to a range of 200-300 animals. As mentioned above, the population at the end of 2018, accounting for reproduction, was estimated at between 350 and 450 animals with a third population reduction planned for spring 2019.

6.1.4 Compliance Assurance and Potential Impact Assessment

The NEPA review process at BNL ensures

that environmental impacts of a proposed action or activity are adequately evaluated and addressed. The Laboratory uses NEPA reviews when identifying potential environmental impacts associated with site activities, especially projects that may result in physical alterations to the landscape and structures. As appropriate, stakeholders such as EPA, NYSDEC, Suffolk County Department of Health Services (SCDHS), BNL's Community Advisory Council, and the Brookhaven Executive Roundtable are involved in reviewing major projects that have the potential for significant environmental impacts. Formal NEPA reviews are coordinated with the State of New York. There were no higher level NEPA reviews started or completed in 2018.

6.2 UPTON ECOLOGICAL AND RESEARCH RESERVE

The Upton Ecological and Research Reserve (Upton Reserve) consists of 530 acres located on the eastern boundary of the BNL site. The reserve has been designated as an area for the protection of sensitive habitats and a place where researchers can study local ecosystems. The Upton Reserve is home to a wide variety of flora and fauna. It contains wetlands and is largely within the core preservation area of the Long Island Central Pine Barrens. Based on information from a 1994-1995 biological survey of the Laboratory, experts believe the reserve is home to more than 200 plant species and at least 162 species of mammals, birds, fish, reptiles, and amphibians (LMS 1995).

The Upton Reserve is managed by BNL and the Foundation for Ecological Research in the Northeast (FERN). Funding is coordinated for research projects that occur within the reserve and the larger pine barrens area of Long Island.

6.3 MONITORING FLORA AND FAUNA

The Laboratory routinely conducts surveillance monitoring of flora and fauna to determine the effects of past and present activities on site. Because soil contaminated with a radioactive isotope of cesium, Cs-137, was used in some BNL landscaping projects in the past, traces of Cs-137 attributable to past practices

and world-wide fallout can be found in deer and other animals and plants. At the cellular level, Cs-137 takes the place of potassium (K), an essential nutrient. Most radionuclide tables in this chapter also list analytical results for potassium-40 (K-40), a naturally occurring radioisotope of potassium that is commonly found in flora and fauna. Studies indicate that Cs-137 out-competes potassium when potassium salts are limited in the environment, which is typical on Long Island. Including K-40 in tables allows for a comparison with Cs-137 levels and is used, in part, to determine the accuracy of analytical results. The results of the annual sampling conducted under the flora and fauna monitoring program follow.

6.3.1 Deer Sampling

White-tailed deer in New York State are typically large, with males weighing, on average, approximately 150 pounds; females typically weigh approximately 100 pounds. However, white-tailed deer on Long Island tend to be much smaller, weighing an average of 80 pounds. The meat available for consumption from local deer ranges from 20 to 40 pounds per animal. Samples of meat and liver are taken from each deer, when possible, and are analyzed for Cs-137. Data are reported on a wet-weight basis, as that is the form most likely used for consumption.

Since 1996, BNL has routinely collected deer samples from on- and off-site areas. While most off-site samples are the result of car/deer accidents near the Laboratory, samples from deer taken by hunters beyond BNL boundaries or samples from car/deer accidents greater than one mile from BNL have also been made available for analysis. In 1998, a statistical analysis suggested that 40 deer from off site and 25 deer from on site are needed to achieve a statistically sound data set. The number obtained each year has not met this preferred level because sample availability depends on accidents between vehicles and deer and people reporting dead deer. In 2018, a total of 18 deer were taken both on and off the BNL site. Figure 6-1 shows the location of all deer samples taken within a five-mile radius of the Laboratory

since 2014. Most of the off-site samples are concentrated along the William Floyd Parkway on the west boundary of BNL, whereas historically most on-site samples are collected near the Laboratory's main entrance gate and the developed portions of the site. This distribution is due to the fact that people on their way to work see and report dead deer. Also, vehicle collisions with deer on site occur primarily early or late in the day, when deer are more active and traffic to and from the Lab's Main Gate is greatest.

Based on more than a decade of sampling, deer taken from more than one mile from BNL are used for comparison with populations on and near the Laboratory that could acquire Cs-137 from a BNL source. In 2018, six deer were obtained on site, eight from off-site locations within one mile of the Laboratory, and four from greater than one mile from the BNL boundary, all from car/deer accidents. The analytical results of deer sampling are shown in Table 6-2.

6.3.1.1 Cesium-137 in White-Tailed Deer

Based on historic and current data, white-tailed deer sampled at or near the Laboratory contain higher concentrations of Cs-137 than deer from greater than one mile off site. This is most likely because the deer graze on vegetation growing in soil where elevated Cs-137 levels are known to exist. Cesium-137 in soil can be transferred to above-ground plant matter via root uptake, where it then becomes available to browsing/grazing animals or is consumed directly with soil while the animal is grazing. Remediation of contaminated soil areas on site has occurred under the Laboratory's CERCLA program, with all major areas of contaminated soil being remediated by September 2005.

In 2018, Cs-137 concentrations in deer meat samples were obtained from six deer on site with a range of values from 0.01 pCi/g, wet weight to 2.19 pCi/g, wet weight, and an arithmetic average of 0.97 pCi/g, wet weight, as shown in Table 6-2. The wet weight concentration is before a sample is dried for analysis and is the form most likely to be consumed. Dry

weight concentrations are typically higher than wet weight values. The highest on-site sample in 2018 (2.19 pCi/g, wet weight) was about 63 percent higher than the highest on-site sample reported in 2017 (1.34 pCi/g, wet weight) and five times lower than the highest level ever reported in 1996 (11.74 pCi/g, wet weight).

Cs-137 concentrations in off-site deer meat samples are typically separated into two groups: samples taken within one mile of BNL (eight samples) and samples taken farther away (four samples), as shown in Table 6-2. Concentrations in meat samples taken within one mile ranged from 0.03 pCi/g, wet weight to 2.14 pCi/g, wet weight, with an arithmetic average of 0.76 pCi/g, wet weight. Because deer on site may routinely travel up to one mile off site, the arithmetic average for deer taken on site and within one mile of the Laboratory is also calculated; for 2018, this was 0.85 pCi/g, wet weight. The four deer sampled from greater than one mile from BNL had Cs-137 concentrations ranging between 0.02 pCi/g, wet weight, to 2.29 pCi/g, wet weight, with an arithmetic average of 1.05 pCi/g, wet weight. Figure 6-2 compares the average values of Cs-137 concentrations in meat samples collected in 2018 from four different location groupings. Beginning in 2013, the average Cs-137 content from deer taken within one mile of the Laboratory was lower than the on-site average, and this pattern has been consistent for the past six years. While no definitive explanation can be given to the difference from past results, it could simply be an artifact of low sample numbers and randomness in sample acquisition. Although not shown on Figure 6-2, Cs-137 concentrations in seven of the 14 meat samples taken both on and off site were below 0.5 pCi/g, wet weight.

Figure 6-3 presents the ten-year trend of on-site and near off-site Cs-137 averages in deer meat. The 2018 average is approximately 25 percent lower than the 2009 average and is three times higher than the 2015 value of 0.28 pCi/g wet weight, which was the lowest average seen since trending began in 2000. The higher averages shown are reflective of a significant number of samples taken in the fall




Figure 6-1. Deer Sample Locations, 2014—2018.

CHAPTER 6: NATURAL AND CULTURAL RESOURCES

Table 6-2. Radiological Analyses of Deer Tissue (2018).

Sample Location	Collection Date	Tissue	K-40 pCi/g (Wet Weight)	Cs-137 pCi/g (Wet Weight)
BNL				
East Fifth Ave. and First St.	3/9/18	Flesh	2.72±0.28	0.06±0.01
		Liver	2.40±0.51	ND
RHIC Bldg. 1005	4/16/18	Flesh	2.27±0.21	0.01±0.01
		Liver	1.93±0.45	ND
BNL Apartment Area, East St.	9/11/18	Flesh	3.25±0.21	0.74±0.02
		Liver	2.23±0.43	0.19±0.03
Bldg. 1005E, front of building	11/9/18	Flesh	3.04±0.26	1.79±0.05
		Liver	2.55±0.26	0.78±0.04
NSLS II Outer Loop Rd, east side	11/28/18	Flesh	3.52±0.21	2.19±0.04
		Liver	2.27±0.16	0.33±0.01
RHIC Bldg. 1005	12/12/18	Flesh	3.18±0.20	1.03±0.03
		Liver	2.15±0.20	0.43±0.02
< 1 Mile from BNL				
LIE Service Rd at south gate	1/19/18	Flesh	2.87±0.33	0.19±0.03
		liver	2.24±0.40	0.04±0.02
LIE South Service Road and WFPKWY	4/7/18	Flesh	2.97±0.29	0.03±0.01
		liver	1.78±0.29	ND
WFPKWY 1/2 mile south of main gate	10/6/18	Flesh	3.38±0.15	2.14±0.03
		liver	1.85±0.16	0.85±0.02
WFPKWY 1/2 mile north of main gate	11/9/18	Flesh	3.35±0.27	0.32±0.02
		liver	2.58±0.24	0.08±0.01
WFPKWY north gate median	11/9/18	Flesh	3.61±0.35	0.20±0.02
		liver	2.66±0.27	0.09±0.01
WFPKWY, 1/2 mile south of main gate	11/12/18	Flesh	3.12±0.29	0.48±0.03
		liver	2.92±0.29	0.13±0.02
LIE Service Rd. south gate deer one	12/5/18	Flesh	3.08±0.12	1.18±0.02
		liver	2.72±0.19	0.26±0.02
LIE Service Rd. south gate deer two	12/5/18	Flesh	3.24±0.13	1.55±0.02
		liver	2.52±0.14	0.25±0.01
> 1 Mile from BNL				
Rte 25 and Pine Bark Rd	2/14/18	Flesh	3.17±0.36	0.11±0.02
		Liver	2.23±0.30	0.03±0.01
Manorville, Hotwater St. and Chapman Blvd	4/29/18	Flesh	2.74±0.31	0.02±0.02
		Liver	2.89±0.37	ND
Rte. 25 1/2 mile west of Wading River Rd	11/11/18	Flesh	2.90±0.28	2.29±0.06
		Liver	2.29±0.27	0.50±0.03
LIE @ Exit 70 Manorville	12/9/18	Flesh	3.18±0.27	1.79±0.05
		Liver	2.57±0.25	0.43±0.03
Averages by Tissue				
Flesh Averages				
All Samples (18)			3.09±1.10	0.90±0.13
BNL Average (6)			3.00±0.56	0.97±0.08
< 1 Mile Average (8)			3.20±0.72	0.76±0.06
BNL + < 1 Mile Average (14)			3.11±0.91	0.85±0.10
> 1 Mile Average (4)			3.00±0.61	1.05±0.08
Liver Averages				
All Samples (18)			2.38±1.30	0.25±0.09
BNL Average (6)			2.26±0.88	0.29±0.06
< 1 Mile Average (8)			2.41±0.73	0.21±0.05
BNL + < 1 Mile Average (14)			2.34±1.15	0.25±0.08
> 1 Mile Average (4)			2.50±0.60	0.24±0.04

Notes: All values are shown with a 95% confidence interval. K-40 Occurs naturally in the environment and is presented as a comparison to Cs-137
All averages are the arithmetic average with confidence limits using a 2 sigma (95%) propagated error. ND = not detected

when Cs-137 levels are typically higher. However, these sample results continue to indicate the effectiveness of cleanup actions across the Laboratory, with the trend being downward from 2009 to 2018 and the ten-year average being 0.79 pCi/g.

The effectiveness of the BNL soil cleanup program and the reduction of Cs-137 in deer meat was evaluated by Rispoli, et al. (2014). The average Cs-137 content was shown to be statistically lower than before cleanup. Samples taken at distances greater than one mile from the BNL site were shown to remain consistent before and after cleanup, while the on-site and near off-site values were shown to decline. In 2017, while preparing for monitoring associated with the reduction of the deer population, the ten-year average for on-site deer samples was calculated to be 1.0 pCi/g, wet weight, and this value was used to establish an administrative release criterion for deer meat made available for donation.

When possible, liver samples are taken concurrently with meat samples. The liver generally accumulates Cs-137 at a lower rate than muscle tissue. The typically lower values in liver allow the results to be used as a validity check for meat values (i.e., if liver values are higher than meat values, results can be considered questionable and should be confirmed). In liver samples collected on site in 2018, Cs-137 concentrations ranged from non-detect to 0.78 pCi/g, wet weight, with an average of 0.29 pCi/g, wet weight. The near off-site Cs-137 concentration in liver ranged from non-detect to 0.85 pCi/g, wet weight, with an arithmetic average for off-site liver samples within one mile of 0.21 pCi/g, wet weight. Liver samples from deer taken greater than one mile from BNL ranged from non-detect to 0.50 pCi/g, wet weight with the arithmetic average being 0.24 pCi/g, wet weight. The potential radiological dose resulting from deer meat consumption is discussed in Chapter 8.

The New York State Department of Health (NYSDOH) has formally considered the potential public health risks associated with elevated Cs-137 levels in on-site deer and determined that neither hunting restrictions nor formal health advisories are warranted (NYSDOH 1999). As mentioned above, BNL has established an

administrative release criterion of 1.0 pCi/g, wet weight for meat from deer removed from the Laboratory and donated for consumption. No deer were taken for population management in 2018.

With respect to the health of on-site deer based on their exposure to radionuclides, the International Atomic Energy Agency (IAEA) has concluded that chronic dose rates of 100 millirad per day to even the most radiosensitive species in terrestrial ecosystems are unlikely to cause detrimental effects in animal populations (IAEA 1992). A deer containing a uniform distribution of Cs-137 within muscle tissue at the highest levels observed to date (11.74 pCi/g, wet weight, reported in 1996) would carry a total amount of approximately 0.2 μ Ci. That animal would receive an absorbed dose of approximately 3 millirad per day, which is only three percent of the IAEA threshold. The deer observed and sampled on site appear to have no health effects from the level of Cs-137 found in their tissues.

6.3.2 Other Animals Sampled

When other animals, such as wild turkey or Canada geese, are found dead along the roads of BNL and the immediate vicinity due to road mortality, they are tested for Cs-137. No other animals were sampled in 2018.

6.3.3 Fish Sampling

BNL maintains an ongoing program for collecting and analyzing fish from the Peconic River and surrounding freshwater bodies. Monitoring of the river has been conducted under the environmental surveillance program and the CERCLA post-cleanup program. Surveillance monitoring had occurred during even-numbered years and post-cleanup monitoring occurred in odd-numbered years. However, with the discontinuance of discharges from the STP to the Peconic River in September 2014 and current below-average amounts of precipitation, the objectives for the fish monitoring program have changed to reflect the current intermittent presence of water in the on-site portions of the river. Fish are now only sampled under the surveillance




Figure 6-2. Comparison of Cs-137 values in deer flesh for onsite, offsite within 1 mile, onsite and near offsite, and offsite greater than 1 mile from the Laboratory.




Figure 6-3. Ten year trend in Cs-137 in deer flesh for samples taken at BNL and within 1 mile of the Laboratory. Average before clean-up (dashed line) 2.57 pCi/g wet weight.

program when there is enough water to support a sufficient population of fish that can be sampled without harm to their population and that are of sufficient size for analysis. Based upon the 2016 CERCLA Five-year Review of the effectiveness of the environmental cleanup and the final supplemental cleanup of a small area within the river during 2017, the Laboratory has discontinued fish monitoring under the CERCLA program. However, when conditions allow, fish sampling will be conducted under the surveillance program. Due to lack of water and fish within the on-site portions of the Peconic River, no fish were sampled in 2018.

6.3.3.1 Fish Population Assessment

The relative sizes of fish caught during annual sampling events are tracked and modifications to future sampling events are made, as necessary, to ensure long-term health of the on-site fish populations. Successful sampling of sufficiently large fish for analysis from 2008 through 2015, even with low water levels in the on-site portion of the Peconic River, indicated that fish populations could maintain themselves. However, the combination of discontinuing STP discharges to the Peconic River and continued drought conditions have resulted in the on-site portions of the Peconic River being totally dry and no longer able to support fish. The river remained dry through July 2018 when groundwater levels began to rise. By the end of 2018, water levels were sufficiently high, resulting in off-site flow at station HQ on the east boundary of the Laboratory. For fish populations to survive and flourish, water levels must be substantial enough to allow migration of fish and maintain their presence for an extended period of time to replenish populations. As mentioned above, new criteria for the collection of fish samples have been developed. These criteria will guide the environmental monitoring approach for fish in the future.

6.3.4 Vegetation Sampling

6.3.4.1 Grassy Plants and Soil

During 2018, grassy vegetation samples were collected from ten locations around the Laboratory (Figure 6-4) and a control location

at the NYSDEC hunter check station in Ridge, New York. All samples were analyzed for Cs-137 (see Table 6-3). Cs-137 content in vegetation ranged from non-detectable to 0.22 pCi/g, wet weight, in two areas on both the south and north boundaries of the Laboratory. Soil samples had Cs-137 levels from non-detect to 1.36 pCi/g, dry weight. All values were consistent with historic monitoring and knowledge of cleanup areas. Monitoring results for grassy vegetation and soils were utilized for the annual dose to biota analysis reported in Chapter 8.

6.4 PRECIPITATION MONITORING

6.4.1 Mercury Monitoring of Precipitation

During 2018, precipitation samples were collected quarterly at air monitoring Stations P4 and S5 (Figure 4-2 for station locations). The samples were analyzed for total mercury (Table 6-4) using low-level mercury analysis. Until 2015, BNL had routinely analyzed precipitation for radiological content. However, with no emissions of significantly long-lived radionuclides from Laboratory operations, the monitoring program objectives were modified to remove testing of precipitation for radiological content beginning in 2016.

Mercury concentrations in precipitation have been measured at BNL since 2007. Analysis of mercury in precipitation is conducted to document mercury deposition that is attributable to off-site sources. This information has been used as a comparison to Peconic River monitoring data and aids in understanding the distribution of mercury within the Peconic River watershed. Mercury was detected in all the precipitation samples collected at both sampling stations. Mercury ranged from 1.58 ng/L at station S5 in January to 8.59 ng/L at station P4 in April. The 8.59 ng/L concentration is five times lower than the highest value of 45.1 ng/L, recorded in 2017.

6.5 WILDLIFE PROGRAMS

BNL sponsors a variety of educational and outreach activities involving natural resources. These programs are designed to help participants understand the ecosystem, foster an interest in science, and provide a meaningful experience for interns in preparation for further

Table 6-3. Radiological analysis of grassy vegetation and associated soils.

Location/Matrix	K-40 pCi/g±95% C.I.	Cs-137 pCi/g±95% C.I.
South of RHIC east of Renaissance Rd		
Vegetation	2.51±0.34	ND
Soil	7.64±1.01	0.28±0.08
Tank Pond		
Vegetation	1.06±0.45	0.22±0.05
Soil	9.86±1.77	1.36±0.18
South Boundary Road, South Side		
Vegetation	2.03±0.56	ND
Soil*	7.60±0.98	0.05±0.02
North Firebreak, south side west of Crescent Path		
Vegetation	0.87±0.43	0.22±0.07
Soil*	3.66±0.57	0.14±0.06
North Firebreak, south side east of Crescent Path		
Vegetation	1.09±0.48	ND
Soil*	4.73±0.78	0.07±0.05
East Firebreak, west side, south of East Fifth Ave.		
Vegetation	1.55±0.67	ND
Soil*	4.55±0.70	0.09±0.04
East Firebreak, east side, north of LISF Array 2		
Vegetation	3.24±0.71	ND
Soil	7.32±1.08	ND
RHIC North Pond		
Vegetation	1.96±0.55	ND
Soil*	7.66±0.90	0.08±0.05
BGRR Hill west of Stack		
Vegetation	4.29±0.52	ND
Soil	6.17±0.93	0.22±0.05
Tandem to AGS Tunnel		
Vegetation*	1.46±0.48	0.02±0.02
Soil*	5.62±0.63	0.13±0.05
NYSDEC Game Farm (Control)		
Vegetation	1.90±0.45	ND
Soil	5.66±0.85	0.22±0.07

Notes:

All values are shown with a 95% confidence interval.
 Radiological values for soils are on a 'dry weight' basis.
 K-40 occurs naturally in the environment and is presented as
 a comparison to Cs-137.

Cs-137 = cesium-137

K-40 = potassium-40

ND = not detected

* = estimated value for Cs-137 based on laboratory qualifiers.

Table 6-4 Precipitation Monitoring (Mercury).

Location/Period	Mercury ng/L
P4	
1/17/18	2.02
4/9/18	8.59
7/18/18	4.39
10/12/18	4.94
S5	
1/17/18	1.58
4/9/18	8.46
7/18/18	3.7
10/12/18	3.85

Notes:

Method detection limit for mercury is 0.2 ng/L.

P4 = precipitation sampler near BNL Apartment area.

S5 = precipitation sampler near BNL Sewage
Treatment Plant.

studies or a career. Wildlife programs are conducted at the Laboratory in collaboration with the Department of Energy, local agencies, colleges, and high schools. Ecological research is also conducted on site to routinely update the natural resource inventory records, gain a better understanding of the ecosystem, and guide management planning.

In 2018, BNL hosted ten student interns and two faculty members within the Natural Resources program. Two of the interns worked with a faculty member from Hofstra University as part of the BNL Visiting Faculty Program (VFP), five worked under a faculty member from Southern University of New Orleans, and three interns participated in research associated with various projects including several related to the LISF, turtles, forest health and deer impacts, and pollinators. All interns and faculty participated in the summer internship programs.

- The VFP team from Hofstra University continued a third year of gathering basic information on small mammals, tick loads, and the incidence of Lyme disease in the ticks. This work is being done in preparation for coyotes (*Canis latrans*) migrating to Long Island. Once established, coyotes




Figure 6-4 Vegetation and Soil Sampling Locations

are expected to alter ecosystems due to competition with other carnivores.

- Work associated with the LISF involved tracking 20 eastern box turtles outfitted with transmitters to determine home range sizes. Many of the turtles were captured in or near the LISF to determine if they utilize habitats found in the facility. Since 2011, student interns have followed a total of 42 turtles; as a result, BNL is building a thorough understanding of their habits. Turtles are also permanently marked to facilitate identification of individual turtles as part of a mark-recapture effort. Turtle research also looked at micro- and macro-habitat use.
- A graduate student from the State University of New York (SUNY) at Stony Brook continued mark-recapture studies of box turtles, updating the database of 600-plus marked and recaptured turtles to determine potential impact to box turtles based on expected impacts to meso-predators when coyotes become established on Long Island.
- Interns conducted pollinator surveys of the LISF to determine use. The research effort focused on pollinator resources and resource partitioning by bumble bees (*Bombus spp.*), honey bees (*Apis mellifera*), and other pollinators by comparing solar fields 1, 2, 4, and 6.
- With efforts to manage the deer population on the BNL site, vegetation surveys were carried out to determine continuing impacts. BNL has established several deer exclosures (i.e., fenced areas that prevent deer from getting into an area) that can be compared to paired areas of open forest. During efforts to gather data, several of the exclosures were found with damage significant enough to allow deer to enter, and one had been totally removed. Long-term monitoring will likely have to rely on use of forest health monitoring plots established in 2005-2006, and periodically revisited.
- BNL has significant data resources related to breeding bird surveys. Survey data was

reviewed and compared by an intern to determine if there was any lasting impact from the construction of the LISF. The effort indicated that construction had an initial impact on use of the area adjacent to solar field 1 for several years during and after construction, but bird populations appeared to return to normal after three to four years post-construction.

The Lab's Natural Resource program within the Environmental Protection Division has had a long-term relationship with Dr. Murty Kambhampati from Southern University at New Orleans. Beginning in 2006, the summer after Hurricane Katrina, Dr. Kambhampati began bringing interns to BNL to gain experience in natural resources, ecology, and environmental sciences. Dr. Kambhampati was featured on BNL media with his interns (see photo) during the summer of 2018 for his mentoring efforts, which have introduced 35 interns to BNL over



Southern University at New Orleans (SUNO) students (left to right) Chelsea Brown, Trevor McIntosh, Raven Williams, and Dulaine Vining with their biology professor, Murty Kambhampati (third from left), and Brookhaven Lab mentor Timothy Green (third from right) of the Environmental Protection Division. Kambhampati has been bringing his undergraduate students to Brookhaven Lab to conduct field research in natural resource management. The Brookhaven Lab-SUNO collaboration began after Hurricane Katrina devastated New Orleans in 2005.

a 12-year period, with most students going on to advanced degrees.

By the end of 2017, BNL finalized a Memorandum of Understanding (MOU) with SUNY School of Environmental Science and Forestry (ESF) located in Syracuse, NY. ESF conducts

significant research in the natural sciences including wildlife and forestry. The concept of the MOU would encourage ESF to utilize the BNL site for Pine Barrens-related research. The first meeting to discuss research was held at SUNY-ESF in January 2018 where a wide variety of research efforts were outlined. Several field trips during 2018 introduced faculty members to the Pine Barrens of Long Island and several research proposals were developed. By mid-summer 2018, a proposal to re-visit the 2005-2006 Forest Health Monitoring of the Central Pine Barrens was chosen as the first research effort to be undertaken, and planning began for the research to take place during summer 2019. In the meantime, faculty began planning courses at ESF that would focus on the Pine Barrens starting during the Spring Semester 2019.

In 2018, BNL continued to participate in several events in support of ecological education programs including: providing on-site ecology tours; hosting the Long Island Natural History Conference; participation in the Ninth Annual Pine Barrens Discovery Day held at the Wertheim National Wildlife Refuge; and assisting the Central Pine Barrens Commission with “A Day in the Life of the Rivers,” which allowed students from multiple school districts to acquire environmental and biological data about ten different rivers on Long Island. On separate days, over 30 partner organizations and agencies, over 40 schools, and approximately 3,000 students collected scientific information for analysis to be used to portray the status of the rivers and estuary systems of Long Island. These events provided students hands-on experience with field techniques in catching fish, invertebrate sampling, biodiversity inventory, and water chemistry.

In 2018, BNL entered its 14th year of the Open Space Stewardship Program (OSSP) and worked with 30 schools and nearly 3,000 students. The OSSP enables students to engage in activities to solve problems within their local community through scientific discovery, conservation, and stewardship. The effort integrates outdoor research with school curricula in language arts, civics, community service,

and media arts. Participation in OSSP creates an opportunity for many students to enhance their educational experiences as well as to promote the realization that a career in science and technology is accessible with the proper academic coursework and interaction with teachers and field experts who have a passion for discovery and mentorship.

The Laboratory also hosts the annual New York Wildfire & Incident Management Academy, offered by NYSDEC and the Central Pine Barrens Commission. Using the Incident Command System of wildfire management, this academy trains firefighters in the methods of wildland fire suppression, prescribed fire, and fire analysis. BNL has developed and is implementing a Wildland Fire Management Plan that includes the use of prescribed fire for fuel and forest management and worked with NYSDEC to conduct a single growing season fire in the northern and eastern sections of the BNL property. This successful fire provided significant experience and training for fire crews working in Pine Barrens habitat, improving capabilities of these crews to conduct and manage fire within the Long Island Central Pine Barrens.

6.6 CULTURAL RESOURCE ACTIVITIES

The BNL Cultural Resource Management (CRM) Program ensures that the Laboratory fully complies with numerous cultural resource regulations. The Cultural Resource Management Plan for Brookhaven National Laboratory (BNL 2013a) guides the management for all the Laboratory’s historical resources. BNL’s cultural resources include buildings and structures, World War I (WWI) earthwork features, the Camp Upton Historical Collection, scientific equipment, photo/audio/video archives, and institutional records. As various cultural resources are identified, plans for their long-term stewardship are developed and implemented. Achieving these goals will ensure that the contributions BNL and the site have made to local and national history and culture are documented and available for interpretation.

The Laboratory has four structures or sites that have been determined to be eligible for listing on the National Register of Historic

Places: the Brookhaven Graphite Research Reactor complex, the High Flux Beam Reactor complex, the 1960s-era efficiency apartments, and the WWI training trenches associated with Camp Upton. The trenches are examples of the few surviving WWI earthworks in the United States. Two buildings, Berkner Hall and Chemistry, are architecturally significant. Other buildings have been identified as being important as examples of periods within the history of the BNL site, such as the Brookhaven Center (built during 1930s Civilian Conservation Corp era) and Building 120 (a former WWII-era barracks buildings).

The year 2018 was a quiet for cultural resource efforts with most activity being focused on the 1960s-era efficiency apartments and cement roads that were established as part of the WWI-era Camp Upton. Since the efficiency apartments are likely to be demolished with development of Discovery Park, the New York State Historic Preservation Officer asked that a formal recordation of apartments be accomplished. The recordation was completed and submitted to the New York State Historic Preservation Office (NYSHPO) which resulted in a request to develop an MOU to meet preservation or mitigation requirements under the National Historic Preservation Act. A draft MOU was developed for submittal to the NYSHPO in 2019.

During the summer of 2018, a request to pave the cement road comprising Fifth Street

located east of the National Synchrotron Light Source II was received triggering a review under Section 106 of the National Historic Preservation Act since the road dated to the period of the original Camp Upton established in WWI. This road, along with several other cement roads on site, were evaluated for their historic significance but were found not to be eligible for listing on the National Register of Historic Places since they could not definitively be linked to the period of 1917-1918 and may have been constructed after WWI, just before Camp Upton was decommissioned.

REFERENCES AND BIBLIOGRAPHY

- BNL. 2013. Cultural Resource Management Plan for Brookhaven National Laboratory. BNL-100708-2013. Brookhaven National Laboratory, Upton, NY. May 2013.
- BNL. 2016a. Natural Resource Management Plan for Brookhaven National Laboratory. BNL-112669-2016. Brookhaven National Laboratory, Upton, NY.
- BNL, 2016b. Five-Year Review Report for Brookhaven National Laboratory Superfund Site, April 2016.
- Dwyer, Norval. 1966. Brookhaven National Laboratory. Long Island Forum (reprint), West Islip, NY.
- IAEA. 1992. Effects of Ionizing Radiation on Plants and Animals at Levels Implied by Current Radiation Protection Standards. Technical Report Series No. 332. International Atomic Energy Agency, Vienna.
- LMS. 1995. Phase II Sitewide Biological Inventory Report, Final. Lawler, Matusky & Skelly Engineers. Pearl River, NY.
- NYSDOH. 1999. Deer Meat Contaminated with Cesium-137 at Brookhaven National Laboratory. Bureau of Environmental Radiation Protection, New York State Department of Health, Albany, NY.
- Rispoli, Fred J., Green Timothy, Fasano, Thomas A., Shah, Vishal, 2014. The effect of environmental remediation on the cesium-137 levels in white-tailed deer. Environmental Science and Pollution Research Oct. 2014, 21(19): 11598-11602.