

Abstract No. scar532

XAS Studies of Coordination Complexes Modeling the Structure and Reactivity of Lipoxygenases

R. Scarrow (Haverford College), C. McKenzie, F. Larsen (University of Southern Denmark, Odense)

Beamlines: X18b and (in previous years) X9b

Introduction: Lipoxygenases catalyze the reaction of polyunsaturated fatty acids with molecular oxygen to give fatty acid hydroperoxides. The generally accepted mechanism involves H-atom abstraction from the fatty acid by an iron(III)-hydroxide group. Earlier EXAFS and XANES studies, performed at beam line X9 at the NSLS (reference below), supported this mechanism by showing the presence of a ca. 1.9Å Fe(III)-O bond in the active form of soybean lipoxygenase-1; this distance is shorter than expected for bonds to any amino acid sidechains at the active site of lipoxygenase, but is expected for a hydroxide ligand. Crystallographic studies have revealed that the other ligands to the iron include three or four (depending on species) histidine sidechains and the carboxylate group from the C-terminus of the polypeptide.

Methods and Materials: We are now engaged in a study of the spectroscopy and reactivity of inorganic complexes with ligands that mimic the active site of lipoxygenases. One such ligand is N'-benzyl-N,N'-bis(2-pyridylmethyl)-ethylenediamine N-acetic acid; because it can coordinate to iron with four nitrogen and one oxygen atoms, we denote this ligand (in its neutral acid form) as HL^{N4O}. When the iron(II) complex of this ligand (FeL^{N4O}Cl) is reacted with *t*-butylhydroperoxide (*t*BuOOH) in a methanol/water solvent system, a mononuclear iron(III) complex is formed in solution; ESR spectroscopy and electrospray mass spectrometry both support the formulation of this species as [FeL^{N4O}OH]⁺. To further confirm this formulation and compare the iron coordination in this species to that in soybean lipoxygenase-1, we reacted ca. 10 mM FeL^{N4O}Cl (in a 4:1 MeOH/H₂O solvent system) with a stoichiometric excess of *t*BuOOH at room temperature. The sample was frozen in approximately 1 minute after mixing, and X-ray absorption spectra were obtained using a Si111 monochromator, a Canberra 13-element solid state fluorescence detector and a helium displax cryostat to maintain the sample at ca. 100 K.

Results: The Fe K-edge spectra show about a 2 eV positive shift in edge energy caused by treatment of FeL^{N4O}Cl with *t*BuOOH, confirming the change of oxidation state of the iron from +2 to +3. The figure below shows that the EXAFS spectrum of the frozen solution of the oxidized compound is very similar to that obtained previously of the active form of soybean lipoxygenase-1. Also, the observed spectra rule out the formation of a μ -oxo diferric species under the millimolar solution concentrations used in the EXAFS experiment, since a different spectrum (analyzing for a shorter Fe-O bond length) is obtained on a solid state sample of the dimeric species.

Conclusions: The X-ray absorption spectroscopy supports the results from EPR and mass spectrometry that the oxidation product is [FeL^{N4O}OH]⁺. This system models the formation of the active form of lipoxygenase by treatment of the reduced (iron(+2)) form of the enzyme with lipid hydroperoxides.

Acknowledgments: The detectors and cryostat are owned and maintained by the X18B-X19A PRT and the Fe foil used for energy calibration was supplied by the NSLS equipment pool. We are grateful to Syed Khalid for help in configuring the X18b beam line for our experiments, and to the General User program of the NSLS (proposal # 4294) for providing the beam time.

Reference: R. C. Scarrow, M. G. Trimitsis, C. P. Buck, G. N. Grove, R. A. Cowling, and M. J. Nelson, "X-ray Spectroscopy of the Iron Site in Soybean Lipoxygenase-1: Changes in Coordination upon Oxidation or Addition of Methanol," *Biochemistry*, **1994**, 33, 15023-15035.

