

BNL- 75820-2006-CP

*Unifying the Mechanisms for Single Spin
Asymmetries in Hard Processes*

Feng Yuan

Physics Department/RIKEN BNL Research Center
Brookhaven National Laboratory, Upton, NY, 11973, USA

Presented at the XIV International Workshop on Deep Inelastic Scattering,
DIS 2006
Tsukuba, Japan
April 20-24, 2006

Physics Department/RIKEN BNL Research Center

Brookhaven National Laboratory

P.O. Box 5000
Upton, NY 11973-5000
www.bnl.gov


Managed by
Brookhaven Science Associates, LLC
for the United States Department of Energy under
Contract No. DE-AC02-98CH10886

Notice: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors or their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of author's expresses herein do not necessarily state to reflect those of the United States Government or any agency thereof.



Printed on recycled paper

UNIFYING THE MECHANISMS FOR SINGLE SPIN ASYMMETRIES IN HARD PROCESSES

FENG YUAN

*RIKEN BNL Research Center, Building 510A, Brookhaven National Laboratory,
Upton, NY 11973*

By studying the single transverse-spin asymmetry at the intermediate transverse momentum region in hard processes, Drell-Yan and SIDIS, we demonstrated that the two mechanisms proposed to explain the large SSAs are unified.

Single transverse-spin asymmetry (SSA) in high energy hadronic scattering has a long history. The size of the observed asymmetries came as a surprise and has posed a challenge for researchers in this field¹. Two mechanisms have been proposed in QCD to explain the the large size of SSAs: One is the so-called (naive) time-reversal-odd (T-odd) and transverse-momentum dependent (TMD) parton distributions²; and the other follows the collinear QCD factorization approach and presents the SSAs in terms of spin-dependent twist-three quark-gluon correlation functions (ETQS mechanism)^{3,4}.

In our recent publications⁵, we demonstrated, at the first time, that these two mechanisms are unified, by studying the SSAs at intermediate transverse momentum in semi-inclusive DIS (SIDIS) and Drell-Yan processes. In both processes, At large $q_{\perp} \sim Q$, the ETQS mechanism applies, and the resulting SSA is of twist-three nature. At small $q_{\perp} \ll Q$, a factorization in terms of TMD parton distribution applies⁶, involving in case of the SSA the Sivers functions. If q_{\perp} is much larger than Λ_{QCD} , the dependence of these functions on transverse momentum may be computed using QCD perturbation theory. At the same time, the result obtained within the ETQS formalism may also be extrapolated into the regime $\Lambda_{\text{QCD}} \ll q_{\perp} \ll Q$, and we demonstrated that the result of this extrapolation is identical to that obtained using the TMD approach⁵. In this sense, we have unified the two mechanisms widely held responsible for the observed SSAs.

In this talk, I will present the result for SIDIS. In single transverse-spin polarized deep inelastic scattering, the relevant leptonic tensor is defined as $L^{\mu\nu}(\ell, q) = 2(\ell^\mu \ell^\nu + \ell^\mu \ell'^\nu - g^{\mu\nu} Q^2/2)$, and the hadronic tensor $W^{\mu\nu}$ can be decomposed into $W^{\mu\nu} = \sum_{i=1}^5 V_i^{\mu\nu} W_i$ ⁷. In this study, are primarily interested in hadron production in an intermediate transverse momentum region, $\Lambda_{\text{QCD}} \ll P_{h\perp} \ll Q$, and we will investigate the limit $P_{h\perp} \ll Q$ of the ETQS result. In that limit, V_1 alone provides the leading behavior, so that all other terms than V_1 will be neglected in the following discussions.

After summing over the contributions from all diagrams, we find the transverse-spin dependent cross section can be written as, in the limit of $\Lambda_{\text{QCD}} \ll P_{h\perp} \ll Q$,

$$\frac{d\Delta\sigma(S_\perp)}{dx_B dy dz_h d^2\vec{P}_{h\perp}} = -\frac{4\pi\alpha_{\text{em}}^2 S_{ep}}{Q^4} \epsilon^{\alpha\beta} S_\perp^\alpha \frac{z_h P_{h\perp}^\beta}{(\vec{P}_{h\perp}^2)^2} \frac{\alpha_s}{2\pi^2} \int \frac{dx dz}{xz} \hat{q}(z) \times \left\{ \delta(\hat{\xi} - 1)A + \delta(\xi - 1)B \right\}, \quad (1)$$

where

$$A = \frac{1}{2N_C} \left\{ \left[x \frac{\partial}{\partial x} T_F(x, x) \right] (1 + \xi^2) + T_F(x, x - \hat{x}_g) \frac{1 + \xi}{(1 - \xi)_+} + T_F(x, x) \frac{(1 - \xi)^2 (2\xi + 1) - 2}{(1 - \xi)_+} \right\} + C_F T_F(x, x - \hat{x}_g) \frac{1 + \xi}{(1 - \xi)_+}, \quad (2)$$

$$B = C_F T_F(x, x) \left[\frac{1 + \hat{\xi}^2}{(1 - \hat{\xi})_+} + 2\delta(\hat{\xi} - 1) \ln \frac{z_h^2 Q^2}{\vec{P}_{h\perp}^2} \right], \quad (3)$$

with $\hat{x}_g \equiv (1 - \xi)x = x - x_B$.

On the other hand, when $P_{h\perp} \ll Q$, we know that a transverse-momentum-dependent factorization applies ⁶. Following this reference, the differential SIDIS cross section may be written as

$$\frac{d\sigma}{dx_B dy dz_h d^2\vec{P}_{h\perp}} = \frac{4\pi\alpha_{\text{em}}^2 S_{ep}}{Q^4} (1 - y + \frac{y^2}{2}) x_B \sin(\phi_h - \phi_S) |S_\perp| F_{UT}^{(1)}, \quad (4)$$

where ϕ_S and ϕ_h are the azimuthal angles of the proton's transverse polarization vector and of the transverse momentum vector of the final-state hadron, respectively. $F_{UT}^{(1)}$ has the following factorized form ⁶:

$$F_{UT}^{(1)} = \sum_{q=u,d,s,\dots} e_q^2 \int d^2\vec{k}_\perp d^2\vec{p}_\perp d^2\vec{\lambda}_\perp \vec{k}_\perp \cdot \hat{\vec{P}}_{h\perp} q_T(x_B, k_\perp) / M_P \times \hat{q}(z_h, p_\perp) \left(S(\vec{\lambda}_\perp) \right)^{-1} H_{UT}^{(1)}(Q^2) \delta^{(2)}(z_h \vec{k}_\perp + \vec{p}_\perp + \vec{\lambda}_\perp - \vec{P}_{h\perp}), \quad (5)$$




Figure 1. The factorization arguments for the consistency between the two mechanisms: left is a generic Feynman diagram in the twist-three quark-gluon correlation formalism; and the right is the corresponding TMD factorization form decomposed into different regions, (a) the Siverson function, (b) the fragmentation function, and (c) the soft factor.

where $\hat{P}_{h\perp}$ is a unit vector in direction of $\vec{P}_{h\perp}$, and \hat{q} and q_T are unpolarized quark fragmentation function and the Siverson TMD quark distribution, respectively. We can compute the various factors in the factorization formulas(5) at large transverse momentum ($P_{h\perp} \gg \Lambda_{\text{QCD}}$). The unpolarized quark fragmentation functions is expressed in terms of the k_{\perp} -integrated one,

$$\hat{q}(z_h, p_{\perp}) = \frac{\alpha_s}{2\pi^2} \frac{1}{\vec{p}_{\perp}^2} C_F \int \frac{dz}{z} \hat{q}(z) \left[\frac{1 + \hat{\xi}^2}{(1 - \hat{\xi})_+} + \delta(\hat{\xi} - 1) \left(\ln \frac{\hat{\xi}^2}{\vec{p}_{\perp}^2} - 1 \right) \right],$$

where $\hat{q}(z)$ is the integrated quark fragmentation function and $\hat{\xi} = z_h/z$.

Similarly, the Siverson function at large k_{\perp} can also be calculated perturbatively. Because it is (naively) time-reversal-odd, the only contribution comes from the twist-three quark-gluon correlation function T_F . Carrying out the calculations accordingly, we find ⁵

$$q_T(x_B, k_{\perp}) = \frac{-\alpha_s}{4\pi^2} \frac{2M_P}{(\vec{k}_{\perp}^2)^2} \int \frac{dx}{x} \left\{ A + C_F T_F(x, x) \delta(\xi - 1) \left(\ln \frac{x_B^2 \zeta^2}{\vec{k}_{\perp}^2} - 1 \right) \right\}$$

where A has been defined in Eq. (3) and where $\xi = x_B/x$. The Siverson function for the Drell-Yan process can be calculated similarly, which is found to be the same with an opposite sign⁵, as expected from its definition⁸. This sign difference comes from the different directions of the gauge links in the two processes: in DIS the gauge link arises from final-state interactions and runs to positive light-cone infinity, while in Drell-Yan it is due to initial-state interactions and goes to $-\infty$.

In order to calculate the explicit $P_{h\perp}$ -dependence generated by the TMD factorization, we let one of the transverse momenta \vec{k}_{\perp} , \vec{p}_{\perp} , and $\vec{\lambda}_{\perp}$ be of the

order of $\vec{P}_{h\perp}$ and the others much smaller. When $\vec{\lambda}_\perp$ is large, for example, we neglect \vec{k}_\perp and \vec{p}_\perp in the delta function, and the integrations over these momenta yield either the ordinary quark distribution, or a k_\perp moment of the Sivers function. The latter is related to the twist-three correlation ⁹:

$$\int d^2\vec{k}_\perp q(x, k_\perp) = q(x), \quad \int d^2\vec{k}_\perp \vec{k}_\perp^2 q_T(k_\perp, x)/M_P = -T_F(x, x). \quad (6)$$

In case $\vec{\lambda}_\perp$ is neglected in the delta function, one makes use of the relation ⁶ $\int d^2\vec{\lambda}_\perp S(\lambda_\perp) = 1$. Substituting the above equations into the factorization formula (5), it is easy to see that we reproduce the twist-three result in Eq. (3). This consistency can also be illustrated from the diagrams in Fig. 1, where we showed a generic twist-three Feynman diagram can be factorized into various factors in the TMD factorization formula (5), according to different momentum region of the radiated gluon: parallel to the polarized proton or the final state observed hadron, or soft.

F. Y. is grateful to RIKEN, Brookhaven National Laboratory and the U.S. Department of Energy (contract number DE-AC02-98CH10886) for providing the facilities essential for the completion of their work.

References

1. M. Anselmino, A. Efremov and E. Leader, Phys. Rept. **261**, 1 (1995) [Erratum-ibid. **281**, 399 (1997)]; Z. t. Liang and C. Boros, Int. J. Mod. Phys. A **15**, 927 (2000); V. Barone, A. Drago and P. G. Ratcliffe, Phys. Rept. **359**, 1 (2002).
2. D. W. Sivers, Phys. Rev. D **41**, 83 (1990); Phys. Rev. D **43**, 261 (1991).
3. A. V. Efremov and O. V. Teryaev, Sov. J. Nucl. Phys. **36**, 140 (1982) [Yad. Fiz. **36**, 242 (1982)]; Phys. Lett. B **150**, 383 (1985).
4. J.W. Qiu and G. Sterman, Phys. Rev. Lett. **67**, 2264 (1991); Nucl. Phys. B **378**, 52 (1992); Phys. Rev. D **59**, 014004 (1999).
5. X. Ji, J. W. Qiu, W. Vogelsang and F. Yuan, arXiv:hep-ph/0602239; arXiv:hep-ph/0604023; arXiv:hep-ph/0604128.
6. X. Ji, J. P. Ma and F. Yuan, Phys. Rev. D **71**, 034005 (2005); Phys. Lett. B **597**, 299 (2004); J. C. Collins and A. Metz, Phys. Rev. Lett. **93**, 252001 (2004).
7. R. b. Meng, F. I. Olness and D. E. Soper, Nucl. Phys. B **371**, 79 (1992); Y. Koike and J. Nagashima, Nucl. Phys. B **660**, 269 (2003).
8. S. J. Brodsky, D. S. Hwang and I. Schmidt, Phys. Lett. B **530**, 99 (2002); Nucl. Phys. B **642**, 344 (2002); J. C. Collins, Phys. Lett. B **536**, 43 (2002); X. Ji and F. Yuan, Phys. Lett. B **543**, 66 (2002); A. V. Belitsky, X. Ji and F. Yuan, Nucl. Phys. B **656**, 165 (2003).
9. D. Boer, P. J. Mulders and F. Pijlman, Nucl. Phys. B **667**, 201 (2003); J. P. Ma and Q. Wang, Eur. Phys. J. C **37**, 293 (2004).