



**BNL-XXXXX-YYYY-CP**

**New ENDF/B-VII.0 evaluations of neutron cross sections  
for 32 fission products**

Hyeong Il Kim<sup>1;a</sup>, Young-Ouk Lee<sup>1</sup>, Mike Herman<sup>2</sup>, Said F. Mughabghab<sup>2</sup>,  
Pavel Oblozinsky<sup>2</sup>, and Dimitri Rochman<sup>2</sup>

<sup>1</sup> Nuclear Data Evaluation Laboratory, Korea Atomic Energy Research Institute, Daejeon 305-353, Korea

<sup>2</sup> National Nuclear Data Center, Brookhaven National Laboratory, NY 11973-5000, USA

Energy Sciences & Technology Department  
National Nuclear Data Center  
**Brookhaven National Laboratory**  
P.O. Box 5000  
Upton, NY 11973-5000  
[www.bnl.gov](http://www.bnl.gov)

Notice: This manuscript has been co-authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

# New ENDF/B-VII.0 evaluations of neutron cross sections for 32 fission products

Hyeon Il Kim<sup>1,a</sup>, Young-Ouk Lee<sup>1</sup>, Mike Herman<sup>2</sup>, Said F. Mughabghab<sup>2</sup>, Pavel Obložinský<sup>2</sup>, and Dimitri Rochman<sup>2</sup>

<sup>1</sup> Nuclear Data Evaluation Laboratory, Korea Atomic Energy Research Institute, Daejeon 305-353, Korea

<sup>2</sup> National Nuclear Data Center, Brookhaven National Laboratory, NY 11973-5000, USA

**Abstract.** Neutron cross sections for fission products play important role not only in the design of extended burn-up core and fast reactors, but also in the study of the backend fuel cycle and the criticality analysis of spent fuel. New evaluations in both the resonance and fast neutron regions were performed by the KAERI-BNL collaboration for 32 fission products. These were <sup>95</sup>Mo, <sup>101</sup>Ru, <sup>103</sup>Rh, <sup>105</sup>Pd, <sup>109</sup>Ag, <sup>131</sup>Xe, <sup>133</sup>Cs, <sup>141</sup>Pr, and complete isotope chains of <sup>142-148,150</sup>Nd, <sup>144,147,148-154</sup>Sm, and <sup>156,158,160-164</sup>Dy. The evaluations cover a large amount of reaction channels, including all those needed for neutronics calculations. Also, they cover the entire energy range, from 10<sup>-5</sup> eV to 20 MeV, including the thermal, resolved, and unresolved resonance regions, and the fast neutron region.

## 1 Introduction

Neutron induced nuclear data for 32 fission products were evaluated under the KAERI-BNL collaboration. All evaluations were adopted by the new U.S. Evaluated Nuclear Data Library, ENDF/B-VII.0, released in December 2006 [1]. The list of fission products consists of the priority materials for several applications, extended to cover complete isotopic chains for three elements. Table 1 shows the full list of fission products evaluated in this work.

**Table 1.** Fission products evaluated in this work. Shown in **bold** are priority materials.

	Nuclides
Individual nuclides	<sup>95</sup> Mo, <sup>101</sup> <b>Ru</b> , <sup>103</sup> <b>Rh</b> , <sup>105</sup> <b>Pd</b> , <sup>109</sup> <b>Ag</b> , <sup>131</sup> <b>Xe</b> , <sup>133</sup> <b>Cs</b> , <sup>141</sup> <b>Pr</b>
Neodymium	<sup>142,143,144,145,146,147,148,150</sup> <b>Nd</b>
Samarium	<sup>144,147,148,149,150,151,152,153,154</sup> <b>Sm</b>
Dysprosium	<sup>156,158,160,161,162,163,164</sup> <b>Dy</b>

Our evaluation methodology covers both the low energy region and the fast neutron region. In the low energy region, our evaluations are based on the latest data published in the Atlas of Neutron Resonances [2]. This resource was used to infer both the thermal values and the resolved resonance parameters that were validated against the capture resonance integrals. In the unresolved resonance region we performed the additional evaluation by using the averages of the resolved resonances and adjusting them to the experimental data.

In the fast neutron region, our evaluations are based on the nuclear reaction model code EMPIRE-2.19 [3] validated against the experimental data. EMPIRE is the modular system of codes consisting of many nuclear reaction models, including the spherical and deformed optical model, Hauser-Feshbach theory with the width fluctuation correction and

complete  $\gamma$ -ray emission cascade, DWBA, Multi-step Direct and Multi-step Compound models, and several versions of the phenomenological preequilibrium models. The code is equipped with a powerful GUI, allowing an easy access to support libraries such as RIPL-2 [4] and CSISRS/EXFOR [5], the graphical package, as well the utility codes for formatting and checking.

In general, we used the Reference Input Parameter Library, RIPL-2, for the initial set model parameters. These parameters were properly adjusted to reproduce the available experimental data taken from the CSISRS/EXFOR library.

## 2 Atlas-EMPIRE Evaluation Methodology


Our evaluations are based on the Atlas-EMPIRE methodology developed over several recent years by the National Nuclear Data Center, BNL with several external collaborators. This methodology covers both the low-energy region (thermal energy, resolved resonances, unresolved resonances) and the fast neutron energy region. In the the low-energy region, the evaluations are based on Mughabghab's new Atlas of Neutron Resonances. In the fast-neutron energy region, the evaluations are based on the nuclear reaction model code system EMPIRE developed by Herman *et al.*.

### 2.1 Low-energy region


A statistical analysis of neutron resonances was used by S. Mughabghab, BNL, to produce the well known BNL-325. Its 5<sup>th</sup> edition was published in 2006 as the "Atlas of Neutron Resonances: Resonance Parameters and Thermal Cross Sections, Z = 1 - 100" [2], representing a considerable update to the 1981 [6] and 1984 [7] editions of BNL-325. These latest thermal values and resonance parameters provided a basis for more than 150 new evaluations included in ENDF/B-VII.0, with the resolved and unresolved resonance parameters adopted from Ref. [2]. Figure 1 shows the evaluated capture cross sections for <sup>133</sup>Cs in thermal and resolved resonance

<sup>a</sup> Presenting author, e-mail: hikim@kaeri.re.kr

regions and compares them with the available experimental data.



**Fig. 1.** Neutron capture cross section for  $^{133}\text{Cs}$  in the thermal and resolved resonance region.



**Fig. 2.** Neutron capture cross section for  $^{141}\text{Pr}$  in the unresolved resonance region extended up to the first excited level.

## 2.2 Fast neutron region

All evaluations in the fast neutron region were performed with the new EMPIRE code [3] that has been used for the first time to provide a number of consistent, complete evaluations to the evaluated nuclear data library. EMPIRE calculates cross

sections for all relevant reaction channels, angular distributions, exclusive and inclusive particle- and  $\gamma$ -spectra, double-differential cross sections, and spectra of recoils. The code observes angular momentum coupling (at least in the statistical decay part) and is, therefore, capable of detailed modeling of the  $\gamma$ -cascade providing  $\gamma$  production spectra, intensities of discrete transitions, and isomeric cross sections.

The evaluation in the fast neutron region was merged with the low energy region at 1<sup>st</sup> excited level as shown in Fig. 2. In order to match two regions in a smooth way, we usually adjusted the scattering radius,  $R'$ , and inserted negative cross sections background on rare occasions. Table 2 shows the boundary energy between the unresolved resonance region and the fast neutron region.

**Table 2.** The boundary energies between the unresolved resonance region and the fast neutron region for 32 fission products.

Isotopes	URR limit keV	Isotopes	URR limit keV
$^{95}\text{Mo}$	206.3	$^{144}\text{Sm}$	540.0
$^{101}\text{Ru}$	128.4	$^{147}\text{Sm}$	122.05
$^{103}\text{Rh}$	40.1	$^{148}\text{Sm}$	427.0
$^{105}\text{Pd}$	283.2	$^{149}\text{Sm}$	22.66
$^{109}\text{Ag}$	88.8	$^{150}\text{Sm}$	336.11
$^{131}\text{Xe}$	80.8	$^{151}\text{Sm}$	66.26
$^{133}\text{Cs}$	81.6	$^{152}\text{Sm}$	122.59
$^{141}\text{Pr}$	146.5	$^{153}\text{Sm}$	6.5
$^{142}\text{Nd}$	200.0	$^{154}\text{Sm}$	82.52
$^{143}\text{Nd}$	225.0	$^{156}\text{Dy}$	138.72
$^{144}\text{Nd}$	250.0	$^{158}\text{Dy}$	99.55
$^{145}\text{Nd}$	67.69	$^{160}\text{Dy}$	87.34
$^{146}\text{Nd}$	456.97	$^{161}\text{Dy}$	25.81
$^{147}\text{Nd}$	50.3	$^{162}\text{Dy}$	81.16
$^{148}\text{Nd}$	300.0	$^{163}\text{Dy}$	73.90
$^{150}\text{Nd}$	130.97	$^{164}\text{Dy}$	73.84

## 2.3 Evaluation of complete isotopic chains

A new feature of our evaluation work is a simultaneous evaluation of the complete isotopic chains for a given element. This is possible due to tremendous progress in the development of evaluation tools in recent years. The combined capabilities of the Atlas of Neutron Resonances [2], the nuclear reaction model code EMPIRE [3], input parameter libraries such as the Reference Input Parameter Library, RIPL-2 [4], experimental cross section library CSISRS/EXFOR [5], and numerous ENDF formatting and checking utilities has facilitated such a complex evaluation work. A considerable advantage of this approach is a full and consistent utilization of data that are often measured on natural elements rather than isotopes, a consistent application of model parameters, and comparison with data by summing up isotopic evaluations into a single elemental representation.






Fig. 7. (n,2n) cross sections for all samarium isotopes compared to the measurements.




Fig. 8. Capture cross sections for all dysprosium isotopes compared to the measurements.

## 4 Conclusions

Neutron cross sections for 32 fission products were evaluated in the neutron-incident energy range from  $10^{-5}$  eV to 20 MeV under the KAERI-BNL collaboration. The list of fission products consists of the priority materials for several applications, extended to cover complete isotopic chains for several elements. The full list includes 8 individual isotopes,  $^{95}\text{Mo}$ ,  $^{101}\text{Ru}$ ,  $^{103}\text{Rh}$ ,  $^{105}\text{Pd}$ ,  $^{109}\text{Ag}$ ,  $^{131}\text{Xe}$ ,  $^{133}\text{Cs}$ ,  $^{141}\text{Pr}$ , and 24 isotopes in complete isotopic chains for Nd (8), Sm (9) and Dy (7). The evaluated data were converted into ENDF-6 formatted files, checked by a set of the CSEWG checking codes, processed with the code NJOY-99.161, and subjected to test runs with the code MCNP5 to ensure that the files can be used in transport calculations. All evaluations were adopted by the new U.S. evaluated library, ENDF/B-VII.0, released in December 2006.

This work has been performed under the auspices of the Korean Ministry of Science and Technology as the long-term R&D project.

## References

1. M. Chadwick, P. Obložinský, M. Herman et al., Nuclear Data Sheets **107**(12), 2931 (2006)
2. S.F. Mughabghab, *Atlas of Neutron Resonances: Thermal Cross Sections and Resonance Parameters* (Elsevier, Amsterdam, 2006)
3. M. Herman, R. Capote, B. Carlson, P. Obložinský, M. Sin, A. Trkov, V. Zerkin, *EMPIRE nuclear reaction model code, version 2.19 (Lodi)*, www.nndc.bnl.gov/empire219/ (2005)
4. P. Young, M. Herman, P. Obložinský, T. Belgya, O. Bersillon, R. Capote, T. Fukahori, G. Zhigang, S. Goriely, A. Ignatyuk et al., TECDOC-1506, IAEA, Vienna (2006)
5. NRDC-Network, *CSISRS/EXFOR library of experimental cross sections*, www.nndc.bnl.gov/exfor (2006)
6. S.F. Mughabghab, M. Divadeenam, N.E. Holden, *Neutron Cross Sections: Z=1-60*, Vol. 1A (Academic Press, New York, 1981)
7. S.F. Mughabghab, *Neutron Cross Sections: Z=61-100*, Vol. 1B (Academic Press, New York, 1984)
8. L.R. Veaser, E.D. Arthur, P.G. Young, Physical Review **C16**, 1792 (1977)
9. J. Frehaut, A. Bertin, R. Bois, J. Jary (1980), private communication
10. D.G. Vallis, Tech. Rep. 76/66, A.W.R.E. Aldermaston (1966)
11. H.A. Tewes, A.A. Caretto, A.E. Miller, D.R. Nethaway, Tech. Rep. 6028, U.C., Lawrence Rad.Lab. (Berkeley and Livermore) (1960)
12. F. Voss, K. Wisshak, C. Arlandini, F. Kaeppler, L. Kazakov, T. Rauscher, Physical Review **C59**(2), 1154 (1999)
13. R.B. Taylor, B.J. Allen, A.R.D. Musgrove, R.L. Macklin, Australian Journal of Physics **32**, 551 (1979)
14. J.H. Gibbons, R.L. Macklin, P.D. Miller, J.H. Neiler, Physical Review **122**, 182 (1961)
15. J. Voignier, S. Joly, G. Grenier, Nuclear Science and Engineering **93**, 43 (1986)
16. D.C. Stupegia, M. Schmidt, C.R. Keedy, A.A. Madson, Journal of Nuclear Energy **22**, 267 (1968)
17. G.G. Zaikin, I.A. Korzh, M.V. Pasechnic, N.T. Skljjar, Ukrainskii Fizichnii Zhurnal **16**(7), 1205 (1971)
18. M. Wagner, H. Warhanek, Acta Physica Austriaca **52**, 23 (1980)