

Analytical studies of coherent electron cooling

G. Wang, M. Blaskiewicz, V.N. Litvinenko

Presented at the Particle Accelerator Conference (PAC09)
Vancouver, B.C., Canada
May 4-8, 2009

Collider-Accelerator Department

Brookhaven National Laboratory

P.O. Box 5000
Upton, NY 11973-5000
www.bnl.gov

Notice: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

This preprint is intended for publication in a journal or proceedings. Since changes may be made before publication, it may not be cited or reproduced without the author's permission.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.




Figure 2: Example of electron density modulation in presence of an instantaneous kick at $t = 0$ as calculated from eq. (3). The abscissa is the longitudinal position and the coordinate is the transverse position, both in units of Debye radius. In this example, $\vec{v}_i = (2, 0, 2)$, $\vec{v}_f = (0, 0, 4)$ and $\psi_0 = 2\pi$. (a) Snapshot at $\psi_1 = 0.05\pi$; (b) Snapshot at $\psi_1 = 0.1\pi$; (c) Snapshot at $\psi_1 = 0.2\pi$; (d) Snapshot at $\psi_1 = 0.5\pi$.

$$\tilde{n}_i(\vec{x}, t) = \frac{Z_i}{\pi^2 r_{Dx} r_{Dy} r_{Dz}} \times \left\{ \int_{\psi_1}^{\psi_1 + \psi_0} \frac{\psi \sin(\psi) d\psi}{\left(\omega^2 + (\bar{x} + \bar{v}_{ix} \psi)^2 + (\bar{y} + \bar{v}_{iy} \psi)^2 + (\bar{z} + \bar{v}_{iz} \psi)^2 \right)^{3/2}} + \int_0^{\psi_1} \frac{\psi \sin(\psi) d\psi}{\left(\omega^2 + (\bar{x} + \bar{v}_{ix} \psi)^2 + (\bar{y} + \bar{v}_{iy} \psi)^2 + (\bar{z} + \bar{v}_{iz} \psi)^2 \right)^{3/2}} \right\} \quad (3)$$

, where r_{Dx} , r_{Dy} and r_{Dz} are Debye radius of the electron plasma, ω_p is the plasma frequency of the electrons $\psi_{0,i} = \omega_p T_{0,i}$, $\bar{x} = x / r_{Dx}$ and $\bar{v}_{ix} = v_{ix} / \beta_x$. As shown in Figure 2, direct numerical integration in equation (3) is straightforward and calculation shows that the modulation before merging becomes negligible after 1/4 of plasma oscillations.

CHARGE DENSITY AND PHASE SPACE DENSITY IN 1D FEL




Figure 3: Electron density amplification in 1D FEL. The abscissa is time in units of nanosecond and the coordinate is the electron charge density with units $Z_i e (r_{Dx} r_{Dy} r_{Dz})^{-1}$.

The top graph is the electron density modulation at 1 gain length and the bottom graph is the electron density modulation at 13 gain length. The electron beam going leftwards.

We adopted the formalism in [6] to study the amplification process in an 1D FEL. If we assume the longitudinal energy distribution of the electrons is Lorentzian and the energy spread is small, the electron current density in the frequency domain is given by [7]

$$\tilde{j}_1(\hat{z}, \hat{C}) = \sum_{i=1}^3 A_i(\hat{C}) \lambda_i(\hat{C}) e^{\lambda_i(\hat{C}) \hat{z}} \tilde{j}_1(0, \hat{C}) \quad (4)$$

, where λ_i are three eigenvalues solved from equation

$$\lambda^3 + 2(\hat{q} + i\hat{C})\lambda^2 + \left[\hat{\Lambda}_p^2 + (\hat{q} + i\hat{C})^2 \right] \lambda - i = 0,$$

\hat{q} is parameter to describe the electron energy spread, $\hat{\Lambda}_p$ is the space charge parameter, \hat{C} is the reduced detune, \hat{z} is the longitudinal location along the FEL and A_i are coefficients determined by initial modulation. The time domain solution is thus given by Fourier transformation of equation (4).

Apart from the density modulation, the energy modulation of the electrons in FEL is also important for the effectiveness of the kicker section. For example, preliminary simulation shows that the modulation tends to grow at the beginning of the kicker which is related to the energy modulation in FEL. For Lorentzian energy distribution, the phase space density in the frequency domain is

$$\tilde{j}_1(\hat{C}, \hat{P}, \hat{z}) = -\frac{e\theta_s n_0}{\pi \Gamma \epsilon_0 \rho} \frac{\hat{P} \hat{q}}{(\hat{q}^2 + \hat{P}^2)^2} \sum_{i=1}^3 A_i \frac{1 + i \hat{\Lambda}_p^2 \lambda_i}{\lambda_i + i(\hat{C} + \hat{P})} e^{\lambda_i \hat{z}} \quad (5)$$




Figure 4: Phase space density modulation in FEL. The abscissa is time for 4 radiation period and the coordinate is the energy deviation from $-2\hat{q}$ to $2\hat{q}$. The graph shows the phase space density at 7 gain lengths. The red areas are with higher electron densities and blue areas with lower electron densities. The electron beam going leftwards.

, where \hat{P} is the energy deviation from the average energy of the electron beam, Γ is the gain parameter and θ_s is the electron rotation angle in FEL [6]. The time domain solution is obtained by Fourier transformation of equation (5). As shown in Figure 4, there are more particles losing energy than gaining energy and the net energy lost by the electrons transfers to the radiation field. The whole pattern tilts to the left with a small angle, which is due to the dispersion effects. In the kicker section, the dense area (red) with higher energy and lower energy will approach each other due to the dispersion effect and electron density modulation is thus increased at the beginning of the kicker.

SUMMARY

The analytical solution of electron modulation is obtained for the simplified single kick merger. The result shows that the effects before the kick dissipated in $1/4$ of plasma oscillation. The electron density and phase space density within FEL is derived analytically in the frequency domain. Time domain solutions are obtained by FFT. The results can be used to benchmarking simulation codes and serves as a tool for fast estimation. Further studies involve the analytical study of the kicker section, the diffraction effects in the FEL and the velocity modulation at the modulator.

REFERENCES

- [1] Y. S. Derbenev, in 7th All Union Conference on Charged Particle Accelerators (1980).
- [2] V. N. Litvinenko and Y. S. Derbenev, in 29th International Free Electron Laser Conference (Novosibirsk, Russia, 2007), p. 268.
- [3] V. N. Litvinenko and Y. S. Derbenev, Physical Review Letters **102**, 114801 (2009).
- [4] G. Wang and M. Blaskiewicz, Phys. Rev. E **78**, 026413 (2008).

- [5] George I. Bell, David L. Bruhwiler, Andrey Sobol, et al., in Proceeding of 11th European Particle Accelerator Conference, EPAC' 08 (Genoa, Italy, 2008), p. 3185.
- [6] E. L. Saldin, E. A. Schneidmiller, and M. V. Yurkov, The Physics of Free Electron Lasers (Springer, New York, 1999).
- [7] G. Wang, Ph. D Dissertation, in Department of Physics and Astronomy, Stony Brook University, 2008.