



BNL-93963-2010-CP

**Complete calculation of evaluated
Maxwellian-averaged cross sections and their
uncertainties for s-process nucleosynthesis**

Boris Pritychenko

*National Nuclear Data Center, Brookhaven National Laboratory, Upton, NY 11973-5000, USA
E-mail: pritychenko@bnl.gov*

**National Nuclear Data Center
Energy Sciences & Technology Department
Brookhaven National Laboratory
P.O. Box 5000
Upton, NY 11973-5000
www.bnl.gov**

Notice: This manuscript has been co-authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

This preprint is intended for publication in a journal or proceedings. Since changes may be made before publication, it may not be cited or reproduced without the author's permission.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Complete calculation of evaluated Maxwellian-averaged cross sections and their uncertainties for s-process nucleosynthesis

Boris Pritychenko*

National Nuclear Data Center, Brookhaven National Laboratory, Upton, NY 11973-5000, USA

E-mail: pritychenko@bnl.gov

Present contribution represents a significant improvement of our previous calculation of Maxwellian-averaged cross sections and astrophysical reaction rates. Addition of newly-evaluated neutron reaction libraries, such as ROSFOND and Low-Fidelity Covariance Project, and improvements in data processing techniques allowed us to extend it for entire range of s-process nuclei, calculate Maxwellian-averaged cross section uncertainties for the first time, and provide additional insights on all currently available neutron-induced reaction data. Nuclear reaction calculations using ENDF libraries and current Java technologies will be discussed and new results will be presented.

11th Symposium on Nuclei in the Cosmos, NIC XI

July 19-23, 2010

Heidelberg, Germany

*Speaker.

1. Introduction

Nuclear reactions play an important role in stellar nucleosynthesis and are responsible for producing heavy chemical elements from light elements that were generated in the Big Bang. Present-day calculations of s-process nucleosynthesis are often based on the dedicated nuclear astrophysics data tables, such as work of Bao et al. [1], however it is essential to produce complementary neutron-induced reaction data sets. ENDF-6 formatted evaluated neutron libraries contain various data for all known nuclei, including neutron capture cross sections for more than 680 individual nuclei from ^1H to ^{257}Fm in the range of neutron energy from 10^{-5} eV to 20 MeV. Nuclear-reactor and national-security application communities have used these data extensively in the eV and MeV energy ranges while keV data were less utilized. This creates a unique opportunity to utilize intermediate-energy evaluated neutron data for non-traditional applications, such as s-process nucleosynthesis, and create a new set of ENDF benchmarks. Present work represents a significant upgrade of our previous calculation of Maxwellian-averaged cross sections (MACS) and astrophysical reaction rates [2]. Addition of newly-evaluated neutron reaction libraries, such as JENDL-4, CENDL-3.1, ROSFOND 2010 and Low-Fidelity Covariance Project [3, 4], and improvements in data processing techniques allowed us to extend it for the entire range of s-process nuclei and calculate Maxwellian-averaged cross section uncertainties for the first time. Nuclear reaction calculations using evaluated neutron libraries and current Java technologies will be discussed and new results will be presented.

2. Calculation of Maxwellian-averaged Cross Sections and Uncertainties

ENDF libraries are based on theoretical calculations that are often adjusted to fit experimental data [5]. There are two kinds of ENDF cross section data representations: groupwise (averaged over energy interval) and pointwise. The first kind is often used in reactor physics calculations, while the second one is better suited for nuclear physics applications. Generic ENDF library cross sections (MF=3) do not contain information on neutron resonances. To resolve this problem for neutron physics calculations, codes PREPRO [6] and NJOY [7] are often used to produce a pointwise version of the libraries that include resonance region data and provide cross section information within ENDF range of energies from 10^{-5} eV to 20 MeV. Here, we used code PREPRO to reconstruct resonance region with 10^{-3} precision.

Maxwellian-averaged cross sections (MACS) can be expressed as [2]

$$\sigma^{Maxw}(kT) = \frac{2}{\sqrt{\pi}} \frac{(m_2/(m_1 + m_2))^2}{(kT)^2} \int_0^\infty \sigma(E_n^L) E_n^L e^{-\frac{E_n^L m_2}{kT(m_1 + m_2)}} dE_n^L \quad (2.1)$$

where k and T are the Boltzmann constant and temperature of the system, respectively and E is an energy of relative motion of the neutron with respect to the target. Here E_n^L is a neutron energy in the laboratory system and m_1 and m_2 are masses of a neutron and target nucleus, respectively.

Previously [2], Maxwellian-averaged cross sections and astrophysical reaction rates were produced using the Simpson method on linearized ENDF cross sections (MF=3). This simple method allowed quick calculated integral values with good precision. However the degree of precision was within $\sim 1\%$ [2, 8]. This general limitation can be solved in the linearized ENDF files because

Table 1: Evaluated nuclear libraries and KADONIS Maxwellian-averaged neutron capture cross sections in mb at $kT=30$ keV for s -process nuclei.

Isotope	JENDL-4.0	ROSFOND 2010	ENDF/B-VII.0	KADONIS [13]
36-Kr- 82	9.582E+1	9.483E+1	1.027E+2±3.308E+1	9.000E+1±6.000E+0
42-Mo- 96	1.052E+2	1.035E+2	1.036E+2±3.281E+1	1.120E+2±8.000E+0
44-Ru-100	2.065E+2	2.062E+2	2.035E+2±6.716E+1	2.060E+2±1.300E+1
46-Pd-104	2.700E+2	2.809E+2	2.809E+2±7.551E+1	2.890E+2±2.900E+1
48-Cd-110	2.260E+2	2.346E+2	2.346E+2±7.266E+1	2.370E+2±2.000E+0
50-Sn-116	9.115E+1	1.002E+2	1.002E+2±3.402E+1	9.160E+1±6.000E-1
52-Te-122	2.644E+2	2.639E+2	2.349E+2±8.648E+1	2.950E+2±3.000E+0
52-Te-123	8.138E+2	8.128E+2	8.063E+2±2.857E+2	8.320E+2±8.000E+0
52-Te-124	1.474E+2	1.473E+2	1.351E+2±4.689E+1	1.550E+2±2.000E+0
54-Xe-128	2.582E+2	2.826E+2	2.826E+2±9.991E+1	2.625E+2±3.700E+0
54-Xe-130	1.333E+2	1.518E+2	1.518E+2±4.913E+1	1.320E+2±2.100E+0
56-Ba-134	2.301E+2	2.270E+2	2.270E+2±8.311E+1	1.760E+2±5.600E+0
56-Ba-136	7.071E+1	7.001E+1	7.001E+1±2.389E+1	6.120E+1±2.000E+0
60-Nd-142	3.557E+1	3.701E+1	3.341E+1±1.147E+1	3.500E+1±7.000E-1
62-Sm-148	2.361E+2	2.444E+2	2.449E+2±7.331E+1	2.410E+2±2.000E+0
62-Sm-150	4.217E+2	4.079E+2	4.227E+2±1.452E+2	4.220E+2±4.000E+0
64-Gd-154	9.926E+2	1.010E+3	9.511E+2±2.901E+2	1.028E+3±1.200E+1
66-Dy-160	8.702E+2	8.293E+2	8.328E+2±2.900E+2	8.900E+2±1.200E+1
72-Hf-176	5.930E+2	4.529E+2	4.571E+2±1.459E+2	6.260E+2±1.100E+1
80-Hg-198	1.612E+2	1.612E+2	1.612E+2±5.392E+1	1.730E+2±1.500E+1
82-Pb-204	8.355E+1	7.242E+1	7.242E+1±2.084E+1	8.100E+1±2.300E+0

cross section value is linearly-dependent on energy within a particular bin [9]

$$\sigma(E) = \sigma_1 + (E - E_1) \frac{\sigma(E_2) - \sigma(E_1)}{E_2 - E_1} \quad (2.2)$$

where $\sigma(E_1), E_1$ and $\sigma(E_2), E_2$ are pointwise cross section and energy values for the energy bin. This allowed us to calculate definite integrals using Wolfram Mathematica online integrator [10]. Summing integrals for all energy bins will produce an exact integral value for Maxwellian-averaged cross section. Low-Fidelity cross section covariances were used to calculate uncertainties for ENDF/B-VII.0 data [4, 11]. Final results for JENDL-4.0, ROSFOND 2010 and ENDF/B-VII.0 libraries [12, 3, 11] are shown in Table 1. Due to space limitations only selected data are shown and CENDL-3.1, JEFF-3.1 cross sections are omitted.

In s -process nucleosynthesis, the product of neutron-capture cross section (at 30 keV in mb) times solar system abundances (relative to Si = 10^6) as a function of atomic mass should be constant for equilibrium nuclei [14]:

$$\sigma_A N_A = \sigma_{A-1} N_{A-1} = \text{constant} \quad (2.3)$$

To verify this phenomenon, the calculated $\langle \sigma_\gamma^{Maxw}(30keV) \rangle$ from the ENDF/B-VII.0 library [11] were multiplied by solar abundances taken from Anders and Grevesse [15], and plotted in




Figure 1: ENDF/B-VII.0 product of neutron-capture cross section (at 30 keV in mb) times solar system abundances (relative to Si = 10^6) as a function of atomic mass for nuclei produced only in the *s*-process.

Figure 1. Visual inspection of the Figure indicates two local equilibrium and ledge-precipice break at $A \sim 138$ for the ENDF/B-VII.0 fit. Relatively high product value for ^{116}Sn is due to the fact that ^{116}Sn has *r*-process contribution [15].

In FY 2011, present contribution results will be used to upgrade ‘Maxwellian-averaged Cross Sections and Astrophysical Reaction Rates’ Web application <http://www.nndc.bnl.gov/astro>. The Web application frontpage is shown in Figure 2.

3. Conclusion & Outlook

Maxwellian-averaged cross sections (MACS) for capture and fission have been calculated using JENDL-4, CENDL-3.1, ROSFOND 2010, ENDF/B-VII.0 and JEFF-3.1 libraries using Low-Fidelity covariance project data. ROSFOND 2010 library (686 materials) [3] provides the most complete coverage along the *s*-process path. In fact, it contains information on 347 out of 354 isotopes (98% coverage). The only *s*-process isotopes that are missing in ROSFOND are: ^{12}C , ^{110}Ag , $^{132,133,135,137}\text{Ce}$ and ^{142}Pr . Carbon and Praseodymium cross sections can be found in the CENDL-3.1 and ENDF/B-VII.0 or JEFF-3.1 libraries, respectively. Typical neutron library, such as ENDF/B-VII.0 (393 materials), provides 80% coverage of the *s*-process path.

The comparison of Maxwellian-averaged cross section values from Table 1 indicates a good agreement between KADONIS and evaluated nuclear libraries. Future work on ENDF libraries will provide additional improvements and benefits for nuclear astrophysics and applications communities.

4. Acknowledgments

The author thanks Drs. V. Zerkin (IAEA), M.W. Herman, C.M. Mattoon, S.F. Mughabghab,




Figure 2: ‘Maxwellian-averaged Cross Sections and Astrophysical Reaction Rates’ Web application, <http://www.nndc.bnl.gov/astro>.

and M. Pigni (BNL) for productive discussions and help on ROSFOND and Low-Fidelity project data, respectively. We are also grateful to M. Blennau for a careful reading of the manuscript. This work was sponsored by the Office of Nuclear Physics, Office of Science of the U.S. Department of Energy, under Contract No. DE-AC02-98CH10886 with Brookhaven Science Associates, LLC.

References

- [1] Z.Y. Bao, H. Beer, F. Käppeler *et al.*, *ATOMIC DATA AND NUCLEAR DATA TABLES* **76**, 70 (2000).
- [2] B. Pritychenko, S.F. Mughabghab, and A.A. Sonzogni, *ATOMIC DATA AND NUCLEAR DATA TABLES* (2010), in press; doi:10.1016/j.adt.2010.05.002.
- [3] S.V. Zabrodskaya, A.V. Ignatyuk, V.N. Koscheev *et al.*, *VANT, Nuclear Constants 1-2 3* (2007).
- [4] R.C. Little, T. Kawano, G.D. Hale *et al.*, *NUCLEAR DATA SHEETS* **109** 2828 (2008). Available from <http://www.nndc.bnl.gov/lowfi/>.
- [5] *EXPERIMENTAL NUCLEAR REACTION DATA (EXFOR/CSISRS)*, Available from <http://www.nndc.bnl.gov/exfor>.

- [6] D.E. Cullen, "The ENDF/B Pre-processing codes (PREPRO)," Available from <http://www-nds.iaea.org/ndspub/endf/prepro/>.
- [7] R.E. MacFarlane and D.W. Muir, "The NJOY nuclear data processing system, version 91," Tech. Rep. LA-12740-M, Los Alamos National Lab, 1994.
- [8] T. Nakagawa, S. Chiba, T. Hayakawa, T. Kajino, *ATOMIC DATA AND NUCLEAR DATA TABLES* **91**, 77 (2005).
- [9] B. Pritychenko, *PROCEEDINGS OF 59th CSEWG MEETING*, Upton, NY, USA, November 3-5 (2009), Available from <http://www.nndc.bnl.gov/proceedings/2009csewgusndp/Tuesday/csewg/PritychenkoCSEWG09.pdf>.
- [10] Wolfram Mathematica Online Integrator, Available from <http://integrals.wolfram.com>.
- [11] M.B. Chadwick, P. Obložinský, M.W. Herman *et al.*, *NUCLEAR DATA SHEETS* **107**, 2931 (2006).
- [12] *JAPANESE EVALUATED NUCLEAR DATA LIBRARY (JENDL-4.0)*, Available from <http://www.ndc.jaea.go.jp/index.html>.
- [13] I. Dillmann, M. Heil, F. Käppeler *et al.*, *AIP Conf. Proc.* **819**, 123 (2006). Data downloaded from <http://www.kadonis.org> on April 7, 2010.
- [14] C.E. Rolfs and W.S. Rodney, *CAULDRONS IN THE COSMOS*, The University of Chicago Press (1988).
- [15] E. Anders and N. Grevesse, *GEOCHIMICA ET COSMOCHIMICA ACTA* **53**, 53 (1989).