***CLOUDS, PRECIPITATION AND MARINE
BOUNDARY LAYER STRUCTURE DURING THE
MAGIC FIELD CAMPAIGN***

Xiaoli Zhou¹, Pavlos Kollias¹, and Ernie R. Lewis²

1. Department of Atmospheric and Oceanic Sciences, McGill University,
Montreal, CA

2. Biological, Environmental & Climate Sciences Department, Brookhaven
National Laboratory, Upton, NY

Corresponding author: Xiaoli Zhou
Department of Atmospheric and Oceanic Sciences
Montreal H3A0B9 Quebec, Canada
email: xiaoli.zhou@mail.mcgill.ca

*Accepted for publication in
the Journal of Climate*

April 2014

Biological, Environmental & Climate Science Dept.

Brookhaven National Laboratory

**U.S. Department of Energy
DOE Office of Science**

Notice: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-SC0012704 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Abstract

The recent ship-based MAGIC field campaign with the marine-capable Second ARM Mobile Facility (AMF2) deployed on the Horizon Lines cargo container M/V *Spirit* provided nearly 200 days of intra-seasonal high-resolution observations of clouds, precipitation and marine boundary layer (MBL) structure on multiple legs between Los Angeles, California, and Honolulu, Hawaii. During the deployment, MBL clouds exhibited a much higher frequency of occurrence than other cloud types and occurred more often in the warm season than in the cold season. MBL clouds demonstrated a propensity to produce precipitation, which often evaporated before reaching the ocean surface. The formation of stratocumuli is strongly correlated to a shallow MBL with a strong inversion and a weak transition, while cumuli formation is associated with a much weaker inversion and stronger transition. The estimated inversion strength (EIS) is shown to depend seasonally on the potential temperature at the 700 hPa. The location of the commencement of systematic MBL decoupling (DE) always occurred eastward of the locations of cloud breakup (CBs), and the systematic decoupling showed a strong moisture stratification. The entrainment of the dry warm air above the inversion appears to be the dominant factor triggering the systematic decoupling, while surface latent heat flux, precipitation and solar radiation did not play major roles. MBL clouds broke up over a short spatial region due to the changes in the synoptic conditions, implying that in real atmospheric conditions, the MBL clouds do not have enough time to evolve as is in the idealized models.

48

49

50 **1. INTRODUCTION**

51

52 On average, near 30% of the global oceans are covered with low-level clouds (ISCCP
53 online data set: <http://isccp.giss.nasa.gov/climanal7.html>). These prevailing marine
54 boundary layer (MBL) clouds are a key component in the earth's radiation budget, from
55 which Stratocumulus (Sc) clouds exert a strong negative net radiative effect due to their
56 low height and high areal coverage, by strongly reflecting incoming solar radiation but
57 only weakly influencing the outgoing longwave radiation (Wood, 2006). Cumulus (Cu)
58 have a reduced effect on the radiation due to their low areal coverage (Wood, 2006;
59 Karlsson et al. 2010) but play a critical role in the vertical redistribution of moisture and
60 energy in the lower troposphere (Tiedke et al., 1988). Thus, it is important for global
61 climate models to accurately represent these cloud regimes.

62 The evolution, with increasing sea surface temperature (SST), from Sc regimes to Cu
63 regimes in the trade wind regions and then eventually to congestus and deep convective
64 Cu over the warmer waters of the intertropical convergence zone are well documented
65 (e.g., Albrecht et al., 1995a,b; Karlsson et al., 2010). An inversion layer that is often
66 thought of as the top of the MBL typically caps the Sc. A regime with Cu-under-Sc
67 usually occurs during the progression between Sc and Cu and is associated with a weakly
68 stable layer below the inversion base that is characterized by a sharp decrease of moisture
69 with height (e.g. Krueger et al. 1995; Bretherton and Wyant, 1997; Jones et al. 2011).

This stable layer, referred here as the transition layer, separates a region below of surface flux-driven turbulence from a region above dominated by radiatively driven convection (Bretherton and Wyant, 1997) and acts to isolate the upper MBL from the surface moisture supply. When this vertical moisture stratification gets sufficiently strong, systematic decoupling occurs, and remains decoupled with further increase in SST. This systematic decoupling is a crucial first step in the Sc-to-Cu transition and is not affected by the diurnal cycle of radiation (Wyant et al. 1997).

Due to the lack of fully understanding of the mechanisms responsible for the evolution of MBL structure and cloud, global weather and climate prediction models still do not accurately reproduce the evolution between these cloud regimes, and the locations of Sc breakup and the rates of change of cloud coverage vary widely among different models, which generally underestimate cloud amounts in the Sc region while overestimating them in the Cu region (Teixeira et al., 2011).

One of the main factors hindering progress in representing these clouds in numerical models has been the lack of observational data. Most of the observational data sets used to evaluate the cloud-related processes are satellite-based. Satellite data have proven valuable in determining the climatological links between MBL inversion base height (MBLH) and cloud cover (e.g. Heck et al. 1990; Wang et al. 1993; Wood and Bretherton, 2004). However, satellite observations cannot provide information on detailed vertical cloud and thermodynamic structure of the MBL especially during decoupling conditions (Wood and Bretherton, 2004; Kalsson et al. 2010). The difficulty in accurately observing low-level clouds with small-scale variability (Xu and Cheng, 2013) further restricts the applicability of satellite data in the understanding of the transition between these cloud

regimes.

In addition to satellite data, several field campaigns have been conducted to study MBL clouds and the mechanisms responsible for the Sc-to-Cu transition. These previous ship-based efforts provided a wealth of information with regard to the vertical structure of the MBL and associated clouds. However, they were primarily conducted in a fairly small region and focused on studying one specific cloud type. For example, the First International Satellite Cloud Climatology Project (ISCCP) Regional Experiment (FIRE; Albrecht et al. 1988) focused on Sc and cirrus cloud regimes, the Tropical Instability Wave Experiment (TIWE; Albrecht et al. 1995b) focused on the trade wind Cu boundary layer structure, and during the Atlantic Stratocumulus Transition Experiment (ASTEX; Albrecht et al., 1995a) a transition region in which Cu form beneath Sc was observed. Albrecht et al. (1995b) compared the large-scale forcing and thermodynamic profiles from these three field experiments and concluded that the increase in sea surface temperature (SST) is important in the thinning of the Sc. The same study provided evidence that the boundary layer structure and the associated transition from Sc to Cu may be more complicated than originally thought. More recently, Jones et al. (2011) examined in detail the coupled and decoupled boundary layers in the Variability of the American Monsoon Systems (VAMOS) Ocean-Cloud-Atmosphere-Land Study Regional Experiment (VOCALS-REx; Wood et al., 2011). One of the major findings in Jones et al, 2011 is that the difference between MBLH and the lifting condensation level (LCL) best predicts decoupling.

Previous numerical studies have demonstrated that the systematic decoupling is mainly driven by the increasing surface latent heat flux (LHF) as a response to the increasing

116 entrainment due to the warmer SST (e.g. Bretherton and Wyant, 1997, Sandu and Stevens,
117 2011). The subsequent Sc breakup was explained as a result of the further increase of the
118 SST that causes the Cu to become deeper and more vigorous, penetrating farther into the
119 inversion and entraining more dry air from above the inversion (Wyant et al., 1997;
120 Sandu and Stevens, 2011). Although numerical studies have advanced our knowledge of
121 MBL structure and clouds, simulations have usually simplified the problem by assuming
122 the constant divergence and free-tropospheric lapse rates (Bretherton and Wyant, 1997),
123 but neither assumption is supported by observations.

124 There are noticeable discrepancies between idealized model simulations and
125 observational findings. For instance, the dominant effect of the LHF on MBL decoupling
126 was not observed in VOCAL-REx (Jones et al. 2011). Additionally, the assumption of a
127 constant free-atmosphere lapse rate might introduce biases because MBLH and boundary
128 layer mixing ratios are very sensitive to above-inversion features (Albrecht 1984;
129 Krueger et al. 1995). At the same time, the availability of comprehensive, long-term
130 observations that document the gradual MBL decoupling and Sc-to-Cu transitions is
131 limited.

132 The recent MAGIC field campaign provided high resolution, profiling observations, from
133 the coast of California to Honolulu for over 200 days. The collected dataset is the most
134 extensive direct, long-term, intra-seasonal set of measurements of MBL structure and
135 cloud evolution from Sc to Cu over large downwind regions. Here, we investigate the
136 potential of the dataset in advancing our understanding of the systematic MBL
137 decoupling and Sc breakup to be compared with and constrain the modeling studies.

The remaining of the manuscript is organized as follows. Brief descriptions of the MAGIC field campaign and of the AMF-2 instruments are provided in Section 2, and the methodology used for this study is introduced in Section 3. Results are presented in Section 4, and a summary of these results and plans for future work are presented in Section 5.

2. OBSERVATIONS

2.1 The MAGIC Field Campaign

The MAGIC field campaign (<http://www.arm.gov/sites/amf/mag/>) deployed the US Department of Energy (DOE) Atmospheric Radiation Measurement (ARM) Mobile Facility 2 (AMF2) on the commercial cargo container Horizon *Spirit* (Fig. 1a). The MAGIC transect is very near the line from the coast of California to the equator (35°N, 125°W to 1°S, 173°W) that was chosen by modelers to compare model results for the Global Energy and Water Experiment (GEWEX) Cloud System Studies (GCSS) Pacific Cross-section Intercomparison (GPCI) study, in which more than twenty climate and weather-prediction models participated (Teixeira et al., 2011). Thus, MAGIC with its unprecedented, intra-seasonal, high-resolution ship-based observations is expected to provide constraint, validation, and support for the aforementioned modeling efforts and at the same time contribute in improving our understanding of the Sc-to-Cu transition along this GPCI transect.

From October, 2012 through September, 2013, the *Spirit* completed 20 round trips between Los Angeles, California, and Honolulu, Hawaii. Each trip is called a “leg”, and legs are numbered sequentially as “LegxxA” for the trips from Los Angeles to Honolulu and “LegxxB” for the return trips (Fig. 1b). During the legs from Los Angeles, the *Spirit* traveled at ~21 kts ($\sim 10.5 \text{ m s}^{-1}$) and covered the 4100 km distance in 4.5 days. The *Spirit* returned to Los Angeles at ~16 kts ($\sim 8 \text{ m s}^{-1}$), making the trip in approximately 6.5 days (the lower speed resulting in lower fuel costs and allowing the ship to remain on a two-week schedule; Lewis et al., 2012). The departure and arrival times of MAGIC legs are listed in Appendix A. Two technicians associated with the MAGIC campaign lived on board the *Spirit* during the deployment to maintain the instrumentation, launch radiosondes, and perform other tasks.

2.2 AMF2 instrumentation and data description

The AMF2 contains a state-of-the-art instrumentation suite and was designed to operate in a wide range of climate conditions and locations, including shipboard deployments. The AMF2 was located on the bridge deck of the *Spirit*, approximately 16 m above sea level (ASL). The primary AMF2 instruments used in the current study are 1) a Ka-band ARM Zenith Radar (KAZR), 2) a laser ceilometer, 3) a Vaisala weather station, 4) a inertial navigational location and attitude system (NAV), 5) the Marine Meteorological System (MARMET) installed on the mast of the *Spirit* approximately 27 m above sea level and an Infrared Sea-surface temperature Autonomous Radiometer (ISAR), and 6)

radiosondes (four or eight per day). AMF2 also contained a motion-stabilized W-band radar, a radar wind profiler, a broadband and spectral radiometer suite, aerosol instrumentation, and other instruments. The operational status of all instruments during the campaign is summarized in Appendix B. There were some time periods when the KAZR was not acquiring data for various reasons (e.g., installation, power outages, etc.), but KAZR measurements were obtained for all of Leg03A through Leg08B, Leg11, and Leg14A through Leg17B. The analyses presented here are based on these data, in total 22 transits between California and Hawaii through the Sc-to-Cu transition region comprising more than 3000 hours. The data are separated into two seasons, the warm season: Leg11 and Leg14 to Leg17 (May 25 to June 6, and July 7 to August 29, 2013), and the cold season: Leg03 to Leg08 (October 6 to December 27, 2012).

2.2.1 Ka-band ARM Zenith Radar (KAZR)

The KAZR, formerly known as the millimeter wavelength cloud radar (MMCR, Moran et al., 1998), is a 35-GHz profiling Doppler radar that retrieves information on the vertical distribution of the hydrometeors in the atmospheric column. Due to its short wavelength (8.6 mm), the KAZR has sufficient sensitivity to detect MBL clouds with little attenuation under moderate drizzle conditions. The KAZR might fail to detect very thin liquid clouds and it can provide inaccurate hydrometeor-layer heights during heavy precipitation because of severe radar signal attenuation (Matrosov, 2007), but because a ceilometer was used (Section 2.2.2) and because there was very little heavy precipitation

during MAGIC, these issues should have little effect on the results of this study. The KAZR utilizes a new digital receiver that provides higher temporal (less than 2 s) and spatial (30 m) resolution than the MMCR (Widener et al., 2012). It is unaffected by Bragg scattering and has small antennas with narrow beamwidths as well as limited sidelobes (Kollias et al., 2007). During MAGIC, the KAZR was configured to have temporal resolution of about 0.4 s to oversample the ship motion, thus enabling compensation of the effects of this motion on the radar observables during data post-processing. In this study, all KAZR measurements have been averaged over 4 s, which allows for the detection of small-scale Cu.

2.2.2 Ceilometer

A ceilometer (Vaisala model CT25K) operating at a wavelength of 910 nm was used to detect the base heights of clouds. The ceilometer's range resolution was 10 m, and its temporal resolution was near 16 s for Leg03 and Leg04 and 3 s for the other legs. In order to maintain the 4-s temporal resolution, it is assumed that each reported base height is representative of the entire original time period.

2.2.3 Vaisala Weather station

A Vaisala weather station WXT-520 installed as part of a suite of meteorological instruments associated with the Aerosol Observing System of the AMF2 (AOSMET)

measured rain intensity at 1 s resolution which was used to detect the presence of precipitation reaching the ground (see Section 3c).

2.2.4 Navigational Location and Attitude (NAV)

NAV provided ship location and attitude with a temporal resolution of 1 s during the period between November 3 and December 3, 2012 (Leg05A to Leg07A), and 0.1 s for the rest of the deployment. As all macroscopic data are averaged over 4 s, both temporal resolutions are sufficiently accurate for the present comparisons.

2.2.5 Marine Meteorological Measurement (MARMETX) & Marine Flux (MARFLUX) data sets

The MARMETX data set (<http://www.arm.gov/campaigns/amf2012magic/>) contains standard surface meteorological parameters measured by the MARMET: temperature (T), pressure (P), relative humidity (RH), and apparent and true wind speed and direction; and the sea-surface skin temperature (SSST) measured by the ISAR (with an accuracy of better than 0.1°C). MARFLUX (<http://www.arm.gov/campaigns/amf2012magic/>) contains the surface fluxes of moisture and sensible and latent heat calculated by the TOGA-COARE air-sea flux algorithm (Fairall et al., 1996) using the MARMETX variables. Both MARMETX and MARFLUX have a time resolution of 1 min.

2.2.6 Atmospheric Soundings

250

251 Standard radiosondes (Vaisala model MW-31, SN E50401) were launched every 6 h to
252 measure vertical profiles of the thermodynamic state of the atmosphere (T, P, RH, and
253 wind speed and direction). During Leg14, which occurred in July 2013, launches were
254 made every 3 h to provide a more detailed picture of the atmospheric structure. Only
255 soundings providing measurements as high as 15 km were used in this study (389 in all).
256 The radiosondes collected data every 2 s during their ascent, providing a typical vertical
257 resolution of 10 m in the troposphere. However, owing to the limited launching frequency
258 (four to eight per day), sounding data can be interpolated to higher-resolution time steps
259 with only limited confidence.

260

261

262 **3. METHODOLOGY**

263

264 **3.1 Hydrometeor mask**

265

266 A hydrometer mask was applied to the raw KAZR reflectivity measurements to identify
267 the radar range gates that contain appreciable returns from hydrometeors. Following
268 Rémillard et al. (2012), this mask uses the algorithm of Hildebrand and Sekhon (1974)
269 and a two-dimensional (time-height) filter to identify the number of hydrometeor layers
270 in the atmospheric column and their corresponding boundaries. The lowest hydrometeor
271 boundary is not necessarily the cloud base because the KAZR cannot distinguish cloud
272 drops from precipitation particles below the cloud base.

3.2 Cloud boundaries and cloud classification

To obtain cloud boundaries, the hydrometeor mask is combined with the ceilometer-generated time series of cloud-base height. The ceilometer cloud-base heights are binned to the KAZR spatial resolution, with an uncertainty of less than 15 m (the KAZR has range resolution of 30 m). Examples of the KAZR hydrometeor mask along with first ceilometer-derived cloud-base heights for Leg04A and Leg04B are shown in Fig. 2. The combination of the KAZR and the ceilometer is sufficient to characterize cloud fraction (CF) and layering in the lower troposphere (i.e., $< \sim 3$ km), although hydrometeor occurrence in the upper troposphere is probably underestimated because the maximum operational range of the ceilometer is 5-6 km. However, as the analysis here pertains to MBL clouds, this underestimation should not introduce any biases.

Cloud tops for each hydrometeor layer throughout the troposphere are determined using the KAZR-derived hydrometeor mask. In liquid clouds, KAZR reflectivity measurements below the first cloud-base height determined by the ceilometer are used to characterize precipitation (see next section). If no ceilometer data are available, no KAZR data below 300 m are used, since they often contain artifacts (especially when no precipitation is present). In cases when the first ceilometer-derived base height is 100 m or more less than the first KAZR-defined hydrometeor base height, the two clouds are considered independent, with the first cloud-top height undetermined; otherwise the first KAZR hydrometeor top is considered to be the first cloud top.

295

296 Once the cloud boundaries are determined, each time-height cluster of KAZR echoes
297 with more than 25 connected pixels (in time-height space) is considered to be a cloud
298 entity. In order to obtain realistic bases of multiple-layer cloud entities, the bases of the
299 second cloud level are further smoothed according to the ceilometer-derived base heights.
300 KAZR echoes below the newly defined cloud bases are neglected. Each cloud entity is
301 categorized into one of four types based on its average base and top heights (Table 1, see
302 Fig. 3a as an example): high-level, mid-level, MBL, or cumulus congestus and deep
303 convective. High-level clouds have average base heights of at least 6 km. Mid-level
304 clouds have average base heights between 3 km and 6 km. MBL clouds have average
305 base heights and top heights less than 3 km, or have undetermined average top heights.
306 Cumulus congestus and deep convective clouds have average base heights less than 3 km
307 but average top heights of at least 3 km. The statistical results of cloud properties
308 presented in this study are not sensitive to the specific values chosen for the thresholds.

309 An MBL cloud layer is detected if more than 10% of cloud bases are measured over a
310 continuous range of heights during one hour (a one-gate gap is allowed). The cloud bases
311 here refer to the first and second ceilometer-derived bases and the first three
312 hydrometeor-mask bases.

313 As the focus of this study is MBL clouds, emphasis is placed on these clouds, which are
314 further divided into three subtypes: stratocumulus (Sc), cumulus (Cu), and indeterminate
315 (e.g., Figs. 3b and 3c). Sc are low clouds composed of an ensemble of individual
316 convective elements that together assume a layered form (Wood, 2012), whereas Cu
317 clouds are separate convective elements. The difference between Sc and Cu in this study

is based on their time durations: a cloud is defined as Sc if it lasts more than 20 min, and as Cu if its duration is less than 20 min. Sc clouds are also required to have a narrow cloud top height distribution that is restricted by a specific standard deviation threshold that depends on its duration (see Table 1 for details). The remaining MBL cloud clusters make up the subtype ‘indeterminate’. Because of the limited nature of the MAGIC observations (1D in distance/time and height), these cloud types are not mutually exclusive.

Because of their ship-based origin, the cloud macroscopic data for each leg depend on both time and location (e.g., Figs. 1,2). To account for slight ship-course deviations between different legs (Fig. 1b), all of the cloud macroscopic data are binned to a uniform great-circle route with 40-m resolution (the approximate distance covered by the ship in 4 s) in order to examine the evolution of cloud properties along this representative great-circle transect. Finally, all the cloud macroscopic data are averaged over 36-km (the approximate distance covered by the ship in 1 hr) and converted to the corresponding latitude along the great-circle route. The frequency of occurrence of MBL cloud every 36-km are considered to represent the CF over that area and are referred to as CF_{36} in this paper.

3.3 Precipitation classification

KAZR observations, the KAZR-derived hydrometer mask, ceilometer-derived cloud-base heights, surface rainfall occurrence from the weather stations, and 0°C isotherm heights

derived from interpolated radiosonde data are used to characterize precipitation. KAZR echoes are classified as either cloud or precipitation; no attempt is made to distinguish ice hydrometeors from liquid cloud drops or precipitation. Because the cloud-base heights are determined from the ceilometer data, every KAZR echo below the first cloud-base height is classified as precipitation. The ceilometer quality control flag was checked to ensure that no water was present on the ceilometer window (which would occur in the case of intense precipitation reaching the surface), since this would strongly attenuate the signal and result in inaccurate readings.

Precipitation is classified into five types (Table 2): virga, drizzle, warm rain, cold rain, and deep convective rain (e.g., Fig. 3d). Precipitation that is not detected at the surface is either virga or drizzle. The distinction between the two is based on the detection of KAZR echoes in the lowest range gate (around 120 m ASL before Leg11A, and 240 m ASL for the later legs): virga is defined as precipitation that is detected at least 50 m below the ceilometer cloud base and does not reach the lowest KAZR range gate, whereas drizzle is detected at the KAZR lowest range gate. This distinction provides a qualitative indicator of light rain intensity and indicates the portion of the subcloud layer affected by evaporation. Under this proposed definition only rain that falls through a cloud and not that from the side of a cloud could be considered virga. Precipitation is defined as warm (cold) rain when it is detected at the surface and when the first KAZR-derived cloud top is above 3 km but lower (higher) than the 0°C isotherm height, regardless of the ceilometer-derived cloud-base height. Precipitation is defined as deep convective rain when it is detected at the surface and when the radar reflectivity at the lowest range gate is greater than 0 dBZ and also greater than that at the first KAZR

cloud-top height. Similar to the cloud macroscopic data, the precipitation data for each leg are averaged over 36-km along the great-circle route.

3.4 Radiosonde analysis

Radiosonde data are used to determine the thermodynamic structure of the lower atmosphere. Emphasis in this study is placed on identifying the inversion and transition layers, which are closely associated with MBL stratification and MBL clouds. The inversion layer is defined as all levels around the maximum increase of temperature with height that occurs between 500 m and 3 km that have an increase of temperature and a decrease in water vapor mixing ratio (r) with height (see Rémillard et al., 2012 for more details).

Following the algorithm of Yin and Albrecht (2000), a parameter μ is defined for each level below the inversion base in terms of the potential temperature (θ) and r :

$$\mu = -\left(\frac{\partial \theta}{\partial P} - \frac{0.608\theta}{1+0.608r} \frac{\partial r}{\partial P}\right), \quad (1)$$

and the transition layer is defined whenever the maximum value of μ is positive and consists of all levels below the base of the inversion at which μ is greater than 1.3 times its mean value over the entire region below the inversion. Examples of the sounding-derived inversion and transition layers under conditions of broken Cu and overcast Sc are illustrated in Figs. 4a and 4b, respectively.

383 The sounding data for each vertical profile are smoothed prior to any analysis using only
 384 the next-nearest points to remove small-scale variability and to provide smooth local
 385 gradients. To preserve the features of the transition and inversion layers, a layer-by-layer
 386 averaging procedure is performed when averaging various profiles together (Augstein et
 387 al., 1974; Yin and Albrecht, 2000; Rémillard et al., 2012). Between the surface and 3 km,
 388 the profiles are broken into 5 layers: the layer below the transition, the transition layer,
 389 the layer between the transition and the inversion, the inversion layer, and the layer above
 390 the inversion. Each layer is averaged separately using a normalized height coordinate
 391 (from 0 to 1) and is combined with other layers using the averaged base and top heights
 392 of the layers as the height coordinate.

393 The difference between equivalent potential temperature at the inversion top ($\theta_{e(inv_top)}$)
 394 and base ($\theta_{e(inv_base)}$),

$$395 \quad \Delta\theta_e = \theta_{e(inv_top)} - \theta_{e(inv_base)}, \quad (2)$$

396 provides information on the stability of the MBL: lower values of $\Delta\theta_e$ indicates greater
 397 cloud-top entrainment instability because the entrained air, after becoming saturated,
 398 would be more negatively buoyant and would continue to sink further (Lilly, 1968;
 399 Deardorff, 1980). Additionally, numerical calculations demonstrate that the entrainment
 400 rate increases abruptly when $\Delta\theta_e$ decreases below a critical value (Deardorff, 1980).
 401 Approximately 10% of the profiles contained an artificial peak in water vapor mixing
 402 ratio immediately above Sc due to the wet-bulb effect of the radiosondes; when this
 403 occurred, $\theta_{e(inv_top)}$ was replaced by the value of θ_e immediately below this peak.

404 The inversion strength is quantified using the lower-tropospheric stability (LTS; Klein
 405 and Hartmann, 1993) and by the Estimated Inversion Strength (EIS; Wood and
 406 Bretherton, 2006). LTS is defined as the difference between the potential temperature at
 407 700 hPa (θ_{700}) and that at the surface ($\theta_{surface}$),

$$408 \quad LTS = \theta_{700} - \theta_{surface} . \quad (3)$$

409 EIS is defined as

$$410 \quad EIS = LTS - \Gamma_m^{850} (Z_{700} - LCL) , \quad (4)$$

411 where Γ_m^{850} is the moist adiabat at 850 hPa, Z_{700} is the 700 hPa height, and LCL is the
 412 lifting condensation level. The LCL can be approximated by Epsy's formula (Bohren and
 413 Albrecht, 1998) in terms of the surface values of the air temperature (T_s), dew point
 414 temperature (T_d), dry adiabatic lapse rate (Γ_d), and pseudo-adiabatic lapse rate (Γ_s) as

$$415 \quad LCL = \frac{T_s - T_d}{\Gamma_d - \Gamma_s} ; \quad (5)$$

416 numerically, the value of LCL (in m) is approximately 125 times the difference $T_s - T_d$ (in
 417 K). EIS, and previously LTS, have been shown to correlate well with the occurrence of
 418 the Sc-to-Cu transition (Wyant et al., 1997); however, EIS is a more physically based
 419 quantity as it accounts for the influence of the accumulated static stability between the
 420 inversion and the 700-hPa level (Wood and Bretherton, 2006).

421 To derive the statistics, all the sounding data are averaged over 36-km and converted to
 422 corresponding latitude along the great-circle route. As the statistics of the inversion layer
 423 presented below are based primarily on measurements from radiosondes, which were

launched four (eight) times every day (see Section 2.2), each sounding profile is assumed to be representative of the atmospheric structure over the 6 h (3 h) time around the sounding. Due to the sparseness of these data, these statistics might not be as robust as those derived from the radar, but nonetheless they are expected to accurately represent the observed trends.

3.5 Systematic MBL decoupling and Sc breakup detection

The commencement of the systematic MBL decoupling (DE) and Sc breakup (CB) towards Hawaii has been determined for each leg in order to better understand their triggering factors. At each location, the difference in the mean water vapor specific humidity (q_v) between the bottom 25% of the MBL and the top 25% of the MBL is calculated:

$$\Delta q = q_v(bot) - q_v(top). \quad (6)$$

This quantity is related to Δz_b , the difference in the Sc cloud bases formed below the inversion (z_b) and the LCL:

$$\Delta z_b = z_b - LCL, \quad (7)$$

where z_b is calculated as the maximum MBL cloud bases (averaged over 36 km) within four degrees longitude surrounding each radiosonde. The linear relationship between Δq and Δz_b (Figure 5a) with the slope (276 m kg g^{-1}) and intercept (200 m) comparable to those found in Jones et al. (2011), demonstrate that Δq is a robust proxy for Δz_b . Some

scatter is introduced since Δq comes from a single profile while Δz_b is an averaged maximum value. Biases might be introduced when no Sc was detected near a radiosonde or when MBLH was not well represented due to the very shallow MBL near the coast of California. Thus only those radiosondes within 1.5 standard deviations of the least square fit and that to the west of 123°W were used (336 in total). A threshold of $\Delta q > 1.5 \text{ g kg}^{-1}$ (or equivalently $\Delta z_i > 600 \text{ m}$) is found appropriate to capture the systematic decoupled MBL (Fig. 5b). Compared to the threshold of $\Delta q > 0.5 \text{ g/kg}$ ($\Delta z_i > 150 \text{ m}$) for all kinds of decoupling in VOCAL-REx (Jones et al., 2011), the systematic decoupling showed much stronger moisture stratification below the inversion. Subsequently, the DE during each transect is then defined as the most easterly profile of a group of profiles with continuous decoupling features ($\Delta q > 1.5 \text{ g kg}^{-1}$). Between the detected and the nearest east radiosonde launches, the Δq criteria for decoupling is replaced by the difference of the instantaneous ceilometer-derived cloud bases height and LCL calculated by the ship-measured T and RH. Compared to the systematic decoupling, the weak decoupled MBL is also studied in this paper and is defined as the MBL with $\Delta z_i > 150 \text{ m}$ (Consistent with Jones et al., 2011).

Due to mesoscale influences, CF_{36} sometimes shows variability and does not represent well the major cloud evolution along the transect. To reduce the effects of mesoscale variability and to more objectively capture CB, the frequency of occurrence of the MBL clouds was averaged over 108 km (CF_{108}). CB is then defined as the location along the transect from California to Hawaii where CF_{108} decreased from being greater than 80% for at least three continuous points (324 km) east of 130°W to being less than 15%. Values of CFMBL were generally not sensitive to the above criteria. No cloud breakup

points are determined if values of CFMBL east of 130°W were not sufficiently high (Leg06A, Leg06B, Leg08A, Leg08B, and Leg17B) or if they did not become sufficiently low (Leg15B). Those legs are associated with either mid-latitude or tropical cyclones, or very strong cloud outbreaks thus not represent the Sc breakup in a general sense, and their exclusion helps to elucidate the more general aspects of the transition.

4. RESULTS

The results presented below are separated into two sections. The first section includes general statistical description of MBL clouds, precipitation, thermodynamics and their seasonal behavior. The second section focuses on the study of MBL systematic decoupling and cloud breakup.

4.1 General statistics of MBL clouds, precipitation and thermodynamics

4.1.1 Cloud and Precipitation Occurrence

The fraction of time that the four main cloud types defined in Section 3.2 are detected (i.e., their frequencies of occurrence over individual legs) is shown in Figure 6a. MBL clouds are by far the most frequently observed cloud type, and all other types contribute less than 10% to the total observed hydrometeor occurrence in the column. The frequency of occurrence of MBL clouds has a broad maximum near 75% between 125°W and

491 135°W and decreases steadily to values near 20% near Hawaii. Lower values of MBL
492 cloud occurrence are also observed near the coast of California. The frequency of
493 occurrence of precipitation from MBL clouds (Fig. 6b) exhibits a very similar pattern to
494 that of MBL cloud occurrence.

495 A seasonal breakdown of MBL cloud occurrence is shown in Fig. 6c. The frequencies of
496 occurrence of MBL clouds in the warm season and the cold season exhibit generally
497 similar spatial patterns. During the warm season a broad maximum near of 90% between
498 ~122°W and ~132°W is observed, whereas that during the cold season a maximum near
499 of 70% between ~128°W and ~132°W is observed. On average, the observed MBL cloud
500 occurrence in the warm season is 20-40% higher than that observed in the cold season.

501 As expected, the frequency of occurrence of precipitation (Fig. 6d) is also generally
502 higher during the warm season than during the cold season. During the warm season
503 precipitation exhibits a maximum of ~80% between ~123°W and ~131°W, but during the
504 cold season the broad maximum, at ~55% is observed between ~128°W and ~136°W. In
505 contrast to the relatively high frequency of occurrence of clouds during the warm season
506 east of 124°W, the corresponding frequency of occurrence of precipitation decreases
507 rapidly. This might be attributed to the presence of thin clouds at this region (average
508 thickness of 180m east of 124°W compared to 300m west of 124°W).

509 The mean cloud-base height (H_b , Fig. 6e) and mean cloud-top height (H_t , Fig. 5f) of the
510 lowest cloud layer show little seasonal variability except for the regions east of 125°W.
511 Mean values of H_b increase gradually from 0.6 km near the coast of California to 1 km
512 near 135°W and remain at around 1 km further west but exhibit increasing fluctuations,

reflecting the intermittent presence of small-scale Cu clouds below the Sc. H_t values are on average 230 m greater than those of H_b and also exhibit fluctuations west of 135°W. East of 125°W we found the most noticeable difference in H_b and H_t between the warm and cloud season. The lower H_b east of 125°W during the warm season might be attributed to the stronger coastal upwelling that results to lower SST (Fig. 6g), while the higher H_t east of 125°W during the cold season might be attributed to a frontal system that occurred during Leg06B and a low-pressure system during Leg07B.

Most of the precipitation produced by MBL clouds is in the form of virga (Fig. 6h). Virga is the dominant precipitation type over the entire transect except for the region east of 126°W. The virga frequency of occurrence peaks at 40% near 130°W, whereas that of drizzle (precipitation that reaches the surface) exhibits a sharp maximum of more than 30% near 124°W, contributing to the noticeable peak of precipitation frequency at this location (Fig. 6b and Fig. 6d). The increasing frequency of occurrence virga and decreasing frequency of occurrence of drizzle from 124°W to 130°W (Fig. 6h) might be attributed to the increasing cloud-base height (Fig. 6e) caused by the warmer SST (Fig. 6g) away from the California coast. The low frequencies of occurrence (less than 10%) of both virga and drizzle near Hawaii are consistent with the low frequency of occurrence of MBL clouds there (Fig. 6a, 6b, 6c). At the same time, the low frequencies of occurrence of both virga and drizzle east of 124°W is associated with the presence of thinner clouds in that region (Fig. 6e, 6f) and to the higher in-cloud cloud droplet concentrations. The mean surface CCN at 119°N is on averaged 150/cm³ higher than that around 122°W (Lohmann and Feichter, 2005).

4.1.2 Spatial and seasonal behavior of MBLH and EIS

The mean and seasonal values of MBLH and EIS (Eq. 4) along the MAGIC transect are shown in Fig. 7. The mean MBLH (Fig. 7a) generally increases from California to Hawaii, with slightly lower values in the warm season. The largest differences in MBLHs between the warm season and cold season are observed east of 125°W, with those observed during the cold season being nearly twice as high as those during the warm season near the coast of California. The low MBLH east of 125°W during the warm season results in thin clouds (see section 4.1.1) and correspondingly a low frequency of precipitation there (Fig. 6d). The deeper MBL during the cold season is consistent with the high H_t (Fig. 6f), which might be due to synoptic influences (see Section 4.1.1). The decrease in MBLH between 145°W and 150°W can also attributed to synoptic influences and is discussed below. The trend in MBLH along the MAGIC transect (Fig. 7a) follows that of H_t (Fig. 6f) east of 135°W, indicating the capped feature of the MBL, while values of H_t west of 135°W are generally less than those of MBLH, indicating MBL decoupling and Cu-under-Sc cloud regimes.

The mean EIS (Fig. 7b) decreases gradually from 9 K near the coast of California to around 2 K near Hawaii, with values in the warm season being 1-3 K higher and those during the cold season 1-3 K lower, the differences decreasing toward Hawaii. EIS shows strong linear relationship with SST along the transect (figure not shown), while in terms of the seasonal variability, leg-mean EIS is mainly determined by the leg-mean potential temperature at 700 hPa (Fig. 7c). This quantity exhibits a larger seasonal variability, ranging from 7 K to 11 K among different legs due to the subsidence of the dry warm air

from above the inversion layer, while θ_{surface} , which depends largely on SST, varies less than 1 K between seasons.

It is likely that the seasonal variability of EIS (Fig. 7b) contributes to the seasonal variability in the frequency of occurrence of MBL clouds (higher amount of MBL cloud in the warm season and lower in the cold season), consistent with the conclusion of Wood and Bretherton (2006) that stratus cloud fraction is largely determined by EIS. The higher frequency of occurrence of MBL clouds during the warm season when EIS was higher (Fig. 6c) reflects the importance of the strong warm-season large-scale Hadley cell (Xu and Cheng, 2013) that brings dry warm air downward, leading to higher values of θ_{700} (Fig. 7c).

4.1.3 Sc and Cu occurrence and thermodynamic features

Frequencies of occurrence of the two important MBL cloud types, Sc and Cu, are examined in this section. Statistics of total and seasonal occurrence of Sc and Cu are shown in Fig. 8. The frequency of occurrence of Sc attains a broad maximum near 60% between 125°W and 135°W, and decreases to near 0% near Hawaii. The decrease in frequency of occurrence of Sc is not uniform along the MAGIC transect, and is greatest near 137°W and near 144°W, consistent with the sharp decreases in MBL cloud occurrence (Fig. 6a) at these locations. In contrast, the frequency of occurrence of Cu is always low, but steadily increases from near 5% near the coast of California to over 10% near Hawaii. Sc are more frequently observed during the warm season than during the cold season, while the occurrence of Cu is almost the same for both seasons, with slightly

more frequent cold-season Cu close to the coast of California and slightly more warm-season Cu close to Hawaii.

The comparison between ceilometer-detected Sc base height and MBLH from corresponding radiosondes indicate that 80% of the Sc clouds formed directly below the MBL inversion. Accordingly ceilometer-detected Cu bases heights show broader distribution but mainly occur near the top of the transition layer detected in corresponding radiosondes (figures not shown). Fig. 9 shows the averaged thermodynamic structure for Cu (including multi-layer MBL cloud) and single-layer Sc. A total of 141 radiosonds were analyzed: 104 with Sc near the inversion (Sc top no more than 200 m below the MBLH) and 37 with Cu near the transition or multilayer cases (Cu base more than 200 m above the transition-layer tops). Application of a layer-by-layer averaging method for the soundings for each cloud category (see section 3.4 for details) requires detectable inversion and transition layers. Analyses of the MAGIC sounding data indicate that both layers are present in the vast majority (94%) of the soundings. Note that the presence of a transition layer does not necessarily indicate a systematic decoupled MBL (see section 3.5).

As seen in Fig. 9, the large vertical gradients in potential temperature and water vapor mixing ratio near 1.5 km indicate the heights and strengths of the inversion layers, while the smaller changes below 1 km correspond to transition layers (Fig. 9). The standard deviations of both potential temperature and mixing ratio for both categories are relatively small below the inversion layer, indicating little seasonal variability in profiles of these quantities. Sc cases exhibit lower inversion- and transition-layer heights than Cu cases, and they have greater potential temperature differences across the inversion

(around 10 K compared to near 5 K for Cu); the mixing ratio differences across the inversion are nearly the same for both cases, around 6 gkg⁻¹. Sc cases exhibit smaller jumps across the transition layer than Cu for both potential temperature (< 0.5 K compared to near 1 K for Cu) and mixing ratio (< 1 gkg⁻¹ compared to 2 gkg⁻¹ for Cu); thus Cu cases are associated with a much stronger transition layer than Sc, implying a greater chance of a decoupled MBL. However, this stronger transition results in part because the cumuli help maintain the transition layer by mixing dry and warm air from the free troposphere downward.

4.2 MBL systematic decoupling and cloud breakup

The locations of DE and CB for each leg are shown in Fig. 10. Consistent with previous studies (e.g. Albrecht et al., 1995a; Bretherton and Pincus, 1995; Wyant et al. 1997; Sandu and Stevens, 2011), DE occurred east of CB (when the latter was determined) on all legs. In this section, legs with both DE and CB detected are further examined with the intention to discuss some of the potential controlling factors that are usually neglected in the numerical simulations.

4.2.1 Possible controlling factors of MBL systematic decoupling

Since DEs and CBs occurred at different locations for individual transects, a normalization of each leg is required in order to develop composites of variables across

these points. Each leg is divided into three regions: east of DE, between DE and CB, and west of CB, and the distance along the transect in each region is normalized (that is, for each leg, the distance from California to any location east of DE is divided by the distance between California and CB, and the CFMBL values for all legs used are averaged over this normalized distance, and similarly for the region between DE and CB, CB and Hawaii).

The radiosonde-derived LCL, shown in Fig. 11a as a function of this normalized distance, increases from 450m to 750m east of DE, decreases slightly to the west of DE, and remains at around 750m to the west of CB. Numerical simulations accurately capture the increasing trend of LCL with increasing SST over the well-mixed MBL (Wyant et al. 1997; Sandu and Stevens, 2011). The LCL height is more sensitive to the surface moisture than to the temperature. Thus, the increasing LCL height implies a gradual drying of the MBL, mainly due to the increasing entrainment rate with higher SST that is necessary to maintain the energy balance (Bretherton and Wyant, 1997). During MAGIC, it is found that the LCL increase rapidly near DE (Fig. 11a), and the maximum increase in LCL with SST near DEs range from 122m/K to 369m/K. The MAGIC observations suggest that this sudden dryness of the MBL is correlated with the entrainment of dryness above the inversion.

Fig. 11b shows the mean $\Delta\theta_e$ (i.e., averaged over legs) over normalized distance from California to DE. $\Delta\theta_e$ initially increased to near -2 K and then decreased to -6 K at DE. This decrease is mainly due to the large mean mixing ratio difference across the inversion (Fig. 11c). Plausible explanations for the drier conditions above the inversion are small displacements of the Hadley cell or cold outbreaks behind trailing cold fronts of mid-

648 latitude cyclones indicated by the increasing sea surface pressure from California to DE
649 (Fig. 10b). The advection due to the large-scale circulation might also contribute.
650 Meanwhile, the increase in both the potential temperature above the inversion (caused by
651 the subsidence of the dry warm air) and that below (due to the increasing SST; Fig. 6g)
652 explains the maintenance of the mean potential temperature difference across the
653 inversion of near 6 K (Fig. 11d), which contributed less to the decrease $\Delta\theta_e$. The drop in
654 $\Delta\theta_e$ east of the DE point increase (Fig. 12) the cloud top entrainment instability and the
655 entrainment rate (Deardorff, 1980) and subsequently less moisture in the MBL.
656 Consistent with the ‘deepening-warming mechanism’ (Bretherton and Wyant, 1997), our
657 analysis concurs that entrainment plays a crucial role in inducing the MBL decoupling
658 and the MAGIC observations also suggest that the dry warm air above the inversion
659 might be an important trigger.

660 One would expect a dramatic increase of the LHF east of DE due to the dryness of the
661 MBL; however, the mean surface latent heat maintained around 120 W/m^2 during that
662 period (Fig. 12a). Therefore, the role of increasing latent heat fluxes in generating the
663 systematic decoupling as suggested in the idealized model (Bretherton and Wyant, 1997)
664 is not captured by the MAGIC observations. The reduction of the mean surface wind
665 speed from California to DE (Fig. 12b) regulate the increase of the LHF, when the ship
666 moved from the edge to the center of the high-pressure system (Fig. 10b). Thus, we
667 conclude that LHF might be important in maintaining the systematic decoupling, but LHF
668 does not play the dominant role in inducing decoupling in MAGIC, which is consistent
669 with the conclusion of Jones et al (2011).

Apart from the systematic decoupling, the weak decoupling ($\Delta z_i > 150\text{m}$) east of DE are also investigated. Broad peaks of the frequency of occurrence of virga and drizzle were found east of DE, while these frequencies decreased to below 20% when MBL was decoupled (figures not shown). We found 87% of which are associated with precipitation. We conclude that precipitation might play a role in inducing decoupling, but this decoupling is usually weak and not continuous, thus precipitation did not show a dominant impact on systematic decoupling. Meanwhile, the diurnal decoupling might also partly explain the weak decoupling east of DE since 73% of which occurred in the local daytime from 6:00 am to 6:00pm.

4.2.3 Possible controlling factors of MBL cloud breakup during MAGIC

Values of CF_{36} are typically high along the eastern part of the MAGIC transect (Fig. 10a), especially during the warm season (Fig. 6). The locations where CF_{108} drop to below 50% are usually close to CB (CF_{108} drops to below 15%), indicating the abrupt MBL cloud breakup (Fig. 10a). At the same time, CB is typically located on the west edge of high-pressure systems (Fig. 10b), implying the role of synoptic interference in the observed MBL cloud breakup.

To further investigate this, we analyze the MBL wind profiles near the CB points. A clear synoptic-induced divergence pattern is found. Fig. 13a shows median profiles of zonal (ΔU) and latitudinal (ΔV) wind difference of the composed U(V) wind east and west of CB in the small and large region. This small region is bounded by two days of radiosonds (~1600 km) surrounding CB while the large region refers to the whole transect. The

profiles shown in Fig. 13a show little sensitivity to the details of the region selected to estimate the wind differences. The well-separated zonal and meridional wind differences around the CB indicate a systematic divergence in the MBL since the wind to the west of CB is more easterly ($\Delta U < 0 \text{ m/s}$), and more southerly ($\Delta V > 0 \text{ m/s}$) than that to the east. We interpret the east of CB wind pattern as the stable high-pressure systems and the west of CB wind pattern as the signature of approaching mid-latitude cyclone systems. Strong uplifting convergence in the east-approaching cyclones was compensated by the divergence nearby. The drop of the MBLH near CBs provides evidence of the compensating subsidence (Fig. 13b), which is absent in the idealized model with uniformed large scale forcing. Moreover, the averaged mixing ratio difference above the inversion was nearly doubled in the small region (figures not shown). We conclude that the switch of the synoptic environment to the unstable cyclone system can fast break up the MBL cloud and drive vigorous Cu or deep convective clouds.

5. SUMMARY AND DISCUSSION

The MAGIC field campaign, with nearly 200 days of ship-based observations during 20 round trips along the 4000 km transect between California and Hawaii, provided an unparalleled opportunity to acquire data on properties of MBL clouds, precipitation and thermodynamic structure. The measurements obtained during that campaign are used in this manuscript to examine the location and potential controlling factors of systematic MBL decoupling and Sc breakup.

MBL clouds were by far the most frequently observed cloud type during the MAGIC campaign. MBL clouds occurred more often during the warm season (Fig. 5), reflecting the importance of the strong warm-season large-scale Hadley cell (Xu and Cheng, 2013). Among the different MBL cloud types, Sc was the dominant MBL cloud type and occurred more frequently during the warm season than during the cold season (Fig. 8b), while the occurrence of Cu was less strongly affected by subsidence and exhibited nearly the same behavior for both seasons (Fig. 8b).

The formation of Sc just below the inversion requires a shallow MBL with a strong inversion and a weak transition (Fig. 9), providing a greater opportunity to have well-mixed MBL conditions. In contrast, Cu and multi-layer clouds are usually associated with a much stronger transition, implying a greater chance of decoupling in the MBL.

There was a high frequency of occurrence of precipitation throughout the campaign. However, the precipitation from the MBL clouds is weak and often evaporated well before reaching the ocean surface (Fig. 6e). EIS experienced a seasonal variation caused by that of θ_{700} (Fig. 7c), and generally decreased due to the increasing SST. MBLH generally increased along the transect from California to Hawaii (Fig. 7). East of 135°W the spatial behavior of MBLH parallels that of the first cloud-top height (Figs. 5f and 6d), indicating the capped feature of the MBL.

The locations of MBL systematic decoupling are determined for individual legs. It is found that a threshold of $\Delta q > 1.5$ g/kg separates the well-mixed profiles from the systematic decoupled ones (Fig. 5b). Compared to the threshold of $\Delta q > 0.5 \text{ g kg}^{-1}$ found in VOCAL-REx (Jones et al., 2011), the MAGIC systematic decoupling showed much

stronger moisture stratification below the inversion. Precipitation and solar radiation correlate well with the weak decoupling points between California and DEs, but neither of them plays a dominant role in the systematic decoupling.

A rapid increase of LCL height was found near DEs (Fig. 11s), indicating more rapid drying of the MBL. Correspondingly, the mean cloud top instability showed a sudden increase mainly due to the large mixing ratio difference across the inversion (Fig. 11b,c). These observations imply that the dry warm air from above the inversion is likely of great importance to trigger the systematic decoupling. Consistent with the results in VOCAL-REx (Jones et al., 2011), LHF does not play the dominant role in inducing systematic decoupling in MAGIC. Meanwhile, the mixed layer cloud thickness during MAGIC did not correlate well with the systematic decoupling due to the sudden change of LCL near DEs, further implying that the strong entrainment was driven more by the cloud top instability than by the in-cloud turbulence.

DEs nearly always occurred east of CBs (if present) (Fig. 10). MBL clouds tend to breakup abruptly at a location that is typically on the west edge of high-pressure systems (Fig. 10b). A change in synoptic pattern (i.e. different air mass) was often found near CB, which is associated with systematic divergence in the MBL. The divergence, together with downdrafts, compensated for the convergent uplifting in the approaching cyclones. We conclude that the cloud evolution in the idealized model seldom occurs in reality due to synoptic interference.

6. ACKNOWLEDGMENTS

The MAGIC deployment was supported and undertaken by the U.S. Department of Energy (DOE) Atmospheric Radiation Measurement (ARM) Program Climate Research Facility. The current research was supported by the DOE Atmospheric System Research (ASR) Program (Office of Science, OBER). ERL was supported by the ASR Program under Contract No. DE-AC02-98CH10886. Data used for the analyses were downloaded from the ARM archive (www.arm.gov) except for the MARMET and MARFLUX data, which were provided by Dr. Michael Reynolds of RMR Co. We thank David Painemal of NASA Langley Research Center for providing surface cloud condensation nuclei (CCN) data. We acknowledge the team of scientists and technicians who made this work possible by collecting the data and maintaining the instruments, and especially Horizon Lines and the Captain and crew of the Horizon *Spirit* for their hospitality. Special thanks go to the cloud research group at McGill University (www.clouds.mcgill.ca) for their helpful comments and constructive criticism.

781 **Appendix:**

782 Table A1 about here.

783

784 B. MAGIC Instrument Status

785 **Instrument Status Table Leg03A to Leg09B**

786 The first leg with MAGIC instrumentation was Leg01B during which the ISAR
787 was installed. During Leg02A and Leg02B, for which instrument status designations are
788 not listed, the radars and other instruments were being set up; some collected data during
789 these legs. During Leg03 most of the instruments were up and collecting data. On
790 Leg09B, the instruments were without power for extended times and were being shut
791 down. The instruments were removed from the ship after Leg09B from January, 2013
792 until May, 2013.

793 Fig. 14 about here.

794 **Instrument Status Table Leg10A to Leg18B**

795

796 MAGIC instruments were redeployed during Leg10A, and the campaign
797 continued until the end of Leg20B. The technicians did not report instrument status
798 designations for Leg19A and Leg19B, but these were probably similar to those for
799 Leg18B. During Leg20A and Leg20B the instruments were being turned off, so few data
800 were collected during these legs (although sonde launches occurred on Leg20A,

801 meteorological data were collected until the ship returned to port, and both radars were
802 operating for most of the two legs).

803 Fig 15 about here.

804

805 Table A2 about here.

806

REFERENCES

Albrecht, B. A., 1984. "A model study of downstream variations of the thermodynamic structure of the trade winds." *Tellus*, **36A**, 187-202.

Albrecht, B. A., Randall, D. A., & Nicholls, S. (1988). "Observations of marine stratocumulus clouds during FIRE." *Bulletin of the American Meteorological Society*, **69**(6), 618-626.

Albrecht, B. A., C. S. Bretherton, D. Johnson, W. H. Scubert and A. S. Frisch (1995a). "The Atlantic stratocumulus transition experiment-ASTEX." *Bulletin of the American Meteorological Society* **76**(6): 889-904.

Albrecht, B. A., Jensen, M. P., & Syrett, W. J. (1995b). "Marine boundary layer structure and fractional cloudiness." *Journal of Geophysical Research: Atmospheres* (1984–2012), **100**(D7), 14209-14222.

Augstein, E., H. Schmidt and F. Ostapoff (1974). "The vertical structure of the atmospheric planetary boundary layer in undisturbed trade winds over the Atlantic Ocean." *Boundary-Layer Meteorology* **6**(1-2): 129-150.

Bohren, C. and B. Albrecht (1998). *Atmospheric Thermodynamics*, Oxford University Press, New York, USA.

830 Boutle, I., R. Beare, S. E. Belcher, A. Brown and R. S. Plant (2010). "The moist
831 boundary layer under a mid-latitude weather system." *Boundary-layer meteorology*
832

833 Bretherton, C. S., & Pincus, R. (1995). "Cloudiness and marine boundary layer dynamics
834 in the ASTEX Lagrangian experiments. Part I: Synoptic setting and vertical structure. "
835 *Journal of the atmospheric sciences*, 52(16), 2707-2723.
836

837 Bretherton, C. S. and M. C. Wyant (1997). "Moisture transport, lower-tropospheric
838 stability, and decoupling of cloud-topped boundary layers." *Journal of the Atmospheric*
839 *Sciences* **54**(1): 148-167.
840

841 Cadeddu, M. (2012). *Microwave Radiometer–3 Channel (MWR3C) Handbook*, DOE
842 *Office of Science Atmospheric Radiation Measurement (ARM) Program (United States)*.
843

844 Cheng, A. and K.-M. Xu (2013). "Evaluating Low Cloud Simulation from an Upgraded
845 Multiscale Modeling Framework Model. Part III: Tropical and Subtropical Cloud
846 Transitions over the Northern Pacific." *Journal of Climate* **26**: 5761-5781.
847

848 Deardorff, J. (1980). "Cloud top entrainment instability." *Journal of the Atmospheric*
849 *Sciences* 37(1): 131-147.
850

851 Fairall, C. W., E. F. Bradley, D. P. Rogers, J. B. Edson, and G. S. Young (1996). "Bulk
852 parameterization of air-sea fluxes for Tropical Ocean-Global Atmosphere Couple-Ocean

853 Atmosphere Response Experiment." *Journal of Geophysical Research* 101 (C2):3747-
854 3764.

855

856 Heck, P. W., B. J. Byars, D. F. Young, P. Minnis, and E. F. Harrison 1990. "A
857 climatology of satellite derived cloud properties over marine stratocumulus regions. "
858 Preprints, Conf. on Cloud Physics, San Francisco, CA, Amer. Meteor. Soc., J1–J7.

859

860 Hildebrand, P. H. and R. Sekhon (1974). "Objective determination of the noise level in
861 Doppler spectra." *Journal of Applied Meteorology* 13(7): 808-811.

862

863 Jones, C. R., Bretherton, C. S., & Leon, D. (2011). "Coupled vs. decoupled boundary
864 layers in VOCALS-REx." *Atmospheric Chemistry and Physics*, 11(14), 7143-7153.

865

866 Karlsson, J., Svensson, G., Cardoso, S., Teixeira, J., & Paradise, S. (2010). "Subtropical
867 cloud-regime transitions: Boundary layer depth and cloud-top height evolution in models
868 and observations." *Journal of Applied Meteorology and Climatology*, 49(9), 1845-1858.

869

870 King, M. D., Y. J. Kaufman, W. Menzel and D. Tanre (1992). "Remote sensing of cloud,
871 aerosol, and water vapor properties from the Moderate Resolution Imaging Spectrometer
872 (MODIS)." *IEEE Transactions on Geoscience and Remote Sensing* 30(1): 2-27.

873

874 Klein, S. A. and D. L. Hartmann (1993). "The seasonal cycle of low stratiform clouds."
875 *Journal of Climate* 6(8): 1587-1606.

876

877 Kollias, P., E. Clothiaux, M. Miller, B. Albrecht, G. Stephens and T. Ackerman (2007).
878 "Millimeter-wavelength radars: New frontier in atmospheric cloud and precipitation
879 research." *Bulletin of the American Meteorological Society* **88**(10): 1608-1624.

880

881 Krueger, S. K., McLean, G. T., & Fu, Q. (1995). "Numerical simulation of the stratus-to-
882 cumulus transition in the subtropical marine boundary layer. Part I: Boundary-layer
883 structure. *Journal of the atmospheric sciences*. " **52**(16), 2839-2850.

884

885 Larson, K., D. L. Hartmann and S. A. Klein (1999). "The role of clouds, water vapor,
886 circulation, and boundary layer structure in the sensitivity of the tropical climate." *Journal*
887 *of Climate* **12**(8): 2359-2374.

888

889 Lewis, E. R., W. J. Wiscombe, B. A. Albrecht, G. L. Bland, C. N. Flagg, S. A. Klein, P.
890 Kollias, G. Mace, R. Reynolds and S. Schwartz (2012). "MAGIC: Marine ARM GPCI
891 Investigation of Clouds."

892

893 Liljegren, J. (1994). Two-channel microwave radiometer for observations of total column
894 precipitable water vapor and cloud liquid water path. *Fifth Symposium on Global Change*
895 *Studies*.

896

897 Lilly, D. K. (1968). "Models of cloud-topped mixed layers under a strong inversion."
898 *Quarterly Journal of the Royal Meteorological Society* **94**(401): 292-309.

899

900 Lohmann, U. and J. Feichter (2005). "Global indirect aerosol effects: a review."
901 Atmospheric Chemistry and Physics **5**(3): 715-737.

902

903 Ma, C.-C., C. R. Mechoso, A. W. Robertson and A. Arakawa (1996). "Peruvian stratus
904 clouds and the tropical Pacific circulation: A coupled ocean-atmosphere GCM study."
905 Journal of Climate **9**(7): 1635-1645.

906

907 Matrosov, S. Y. (2007). "Potential for attenuation-based estimations of rainfall rate from
908 CloudSat." Geophysical Research Letters **34**(5): L05817.

909

910 Miller, M. A., M. P. Jensen and E. E. Clothiaux (1998). "Diurnal cloud and
911 thermodynamic variations in the stratocumulus transition regime: A case study using in
912 situ and remote sensors." Journal of the Atmospheric Sciences **55**(13): 2294-2310.

913

914 Morris, V. (2006). "Microwave radiometer (MWR) handbook." ARM-TR016.

915

916 Neiburger, M. (1961). Studies of the structure of the atmosphere over the eastern Pacific
917 ocean in summer, University of California Press.

918

919 Norris, J. R. and C. B. Leovy (1994). "Interannual variability in stratiform cloudiness and
920 sea surface temperature." Journal of Climate **7**(12): 1915-1925.

921

922 Philander, S., D. Gu, G. Lambert, T. Li, D. Halpern, N. Lau and R. Pacanowski (1996).
 923 "Why the ITCZ is mostly north of the equator." *Journal of Climate* **9**(12): 2958-2972.
 924

925 Rémillard, J., P. Kollias, E. Luke and R. Wood (2012). "Marine Boundary Layer Cloud
 926 Observations in the Azores." *Journal of Climate* **25**(21): 7381-7398.
 927

928 Riehl, H., T. Yeh, J. Malkus and N. La Seur (1951). "The north-east trade of the Pacific
 929 Ocean." *Quarterly Journal of the Royal Meteorological Society* **77**(334): 598-626.
 930

931 Rossow, W. B. and R. A. Schiffer (1999). "Advances in understanding clouds from
 932 ISCCP." *Bulletin of the American Meteorological Society* **80**(11): 2261-2287.
 933

934 Sandu, I. and B. Stevens (2011). "On the factors modulating the stratocumulus to
 935 cumulus transitions." *Journal of the Atmospheric Sciences* **68**(9): 1865-1881.
 936

937 Stephens, G. L. (2005). "Cloud feedbacks in the climate system: A critical review."
 938 *Journal of Climate* **18**(2): 237-273.
 939

940 Teixeira, J., S. Cardoso, M. Bonazzola, J. Cole, A. DelGenio, C. DeMott, C. Franklin, C.
 941 Hannay, C. Jakob, Y. Hiao, J. Karlsson, H. Kitagawa, M. Köhler, A. Kuwano-Hoshida, C.
 942 LeDrain, A. Lock, M. H. Miller, P. Marquet, J. Martins, C. R. Mechoso, E. V. Meijgaard,
 943 I. Meinke, A. M. A. Miranda, D. Mironov, R. Neggers, H. L. Pan, D. A. Randall, P. J.
 944 Rasch, B. Rockel, W. B. Rossow, B. Ritter, A. P. Siebesma, P. Soares, F. J. Turk, P. A.

945 Vaillancourt, A. Von Engeln, and M. Zhao (2011). "Tropical and subtropical cloud
 946 transitions in weather and climate prediction models: The GCSS/WGNE Pacific Cross-
 947 Section Intercomparison (GPCI)." *Journal of Climate* **24**(20): 5223-5256.

948

949 Tiedtke, M., W. A. Hackley and J. Slingo, 1988. "Tropical forecasting at ecmwf: The
 950 influence of physical parameterization on the mean structure of forecasts and analyses."
 951 *Q. J. R. Meteorol. Soc.*, 114, 639–664.

952

953 Wang, S., B. A. Albrecht, and P. Minnis, 1993. "A regional simulation of marine
 954 boundary layer clouds." *J. Atmos. Sci.*, **50**, 4022–4043.

955

956 Wyant, M. C., C. S. Bretherton, H. A. Rand, and D. E. Stevens, 1997. "Numerical
 957 simulations and a conceptual model of the subtropical marine stratocumulus to trade
 958 cumulus transition." *J. Atmos. Sci.*, **54**, 168–192.

959

960 Widener, K., N. Bharadwaj and K. Johnson (2012). "Ka-Band ARM Zenith Radar
 961 (KAZR) Instrument Handbook." DOE Office of Science Atmospheric Radiation
 962 Measurement (ARM) Program (United States).

963

964 Wood, R. (2012). "Stratocumulus clouds." *Monthly Weather Review* **140**(8): 2373-2423.

965

Wood, R. and C. S. Bretherton (2004). "Boundary layer depth, entrainment, and decoupling in the cloud-capped subtropical and tropical marine boundary layer." *Journal of Climate* **17**(18): 3576-3588.

Wood, R. and C. S. Bretherton (2006). "On the relationship between stratiform low cloud cover and lower-tropospheric stability." *Journal of Climate* **19**(24): 6425-6432.

Wood, R., C. Mechoso, C. Bretherton, R. Weller, B. Huebert, F. Straneo, B. Albrecht, H. Coe, G. Allen and G. Vaughan (2011). "The vamos ocean-cloud-atmosphere-land study regional experiment (VOCALS-REx): Goals, platforms, and field operations." *Atmospheric Chemistry and Physics* **11**(2): 627-654.

Xu, K.-M. and A. Cheng (2013). "Evaluating Low Cloud Simulation from an Upgraded Multiscale Modeling Framework Model. Part II: Seasonal Variations over the Eastern Pacific." *Journal of Climate* **26**: 5717-5740.

Yin, B. and B. A. Albrecht (2000). "Spatial variability of atmospheric boundary layer structure over the eastern equatorial Pacific." *Journal of Climate* **13**(9): 1574-1592.

7. TABLES

Table 1. Cloud types and characteristics used to differentiate them in the identification algorithm (Ind: indeterminate).

		Type					
		High-level	Mid-level	MBL			Cu congestus and deep convective
				Sc	Cu	Ind	
Minimum cloud base		≥ 6 km	3-6 km	< 3 km*	< 3 km*	< 3 km*	< 3 km
Maximum cloud top		—	—	< 3 km	< 3 km	< 3 km	≥ 3 km
Duration		—	—	≥ 20 min	< 20 min	≥ 20 min	—
Cloud top height standard deviation	[20 min, 2 h]	—	—	< 100 m	—	≥ 100 m	—
	(2 h, 10h]	—	—	< 160 m	—	≥ 160 m	—
	> 10 h	—	—	< 200 m	—	≥ 200 m	—

* Minimum cloud base for MBL clouds are either below 3 km or undetermined

1003 Table 2. List of liquid precipitation types and their main characteristics used to
 1004 differentiate them.

1005

1006

	Type				
	Virga	Drizzle	Warm Rain	Cold Rain	Convective Rain
Echo base	> First gate	= First gate	= First gate	= First gate	= First gate
Base reflectivity	—	<0 dBZ	>0 dBZ	>0 dBZ	>0 dBZ & > First Top reflectivity
Surface Rain	No	No	Yes	Yes	Yes
First cloud top height	Below 0°C isotherm	Below 0°C isotherm	Below 0°C isotherm	Above 0°C isotherm	—
Echo below cloud base	Yes (50 m lower)	Yes (50 m lower)	Possible	Possible	Possible

1007

1008

1009

1010

1011

1012

1013

1014

1015 Table A1. Departure and arrival times of each leg.

1016 **All times are UTC.**

1017

	A			B	
	Depart LA	Arrive HI		Depart HI	Arrive LA
Leg00	2012-02-11, 13:30	2012-02-16, 09:00		2012-02-17, 09:00	2012-02-23, 15:00
Leg01				2012-09-14, 23:20	2012-09-20, 13:40
Leg02	2012-09-22, 12:15	2012-09-27, 05:50		2012-09-28, 09:50	2012-10-04, 14:20
Leg03	2012-10-06, 11:30	2012-10-11, 06:30		2012-10-12, 08:00	2012-10-18, 13:30
Leg04	2012-10-20, 11:25	2012-10-25, 06:15		2012-10-26, 06:40	2012-11-01, 13:20
Leg05	2012-11-03, 17:50	2012-11-08, 15:00		2012-11-09, 17:30	2012-11-15, 15:00
Leg06	2012-11-17, 12:20	2012-11-22, 07:30		2012-11-24, 10:15	2012-11-30, 01:20
Leg07	2012-12-01, 13:30	2012-12-06, 09:00		2012-12-07, 08:20	2012-12-13, 14:45
Leg08	2012-12-15, 13:00	2012-12-20, 08:30		2012-12-22, 08:15	2012-12-28, 00:00
Leg09	2012-12-29, 12:30	2013-01-03, 07:00		2013-01-05, 04:50	2013-01-13, 03:35 SEE NOTE 1
Leg10	2013-05-11, 11:20	2013-05-16, 06:20		2013-05-17, 16:30	2013-05-23, 14:00
Leg11	2013-05-25, 11:25	2013-05-30, 06:30		2013-05-31, 11:15	2013-06-06, 13:30
Leg12	2013-06-08, 11:25	2013-06-13, 06:35		2013-06-14, 16:35	2013-06-20, 13:35
Leg13	2013-06-22, 11:30	2013-06-27, 07:45		2013-06-28, 17:30	2013-07-03, 23:10
Leg14	2013-07-07, 17:35	2013-07-12, 06:50		2013-07-13, 11:45	2013-07-18, 23:15
Leg15	2013-07-20, 12:00	2013-07-25, 05:45		2013-07-26, 13:10	2013-08-01, 13:30
Leg16	2013-08-03, 13:30	2013-08-08, 05:40		2013-08-09, 10:15	2013-08-15, 14:30
Leg17	2013-08-17, 18:30	2013-08-22, 10:15		2013-08-23, 17:15	2013-08-29, 13:20
Leg18	2013-08-31, 11:45	2013-09-05, 06:35		2013-09-06, 12:00	2013-09-12, 13:40
Leg19	2013-09-14, 12:25	2013-09-19, 06:20		2013-09-20, 11:20	2103-09-26, 13:55
Leg20	2013-09-28, 11:35	2013-10-03, 06:35		2013-10-04, 10:05	2013-10-10, 13:35 SEE NOTE 2

1018

1019 NOTE 1: During Leg09B the *Spirit* had its engines off for approximately 14 hours on
1020 2013-01-06 and 2013-01-07; thus the trajectory will look abnormal for this time as the
1021 ship was drifting. Soon thereafter the entire ship, including the AMF2, was without
1022 power for approximately one hour, and some instruments might not have resumed
1023 operation before the end of the leg. Data acquisition stopped 2013-01-11 for some
1024 instruments and on 2013-01-12 for all instruments. After the *Spirit* arrived in port in Los

1025 Angeles after Leg09B, the AMF2 was removed from the ship (it was completely off the
1026 Spirit by 2013-01-13, 21:00 UTC) and placed in storage, where it remained until
1027 reinstallation on 2013-05-09.

1028 NOTE 2: MAGIC instrumentation was being turned off and packed during Leg20 and all
1029 MAGIC instrumentation was removed from the *Spirit* on 2013-10-10 by 22:00.

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

Table A2. Statistics (means and standard deviations) of MBL clouds macroscopic characteristics. The 1st cloud base, top and thickness in the table refer to those of the lowest clouds.

Distance from California Coast		100 km	500 km	1000 km	2000 km	3000 km	3600 km	4000 km
Cloud fractional coverage	Total	0.62±0.44	0.69±0.43	0.74±0.41	0.62±0.38	0.30±0.38	0.33±0.28	0.22±0.28
	Cold	0.49±0.47	0.54±0.46	0.71±0.44	0.54±0.44	0.21±0.27	0.29±0.26	0.17±0.27
	Warm	0.76±0.37	0.88±0.31	0.79±0.40	0.71±0.29	0.40±0.48	0.38±0.31	0.27±0.29
1 st Cloud base [km]	Total	0.51±0.42	0.62±0.34	0.98±0.38	1.14±0.28	1.05±0.40	1.20±0.43	1.11±0.46
	Cold	0.85±0.49	0.69±0.47	1.04±0.52	1.10±0.20	1.09±0.49	1.09±0.46	1.17±0.53
	Warm	0.27±0.08	0.55±0.15	0.92±0.17	1.18±0.36	1.00±0.26	1.32±0.39	1.04±0.37
1 st Cloud top [km]	Total	0.89±0.75	0.91±0.42	1.19±0.28	1.39±0.28	1.20±0.47	1.43±0.47	1.24±0.35
	Cold	1.36±0.90	1.02±0.56	1.17±0.35	1.39±0.24	1.19±0.57	1.31±0.52	1.15±0.31
	Warm	0.48±0.13	0.80±0.16	1.20±0.19	1.39±0.33	1.23±0.34	1.57±0.39	1.33±0.39
1 st Cloud thickness [km]	Total	0.35±0.39	0.27±0.22	0.23±0.15	0.23±0.16	0.19±0.15	0.17±0.13	0.24±0.18
	Cold	0.60±0.51	0.30±0.30	0.18±0.11	0.25±0.16	0.18±0.16	0.15±0.15	0.26±0.19
	Warm	0.18±0.14	0.24±0.11	0.27±0.17	0.21±0.16	0.21±0.15	0.21±0.08	0.21±0.17

Means and standard deviations of the cloud macroscopic properties at different locations along the MAGIC transect (100, 500, 1000, 2000, 3000, 3600 and 4000 km from the coast of California) for all applicable MAGIC legs (Leg3-8, Leg11, Leg14-17). Data during each leg are binned to a uniform great circle route with 36-km resolution.

Figure Caption List:

1. (a) Horizon *Spirit* showing location of the bridge region where the AMF2 was located, and (b) Tracks of MAGIC legs between California and Hawaii (red lines) and great circle route (blue line).
2. KAZR reflectivity observations for Leg04A (top panel) and Leg04B (bottom panel) with the first ceilometer cloud base shown as black dots. The *Spirit* departed from California on Oct. 21 (2012) and arrived in Hawaii on Oct. 25, then left on Oct. 26 and returned to California on Nov. 1. Graphs are shown with Los Angeles to the right and Honolulu to the left. The ceilometer data for 00:00 to ~21:16 UTC on Oct. 22, 2012 are not available.
3. (a) Cloud classifications for Leg04 with the first ceilometer cloud base shown as black dots. The corresponding KAZR reflectivities can be seen in Fig 2. (b) KAZR reflectivities with the first and second ceilometer cloud base shown as black and red dots respectively for a short time segment on Oct. 21, 2012 during Leg04A. (c) MBL cloud classifications with the first and second ceilometer cloud base shown as black and red dots for the same time period as that shown in Fig. 3b during Leg04A. (d) KAZR reflectivities with the first ceilometer cloud base shown as black dots for a short time period on Oct. 28, 2012 during Leg04B. The pink, light blue and purple backgrounds indicate the occurrence of virga, MBL drizzle, and heavy MBL drizzle, respectively; the yellow background indicates no precipitation during that time period.

1075 4. Profiles of temperature (T), potential temperature (θ), and mixing ratio (r)
1076 from soundings. Green and blue dashed lines show the inversion tops and
1077 bases, while black and red dashed lines show the transition tops and bases. (a)
1078 Sounding launched at 18:08 UTC, October 26, 2012, with broken Cu
1079 overhead. (b) Sounding launched at 12:00 UTC, October 29, 2012, with
1080 overcast Sc overhead.

1081

1082

1083 5. (a) The scatter plot of radiosonde-derived Δq for all the legs and maximum
1084 Δz_b within four degrees longitude surrounding each radiosonde. The solid
1085 black line is the least square fit with slope 276 m kg g⁻¹ and intercept 200m.
1086 The dashed black line represents the thermodynamic argument derived in Jone
1087 et al 2011. The red dots indicate outliers outside 1.5 standard deviation of the
1088 least-square fit. (b) Radiosonde-derived Δq for all the legs along the
1089 normalized path from California to DE to Hawaii.

1090

1091 6. Frequencies of occurrence along the MAGIC transect of (a) the four main
1092 cloud types, (b) MBL clouds and observed MBL-cloud liquid precipitation,
1093 and (c) MBL clouds and (d) observed MBL-cloud liquid precipitation during
1094 the warm and cold season during the warm season (Leg11, Leg14-Leg17:
1095 May 25 -June 6, 2013, July 7-Aug. 29, 2013) and during the cold season
1096 (Leg03-Leg08: Oct. 6-Dec. 27, 2012). Total and seasonal mean along the
1097 MAGIC transect of (e) first cloud base heights, (f) first cloud top heights, and

1098 (g) SSST. The black line shows the total mean, the blue and the red line shows
1099 the mean for cold and warm season respectively. The gray shaded region
1100 indicates one standard deviation of the mean. Frequencies of occurrence along
1101 the MAGIC transect of (h) liquid precipitation types.

1102

1103 7. Total and seasonal mean of (a) MBLH, and (b) EIS along the MAGIC transect.
1104 The gray shaded region indicates one standard deviation of total. The black
1105 line shows the total mean, the blue line shows the mean of the cold season,
1106 and the red line the mean for the warm season, and (c) Leg-mean values of
1107 EIS, SSST, potential temperature at 700 hPa, and potential temperature at the
1108 surface.

1109

1110 8. Frequencies of occurrence of cloud types (a) over the entire deployment and
1111 (b) during the cold season and warm season.

1112

1113 9. Profiles of means and standard deviations of (a) potential temperature and (b)
1114 water vapor mixing ratio composed over the cases with single stratocumulus
1115 at the inversion level (blue dotted line), and single cumulus at the transition
1116 level together with multiple MBL clouds (red solid line), both of which must
1117 contain a transition layer.

1118

1119 10. (a) CF₃₆, (b) the sea surface pressure along the MAGIC transect for individual
1120 legs. The black triangle and the black dots indicate the cloud breakup points

1121 (CB) and the starting points of the MBL systematic decoupling (DE). The red
1122 crosses in (a) indicate the location when CF_{108} drops to below 50% before CB.
1123 White spaces in (a) denote times when CF_{36} was less than 10%, and those in
1124 (b) denote missing data.

1125

1126 11. Total mean of (a) the radiosonde-derived LCL along the normalized path from
1127 California to Hawaii, (b) the equivalent potential temperature difference
1128 across the inversion, (c) the mixing ratio difference at the inversion, and (d)
1129 the potential temperature difference at the inversion along the normalized path
1130 from California to DE. Gray shaded region indicates interquartile range. The
1131 boxplot in (b) and (c) indicates the location of the maximum and minimum
1132 value near DE, the difference between the two is significant at 95% confident
1133 level.

1134

1135 12. The total mean (a) surface latent heat flux, and (b) surface wind speed along
1136 the normalized path from California to Hawaii. Gray shaded region indicates
1137 interquartile range.

1138

1139 13. (a) Profiles of medians of U wind difference of the composed U wind east of
1140 CB and that west of CB in the small region (solid black line) and in the large
1141 region (dashed black line); same for V wind difference in the small region
1142 (solid red line) and in the large region (dashed red line). Shaded area indicates

1143 interquartile range, and (b) MBLH along normalized path from California to
1144 Hawaii.

1145 14. MAGIC Instrument Status from Leg03A to Leg09B

1146 15. MAGIC Instrument Status from Leg10A to Leg18B

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

Figures

1166

1167

1168

1169

1170

1171

Fig. 1. (a) *Horizon Spirit* showing location of the bridge region where the AMF2 was located, and (b) Tracks of MAGIC legs between California and Hawaii (red lines) and great circle route (blue line).

Fig. 2. KAZR reflectivity observations for Leg04A (top panel) and Leg04B (bottom panel) with the first ceilometer cloud base shown as black dots. The *Spirit* departed from California on Oct. 21 (2012) and arrived in Hawaii on Oct. 25, then left on Oct. 26 and returned to California on Nov. 1. Graphs are shown with Los Angeles to the right and Honolulu to the left. The ceilometer data for 00:00 to ~21:16 UTC on Oct. 22, 2012 are not available.

Fig. 3. (a) Cloud classifications for Leg04 with the first ceilometer cloud base shown as black dots. The corresponding KAZR reflectivities can be seen in Fig 2. (b) KAZR reflectivities with the first and second ceilometer cloud base shown as black and red dots respectively for a short time segment on Oct. 21, 2012 during Leg04A. (c) MBL cloud classifications with the first and second ceilometer cloud base shown as black and red dots for the same time period as that shown in Fig. 3b during Leg04A. (d) KAZR

1189 reflectivities with the first ceilometer cloud base shown as black dots for a short time
1190 period on Oct. 28, 2012 during Leg04B. The pink, light blue and purple backgrounds
1191 indicate the occurrence of virga, MBL drizzle, and heavy MBL drizzle, respectively; the
1192 yellow background indicates no precipitation during that time period.

1193

Fig. 4. Profiles of temperature (T), potential temperature (θ), and mixing ratio (r) from soundings. Green and blue dashed lines show the inversion tops and bases, while black and red dashed lines show the transition tops and bases. (a) Sounding launched at 18:08 UTC, October 26, 2012, with broken Cu overhead. (b) Sounding launched at 12:00 UTC, October 29, 2012, with overcast Sc overhead.

Fig. 5(a) The scatter plot of radiosonde-derived Δq for all the legs and maximum Δz_b within four degrees longitude surrounding each radiosonde. The solid black line is the least square fit with slope 276 m kg g^{-1} and intercept 200m . The dashed black line represents the thermodynamic argument derived in Jones et al 2011. The red dots indicate outliers outside 1.5 standard deviation of the least-square fit. (b) Radiosonde-derived Δq for all the legs along the normalized path from California to DE to Hawaii.

Fig. 6. Frequencies of occurrence along the MAGIC transect of (a) the four main cloud types, (b) MBL clouds and observed MBL-cloud liquid precipitation, and (c) MBL clouds and (d) observed MBL-cloud liquid precipitation during the warm season (Leg11, Leg14-Leg17: May 25 -June 6, 2013, July 7-Aug. 29, 2013) and during the cold season (Leg03-Leg08: Oct. 6-Dec. 27, 2012). Total and seasonal mean along the MAGIC transect of (e) first cloud base heights, (f) first cloud top heights, and (g) SSST. The black line shows the total mean, the blue and the red line shows the mean for cold and warm season respectively. The gray shaded region indicates one standard deviation of the mean. Frequencies of occurrence along the MAGIC transect of (h) liquid precipitation types.

1222

1223

1224 Fig. 7. Total and seasonal mean of (a) MBLH, and (b) EIS along the MAGIC transect.
 1225 The gray shaded region indicates one standard deviation of total. The black line shows
 1226 the total mean, the blue line shows the mean of the cold season, and the red line the mean
 1227 for the warm season, and (c) Leg-mean values of EIS, SSST, potential temperature at 700
 1228 hPa, and potential temperature at the surface.

1229

1230 Fig. 8. Frequencies of occurrence of cloud types (a) over the entire deployment and (b)

1231 during the cold season and warm season.

Fig. 9. Profiles of means and standard deviations of (a) potential temperature and (b) water vapor mixing ratio composed over the cases with single stratocumulus at the inversion level (blue dotted line), and single cumulus at the transition level together with multiple MBL clouds (red solid line), both of which must contain a transition layer.

Fig. 10(a) CF_{36} , (b) the sea surface pressure along the MAGIC transect for individual legs.

The black triangle and the black dots indicate the cloud breakup points (CB) and the starting points of the MBL systematic decoupling (DE). The red crosses in (a) indicate the location when CF_{108} drops to below 50% before CB. White spaces in (a) denote times when CF_{36} was less than 10%, and those in (b) denote missing data.

Fig. 11. Total mean of (a) the radiosonde-derived LCL along the normalized path from California to Hawaii, (b) the equivalent potential temperature difference across the inversion, (c) the mixing ratio difference at the inversion, and (d) the potential temperature difference at the inversion along the normalized path from California to DE. Gray shaded region indicates interquartile range. The boxplot in (b) and (c) indicates the location of the maximum and minimum value near DE, the difference between the two is significant at 95% confident level.

1260

1261

1262

1263 Fig. 12 The total mean (a) surface latent heat flux, and (b) surface wind speed along the
1264 normalized path from California to Hawaii. Gray shaded region indicates interquartile
1265 range.

1266

1267

1268

1269

Fig. 13. (a) Profiles of medians of U wind difference of the composed U wind east of CB and that west of CB in the small region (solid black line) and in the large region (dashed black line); same for V wind difference in the small region (solid red line) and in the large region (dashed red line). Shaded area indicates interquartile range, and (b) MBLH along normalized path from California to Hawaii.

1280

1281

	Leg												
Instrument	03A	03B	04A	04B	05A	05B	06A	06B	07A	07B	08A	08B	09A
Ka-band reflectivity													
Ka-band spectra													
W-band reflectivity													
W-band spectra													
Radar wind profiler													
HSRL													
Multipulse lidar													
MWR 2-channel													
MWR 3-channel													
ASSIST													
Total Sky Imager													
Ceiliometer													
Portable Radiation Package													
Microtops readings													
CIMEL sun photometer													
Solar Array Spectrophotometer													
Solar Spectral Flux Radiometer													
CPC													
CCN													
UHSAS													
HTDMA													
Wet/dry nephelometer													
PSAP													
Ozone													
Aerosol sampling													
Navigational information													
Meteorology													
Radiosonde launches													
Disdrometers													
IR thermometer													
ISAR													

No issue
Corrective maintenance or partial data
Questionable data
Instrument down
Not in service

1282

1283 Fig. 14 MAGIC Instrument Status from Leg03A to Leg09B

Instrument	Leg																	
	10 A	10 B	11 A	11 B	12 A	12 B	13 A	13 B	14 A	14 B	15 A	15 B	16 A	16 B	17 A	17 B	18 A	18 B
Ka-band reflectivity																		
Ka-band spectra																		
W-band reflectivity																		
W-band spectra																		
Radar wind profiler																		
HSRL																		
Multipulse lidar																		
MWR 2-channel																		
MWR 3-channel																		
ASSIST																		
Total Sky Imager																		
Ceiliometer																		
Portable Radiation Package																		
Microtops readings																		
CIMEL sun photometer																		
Solar Array Spectrophotometer																		
Solar Spectral Flux Radiometer																		
CPC																		
CCN																		
UHSAS																		
HTDMA																		
Wet/dry nephelometer																		
PSAP																		
Ozone																		
Aerosol sampling																		
Navigational information																		
Meteorology																		
Radiosonde launches																		
Disdrometers																		
IR thermometer																		
ISAR																		

No issue
Corrective maintenance or partial data
Questionable data
Instrument down
Not in service

Fig. 15 MAGIC Instrument Status from Leg10A to Leg18B

1292

1293

1294