

HOW TO FIND
THE QCD
CRITICAL POINT
AT RHIC,

IF IT IS AT $\mu_B \leq 400$ MeV

KRISHNA RAJAGOPAL (MIT)

Talks and discussions at:

INT program, August 2008

CPOD conference, June 2009

and reviews:

Koch 0810.2520; Lombardo 0808.3101;


Philipsen 0808.0672; Karsch 0711.0653;

Stephanov hep-lat/0701002


have all been very helpful as I prepared this talk.

This talk is my attempt to describe a suite of measurements that RHIC can perform that will either find experimental evidence for the QCD critical point or demonstrate convincingly that it is not at $\mu \leq 400$ MeV.

EXPLORING *the* PHASES of QCD





WHAT WE KNOW, SO FAR





WHY EXPECT A CRITICAL POINT?

- Models; lattice QCD calculations at $\mu=0$ with varying quark masses suggest:




✓ : known




- Universality class of the QCD critical point is known. (ISING)
- Experiments, and lattice calculations with $T \neq 0, \mu \neq 0$, needed to locate it.

WARNING

Nothing we know precludes pushing the critical point so far to the right that:



although models and some lattice calculations favor



LOCATING THE CRITICAL POINT...

... either via lattice calculations, or via detection of its signatures in heavy ion collision experiments,

would add a point and a line to the known QCD phase diagram.

- A qualitative leap in our understanding of QCD in the interior of its phase diagram, currently terra incognita.
- An opportunity for RHIC to write another new chapter in any future book on QCD.

LATTICE QCD WITH $T \neq 0, \mu \neq 0, \mu/T$ NOT LARGE

- $\mu \neq 0 \rightarrow$ complex Euclidean action
 - \rightarrow sign problem
 - \rightarrow difficulty of standard monte carlo $\sim \exp V$
 - Several lattice methods now in use rely on smallness of $M_q/T = \mu_B/3T$ to control the sign problem:
 - reweighting (Fodor + Katz)
 - continue from imaginary μ (de Forcrand & Philipsen; D'Elia & Lombardo)
 - Taylor expansion of P ; radius of convergence (RBC-Bielefeld; Gavai & Gupta)
- Uncertainties still dominated by systematics; (different systematics for different methods, but in all cases includes coarseness of lattice spacing)
- Steady progress; "crawling towards the continuum limit".
- Several groups exploring calculations at fixed n_B , instead of μ . (de Forcrand & Kratochvila; Li, Alexandru & Liu; ...)

LATTICE RESULTS

- via reweighting (Fodor & Katz)

$$\mu_0 = 360 \pm 40 \text{ (stat) MeV}$$

- via calculating $dmc/d\mu$ (de Forcrand + Philipsen)

$$\frac{\mu_0}{T_0} > 3 \rightarrow \mu_0 > 500 \text{ MeV}$$

- via radius of convergence of Taylor expansion

$$\frac{\mu_0}{T_c(\mu=0)} = 1.7 \pm .1 \text{ (stat) (Gavai & Gupta)}$$

$$\rightarrow 250 < \mu_0 < 400 \text{ MeV}$$

(with a "very naive" estimate of systematics)

$$\mu_0 / T_c(\mu=0) > 1.5 \text{ (RBC-Bielefeld)}$$

- STILL SYSTEMATICS DOMINATED

Nevertheless,

ONE CLEAR LESSON

Lattice calculations provide strong indications, via all algorithms employed to date, that:

$$\mu_0 > 200 \text{ MeV}$$


In the race between lattice calculations and experimental searches to locate the critical point, the lattice team is running strongly but not yet threatening to end the race.

So, let's turn to experimental searches

HOW EXPERIMENTS CAN LOCATE •

- ① At large \sqrt{s} , ie small μ , collisions equilibrate well above the crossover....
- ② Decrease \sqrt{s} in steps, moving freezeout to larger and larger μ .
- ③ Look for signatures:
 - a) Of the "lumpy" non-equilibrium final state expected after cooling through a first order transition. (Mishustin; Dumitru Paech Stöcker; Randrup; Koch Mojumder Randrup;...)
 - proton-proton angular correlations ($\Delta\phi, \Delta\eta$).
[flow turns lumps into clustering in angle]
Mocsy & Sorensen
 - enhanced event-by-event v_2 fluctuations?
 - b) Of the critical point itself. I.e. signatures of the long wavelength fluctuations occurring only near •.
Rise and then fall as $\mu \uparrow, \sqrt{s} \downarrow$.
I shall describe several such analogues of critical opalescence.


SEARCHING FOR THE CRITICAL POINT



Decreasing \sqrt{s} : decreases T and increases μ at which collision equilibrates, "landing on the phase diagram".
 \Rightarrow increases μ at which the trajectory followed by the cooling plasma crosses the transition or crossover.

TNB: location of \bullet in fig. is merely illustrative - we don't know where \bullet is!

CHEMICAL FREEZEOUT T and μ



Parametrization from Cleymans et al, 2005

ISENTROPIC TRAJECTORIES




FIG. 3: Lines of constant entropy per quark number versus μ_q/T (left) and in physical units using $T_0 = 175$ MeV to set the scales (right). In the left hand figure we show results obtained using a 4th (full symbols) and 6th (open symbols) order Taylor expansion of the pressure, respectively. Data points correspond to $S/N_B = 300, 150, 90, 60, 45, 30$ (from left to right). The vertical lines indicate the corresponding ideal gas results, $\mu_q/T = 0.08, 0.16, 0.27, 0.41, 0.54$ and 0.82 in decreasing order of values for S/N_B . For a detailed description of the right hand figure see the discussion given in the text.


Ejri Karsch Laermann Schmidt

- shape of isentropic trajectories in QGP phase and in crossover region is known from lattice calculations
- isentropic trajectories zigzag as they cross first order line

HOW EXPERIMENTS CAN LOCATE •

- ① At large \sqrt{s} , ie small μ , collisions equilibrate well above the crossover....
- ② Decrease \sqrt{s} in steps, moving freezeout to larger and larger μ .
- ③ Look for signatures:
 - a) Of the "lumpy" non-equilibrium final state expected after cooling through a first order transition. (Mishustin; Dumitru Paech Stöcker; Randrup; Koch Mojumder Randrup;...)
 - proton-proton angular correlations ($\Delta\phi, \Delta\eta$).
[flow turns lumps into clustering in angle]
Mocsy & Sorensen
 - enhanced event-by-event v_2 fluctuations?
 - b) Of the critical point itself. I.e. signatures of the long wavelength fluctuations occurring only near •.
Rise and then fall as $\mu \uparrow, \sqrt{s} \downarrow$.
I shall describe several such analogues of critical opalescence.

SEARCHING FOR THE CRITICAL POINT



Decreasing \sqrt{s} : decreases T and increases μ at which collision equilibrates, "landing on the phase diagram".
 \Rightarrow increases μ at which the trajectory followed by the cooling plasma crosses the transition or crossover.

TNB: location of \bullet in fig. is merely illustrative - we don't know where \bullet is!

HOW LOW TO GO?

Down to what \sqrt{s} should we look?

⇒ Up to what μ can we look?


This question should be answered experimentally.

Need an effect that is:

- well-measured at $\sqrt{s} = 200$ GeV
- expected only in collisions that do begin above the crossover/transition
- expected at lower \sqrt{s} , as long as collisions do begin above crossover.
i.e. jet quenching won't do since that can turn off due to absence of jets

Here are two suggestions....

n_q - SCALING OF V_2




STAR B.U.R.

- V_2 same for mesons of varying mass; same for baryons of varying mass $\Rightarrow V_2$ developed before hadrons formed
- Measurement can be done for $\pi/K/p/\Lambda$ with 5M min. bias events @ $\sqrt{s} = 11.5$ GeV with errors shown

PARITY VIOLATING FLUCTUATIONS

Need 5M min. bias events per \sqrt{s} to measure. (STAR B.U.R.)



- Data \Rightarrow charge separation!
 \Rightarrow an electric field (\perp to reaction plane, parallel or antiparallel to $\vec{L} \times \vec{B}$) that is coherent over a volume corresponding to many charged particles
- No hadronic explanation of data
- Kharzeev's explanation requires deconfinement

So.....

① Decrease \sqrt{s} in steps

② Measure the \sqrt{s} at which

- n_q scaling of v_2

- parity violating fluctuations
(charge separation)

turn off.

③ You can only look for signatures of \bullet down to, or perhaps slightly below, that \sqrt{s} .

④ In this way, learn up to what μ heavy ion collision experiments can find \bullet .

SIGNATURES OF THE CRITICAL POINT

In those collisions that pass near the critical point as they cool, find long wavelength oscillations of a mode that is a linear combination of σ (ie fluctuations couple to $\pi\pi$ and pp) and baryon number.

Fujii Ohtani; Son Stephanov

The longer the correlation length ξ gets, the bigger the signatures.

Signatures are event-by-event fluctuations of specific observables, calculable in magnitude in terms of ξ . Stephanov KR Shuryak

- Vary μ by varying \sqrt{s}
- Search for enhancement of these fluctuations in a window in \sqrt{s} , ie μ
- Analogue of critical opalescence
- Long wavelength fluctuations \Rightarrow effects greatest at low P_{\perp} .

Examples

But, first:


HOW LARGE CAN ξ GET?

HOW CLOSE TO \bullet NEED WE BE?

- Obviously ξ limited by finite size of system. But, turns out that finite time is a more severe limitation.

Berdnikov KR; Asakawa Nonaka

- Finite time spent in critical region means that even if equilibrium value of ξ is much larger, actual ξ won't grow bigger than 2-3 fm.
- Means no need to hit \bullet precisely.




Signatures will be just as big if you pass anywhere in \circ . No bigger, even if you hit \bullet .

- Hatta + Ikeda calculated "s" in a model, but did so with contours of χ_B rather than ξ . → Figs.

The robust point is that the extent of these s in μ_B is not small. Width in μ_B is ~ 100 MeV, an estimate that is both crude and uncertain. Can this be obtained on lattice??

- NB also: since ξ cannot be $> 2-3$ fm, heavy ion collision experiments can never be used to measure the critical exponents of the 2nd order critical point. That's OK: we know it is Ising. What we don't know, and need experiments for, is where it is located.

$$m_u = m_d = 5 \text{ MeV}$$



$$= \mu_B / 3$$

Hatta Ikeda

SIGNATURES OF CRITICAL POINT

- Decreasing \sqrt{s} \rightarrow Increasing μ
- Vary \sqrt{s} , and hence μ , and look for nonmonotonic enhancement (rise and then fall) of:
 - i) Event-by-event fluctuations of mean P_T of low P_T pions
 - ii) Event-by-event fluctuations of net proton number ($N_p - N_{\bar{p}}$)
 - iii) Event-by-event fluctuations of particle ratios involving pions and/or protons
 - iv) Kurtosis of the N_p or $(N_p - N_{\bar{p}})$ event-by-event distribution

⋮
In all cases, the enhancement should be greater at lower P_T .

MEAN P_T OF LOW P_T PIONS

Stephanov KR Shuryak (1999)

First example of a quantitative connection between long wavelength fluctuations of the chiral order parameter with correlation length ξ and magnitude of event-by-event fluctuations of an experimental observable.


DISADVANTAGES:

- Effect predicted is not large for $\xi = 3 \text{ fm}$
- Will fluctuations in P_T survive the late time hadronic gas? Will they get washed out between chemical and kinetic freezeout?

RESULT:




NA49 has done a beautiful analysis and sees no \sqrt{s} dependence....
CERES has done a beautiful analysis and sees no \sqrt{s} dependence....

EVENT-BY-EVENT FLUCTUATIONS OF MEAN p_T



CERES, Nucl Phys A, 2008


EVENT-BY-EVENT FLUCTUATIONS OF MEAN p_T OF PARTICLES WITH p_T BELOW A CUT



No enhancement at low p_T .

NA49, 2008

MEAN p_T FLUCTUATIONS, vs. μ



NA49, 2008

Height of solid (dashed) curve is magnitude of effect predicted for $\xi = 3$ fm ($\xi = 6$ fm).

POSSIBLE CONCLUSIONS

- $\mu_0 > 470 \text{ MeV}$?
 - P_T fluctuations washed out.
Predicted effect was not large, and was susceptible to being erased after chemical freezeout.
- ⇒ Look for event-by-event fluctuations of other observables
- a) for which the predicted effect of proximity to the critical point is larger
 - b) which cannot be washed out after chemical freezeout

BARYON, AND PROTON, NUMBER FLUCTUATIONS


Hatta Ikeda; Hatta Stephanov

- seen on the lattice \rightarrow FIG
- should be looked for in
experimental data


$\frac{\partial^2 \Omega}{\partial \mu_B^2} \rightarrow B\text{-fluctuations}$
 $\frac{\partial^2 \Omega}{\partial \mu_B^2} \sim \xi^2 + \text{nonsingular}$

$\frac{\partial^2 \Omega}{\partial \mu_I^2} \rightarrow \text{no enhanced (u-d)}$
 $\frac{\partial^2 \Omega}{\partial \mu_I^2} \sim \text{nonsingular}$

Suggests
 $\mu_q \sim 1$
 T
 getting
 close
 to \bullet .



(a)



(b)

Figure 3.3: The quark number susceptibility χ_q/T^2 (left) and isovector susceptibility χ_I/T^2 (right) as functions of T/T_0 for various μ_q/T ranging from $\mu_q/T = 0$ (lowest curve) rising in steps of 0.2 to $\mu_q/T = 1$, calculated from a Taylor series in 6th order. Also shown as dashed lines are results from a 4th order expansion in μ_q/T .

(Because B fluctuates while isospin does not, proton fluctuations \sim B fluctuations)

Hatta Stephanov

Ejiri et al

PARTICLE RATIOS

NA 49

- Originally motivated by peak in $\langle K \rangle / \langle \pi \rangle$ at $\sqrt{s} = 7.6$ GeV.

To better understand this, look at fluctuations of K/π ratio.

- Now motivated by observation that these fluctuations will better survive the late time hadron gas.

RESULT:

- Large K/π fluctuations at $\mu_B \sim 400 - 450$ MeV

- Why no P/π fluctuations ???

EVENT-BY-EVENT FLUCTUATIONS OF K/ π AND p/ π




FIG. 8: Energy dependence of the event-by-event non-statistical fluctuations of the K/π ratio (top panel) and the $(p + \bar{p})/\pi$ ratio (bottom panel). Filled symbols show data, open symbols show calculations with the UrQMD transport code, using NA49 acceptance tables. Systematic uncertainties are shown as brackets.

NA49, 2008

- Large event-by-event fluctuations in K/π ratio at $\mu \sim 350-450$ MeV
- Intriguing, but puzzling
 - error bars still substantial
 - interpretation is complicated by change in # of accepted kaons (& pions) as \sqrt{s} changes. ^{Koch Schuster Westfall}
 - why no enhancement in p/π fluctuations?
 - K/π fluctuations apparently not driven by low p_T pions


STAR has used this observable as a case study to see how they can improve on this measurement with a beam energy scan at RHIC.

A collider has advantages...

RHIC Critical Point Search Program - Advantage

Hadron Mass

Beam Energy



200 GeV

62.4 GeV


9.2 GeV

Uniform acceptance for different particle species and for different beam energies in the same experimental setup (advantage over fixed target expt.)³⁶

Mohanty, QM 09

Old Program : Fixed Target

Hadron Mass



NA49 : arXiv : 0808.1237

6.3 GeV

7.6 GeV

8.7 GeV

12.3 GeV

17.3 GeV

Beam Energy

Non-Uniform acceptance for different particle species
and for different beam energies in the same experimental setup

K/ π FLUCTUATIONS WITH A RHIC BEAM ENERGY SCAN




Fig. 3-7: Estimate of the error in σ_{dyn} for charge-integrated K/ π fluctuations, based on 100K central events analyzed in the STAR detector (with the newly completed ToF). Shown for comparison are the current measurements from NA49 and STAR.

STAR B.U.R.

Error bars with 1M min bias events
per energy

KURTOSIS OF EVENT-BY-EVENT

DISTRIBUTION OF (NET) PROTON NUMBER

Stephanov; "a direct consequence of discussions" at Aug 2008 INT workshop

Critical fluctuations couple to $\pi\pi$, PP
→ event-by-event fluctuations in their multiplicities, multiplicity ratios, P_T , that are $\propto \xi^2$

Higher moments of the event-by-event distributions receive effects that are more sensitive to ξ .

Skewness $\propto \xi^{4.5}$

Kurtosis $\propto \xi^7$!!!

The prefactors work out particularly nicely for kurtosis of proton distribution, but Stephanov makes predictions for π & p , skewness & kurtosis.

DEFINITIONS

$N = \#$ of protons (or π ; or p - \bar{p}) in 1 event

$\bar{N} \equiv \langle N \rangle = \text{mean}$

$\delta N \equiv N - \bar{N}$ in one event


$\langle (\delta N)^2 \rangle = \text{variance}$

$$K^{\text{eff}} \equiv K \langle (\delta N)^2 \rangle \equiv \frac{\langle (\delta N)^4 \rangle - 3 \langle (\delta N)^2 \rangle^2}{\langle (\delta N)^2 \rangle}$$

Kurtosis $\sim \frac{1}{N}$

Variance $\sim \bar{N}$


In lattice QCD literature, K^{eff} is called χ_4 / χ_2 or χ_4 / χ_2 . It is calculated for $N = \text{baryon number}$, or $N = \text{charge}$.



Gaussian: $K = 0$

EFFECT OF CRITICAL FLUCTUATIONS

Stephanov



$$K^{\text{eff}} \left(\frac{\bar{N}}{\langle (SN)^2 \rangle} \right) = 23 \left(\frac{2\tilde{\lambda}_3^2 - \tilde{\lambda}_4}{50} \right) \left(\frac{g}{10} \right)^4 \left(\frac{\xi}{1 \text{ fm}} \right)^7$$

$\underbrace{\hspace{10em}}_{\approx 1}$
 $\underbrace{\hspace{10em}}_{\theta(1)}$
 $\underbrace{\hspace{10em}}_{!!!}$


$\tilde{\lambda}_3, \tilde{\lambda}_4$: Obscure but universal constants that depend on θ . Known for Ising \bullet .

g : σ PP coupling. $g \sim m_p/f_\pi \sim 10$

So, how big is the background? Theory suggests $\theta(1)$, but better to determine it experimentally....

K_{eff} AT $\sqrt{s} = 200 \text{ GeV}$ AND 62 GeV

ie at very low μ , far from ●



Nayak, QM 09

- Background K_{eff} measured, $\mathcal{O}(1)$.
- Error bars $\approx \pm 0.1$.

PROTON NUMBER KURTOSIS

- Predicted effect of proximity to \bullet is:
 - large
 - strongly \sqrt{s} -dependent, $\therefore \sqrt{s}$ -dependent
 - larger at lower P_T
- How you know its \bullet if you see it.
- Cannot be washed out between chemical & kinetic freezeout
- Background measured, and small, at $\sqrt{s} = 62 \text{ \& } 200$.
- Error bars $\sim \pm 1$ with 5M events STAR BULK
- Calculations done for $\bar{N} \rightarrow \infty$. Finite \bar{N} and acceptance corrections need to be assessed. (Latter minimal for STAR)
- Analogous analyses can be done for skewness, and for pions. Many signatures will be in play if/when \bullet found.

A BEAM ENERGY SCAN TO SEARCH FOR ●

\sqrt{s}/A (GeV)	μ (MeV)*	8 hr days † per 5M events
5	550	
6.2	485	
7.7	420	56
9.8	355	30
12.7	290	13
17.3	230	5
27	155	2
39	110	1
200	25	
		<u>107 days</u>

*: from Cleymans et al's 2005
empirical fit to compilation of data

†: from STAR B.U.R.

WHAT NEED BE MEASURED AT EACH \sqrt{s}

- Enough \langle particle ratios \rangle to evaluate μ .
So you know where on phase diagram you are freezing out.
- n_q scaling of V_2 for $\pi/K/P/\Lambda$
parity violating fluctuations
So you know whether collision got above the crossover/transition
- Event-by-event fluctuations of K/π and P/π ratios with significantly smaller error bars than in NA49 data
- Variance, skewness & kurtosis of event-by-event distributions of N_p , $N_p - N_{\bar{p}}$, N_π .
- All can be done with 5M min bias events per \sqrt{s} . STAR B.U.R.

CAN WE DISCOVER THE QCD CRITICAL POINT IN RUN 10 AT RHIC?

YES, IF:

Nature is kind, and puts $\mu_c \lesssim 420 \text{ MeV}$

IF YES:

- The landmark discovered. Our map of the QCD phase diagram then anchored by experiment.
- Assuming reasonable progress in lattice QCD, quantitative comparison between theory & experiment for μ_c will come.
- RHIC (and FAIR) can study the first order phase transition. RHIC can further probe the region around \bullet .

IF NO:

- We learn that $\mu_c > 420 \text{ MeV}$
- We will be able to make a data-driven decision about whether to run at $\sqrt{s} = 6.2$ and 5 in a future year.

Collision Energies (GeV)		5	7.7	11.5	17.3	27	39
Section	Observables	Millions of Events Needed					
A1	v_2 (up to ~ 1.5 GeV/c)	0.3	0.2	0.1	0.1	0.1	0.1
A1	v_1	0.5	0.5	0.5	0.5	0.5	0.5
A2	Azimuthally sensitive HBT	4	4	3.5	3.5	3	3
A3	PID fluctuations (K/ π)	1	1	1	1	1	1
A3	net-proton kurtosis	5	5	5	5	5	5
A3	differential corr & fluct vs. centrality	4	5	5	5	5	5
A3	integrated p_T fluct (T fluct)						
B1	n_q scaling $\pi/K/p/\Lambda$ ($m_T - m_0$)/ $n < 2$ GeV		6	5	5	4.5	4.5
B1	ϕ/Ω up to $p_T/n_q = 2$ GeV/c		56	25	18	13	12
B2	R_{CP} up to $p_T \sim 4.5$ GeV/c (at 17.3) 5.5 (at 27) & 6 GeV/c (at 39)				15	33	24
B3	untriggered ridge correlations		27	13	8	6	6
B4	parity violation		5	5	5	5	5
See[1]:	charge-photon fluctuations (DCC)	1	1	1	1	1	1
	kink/step/horn	0.1	0.1	0.1	0.1	0.1	0.1
	v_2 fluctuations	0.5	0.5	0.5	0.5	0.5	0.5
	HBT ($R_l, R_o/R_s$)	0.8	0.8	0.5	0.5	0.5	0.5
	Jet/ridge $2 < \text{trig} < 4, 1 < \text{assoc} < \text{trig}$				30	8.8	4.5
	Jet/ridge $3 < \text{trig} < 6, 1.5 < \text{assoc} < \text{trig}$					53	24
	Baryon-Strangeness cor (hypernuc)						50
	Forward π^- yield (rapidity scaling)						
	Forw. $\gamma(\pi^0)$ yield (rapidity scaling)						
	Long-range forward-backward corr.						
	Other PID fluctuations (esp. K/p)						
	Particle ratios (many examples)						
	p_T spectra						
	Prod. of light nuclei & antinuclei						
	Yields of species & stat model fits						

Table 3-2: Observables and statistics needed for the first BES run. The observables in the yellow-shaded area relate to the search for a phase transition or critical point (see section A), while observables in the blue-shaded area search for turn-off of new phenomena already established at higher RHIC energies (see section B). The numbers listed in boldface above are all within reach (nominally require no more than 1.5 times the proposed statistics) in the first BES run plan as set out in Table 3-1. The remaining numbers (not boldface) will need to wait for higher

MULTIPLICITY FLUCTUATIONS

Also coupled to order parameter fluctuations. In fact their effect on multiplicity fluctuations is greater than on P_T fluctuations.

Stephanov KR Shuryak

BUT: large "background", due to impact parameter fluctuations.

Still, nonmonotonic variation with \sqrt{s} would be suggestive.

Order of Phase Transition for $\mu_B \sim 0$

Physical quark masses

Continuum limit



Simulations along Lines of Constant Physics

$m_K/m_\pi = 3.689$; $f_K/m_\pi = 1.185$

Staggered fermionic action

$$\chi(N_s, N_t) = \partial^2 / (\partial m_{ud}^2) (T/V) \cdot \log Z$$

Y. Aoki et al., Nature443:675-678,2006



T grows with $6/g^2$, g : gauge coupling

No significant volume dependence (8 times difference in volumes)

Phase transition at high T and $\mu_B = 0$ is a cross over

1st order :

Peak height $\sim V$

Peak width $\sim 1/V$

Cross over :

Peak height $\sim \text{const.}$

Peak width $\sim \text{const.}$

2nd order :

Peak height $\sim V^\alpha$

Lattice results on electroweak transition in standard model


is an analytic cross-over for large Higgs mass

Relevant to LHC and
current RHIC regimes


K. Kajantie et al., PRL 77, 2887-2890,2006

Method to determine the CEP:

- find largest temperature where all expansion coefficients are positive $\rightarrow T^{\text{CEP}}$
- determine the radius of convergence at that temperature $\rightarrow \mu^{\text{CEP}}$



\rightarrow first non-trivial estimate of T^{CEP} from c_8
 second non-trivial estimate of T^{CEP} from c_{10}



$$\mu^{\text{CEP}} = \lim_{n \rightarrow \infty} \mu_n^{\text{CEP}}$$


$$\mu_n^{\text{CEP}} = T^{\text{CEP}} \sqrt{c_n / c_{n+2}}$$

The Resonance gas limit:

$$\frac{p}{T^4} = G(T) + F(T) \cosh\left(\frac{\mu_B}{T}\right)$$

$$\rightarrow \rho_n = \sqrt{(n+2)(n+1)}$$

K_{eff}



Nayak, QM09

$m_u = m_d = 0$

MODEL ANALYSIS OF EXTENT OF CRITICAL REGION

