The RHIC Experiments: *what can they do (forward)?*

Kenneth N. Barish
UC Riverside

Forward physics at RHIC workshop
2004 RHIC/AGS Users Meeting, BNL
14 May, 2004

Regions of a nucleus



CGC boundary

$Q_S^2 = Q_{S(y=0)}^2 e^{\lambda y}$

$\lambda \sim 0.3$

CQF boundary

$Q_S^2 = Q_{S(y=0)}^4$

Understanding proton spin structure

- Need to understand dynamics of the nucleon spin:
  - Contribution of gluons to proton spin (ΔG)
  - Gluon feed-down to quark sea (measure $\Delta\bar{q}/q$ and $\Delta q/q$)
    - W production unique in providing theoretically clean access

- A_N possibly gives us access to the transversity distribution, orbital angular momentum, and higher twist effects



Spin $\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L_z$

Quark Spin

Gluon Spin

Orbital Angular Momentum

Flavor decomposition

Drell-Yan production of lepton pairs

- Maximal parton level asymmetry: $a_{LL} = -1$
- Possible severe background from semi-leptonic decays of open charm productions – can be improved with forward silicon tracking.



$$A_{LL}^{q\bar{q}\to\ell\bar{\ell}} = \frac{\Delta q(x_A)}{q(x_A)} \otimes \frac{\Delta\bar{q}(x_B)}{\bar{q}(x_B)} \otimes \hat{a}_{LL}^{q\bar{q}\to\ell\bar{\ell}}$$

W production

- Produced in parity violating *V-A* process
  - Chirality / helicity of quarks defined
- Couples to weak charge
  - Flavor almost fixed: flavor analysis possible
  - Flavor ID reduces uncertainty in current pol-PDF models.
- E.g., PHENIX-Muon Arms



$$A_L^{W^+\to\mu^+\nu_\mu} = \frac{\Delta u(x_a)\bar{d}(x_b) - \Delta\bar{d}(x_a)u(x_b)}{u(x_a)\bar{d}(x_b) + \bar{d}(x_a)u(x_b)}$$



ΔG and forward detectors



Experimental access to low-x gluons

Gluons at low-x

- » pp & pA
  - – $\gamma^* \rightarrow \mu\mu$, ee
  - – Prompt photons
  - – Open Charm
  - – Hadrons
  - – J/ψ production
  - – correlations

Map out previous diagrams



Coverage of experiments



charged hadrons
A_N

γ, π^0, jets, charm/beauty
A_N, A_L, A_{LL}

γ, π^0, jets, e, μ, $\mu\mu$, charm/beauty, eμ
A_N, A_L, A_{LL}

Large η coverage is needed



by Anuj K. Purwar

PHOBOS Detector System



Pseudorapidity Distributions at 200 GeV

• d + Au at 200 GeV Min-Bias



• p + p at 200 GeV



Cronin Effect as a Function of η (d+Au)





The data at forward rapidities was collected with FFS at 4º ($\eta \sim 3$) and 12º ($\eta \sim 2$)

High p_T suppression for different η's









STAR Detector



TPC: -1.0 < η < 1.0

FTPC: 2.8 < |η| < 3.8

FPD: |η| ~ 3.8 (p + p)

|η| ~ 4.0 (p + p and d + Au)



Forward π^0 Detector (FPD)

- **Pb-glass EM calorimeter**
- **Shower-Maximum Detector (SMD)**
- **Preshower**

Forward π^0 Inclusive Cross Section



- STAR data at
  - ⟨η⟩= 3.8 (hep-ex/0310058, accepted to PRL, in press)
  - ⟨η⟩= 3.3 (hep-ex/0403012, Preliminary)
- NLO pQCD calculations at fixed η with equal factorization and renormalization scales = p_T
- Solid and dashed curves differ primarily in the g → π fragmentation function

STAR data consistent with Next-to-Leading Order pQCD calculations

in contrast to data at lower √s (Bourelly and Soffer, hep-ph/0311110)

A_N for forward π^0 production at $\sqrt{s}$=200GeV

STAR collaboration, hep-ex/0310058,
accepted by Phys. Rev. Lett. (in press)



In agreement with several models including different dynamics:

- Sivers: spin and $k_\perp$ correlation in initial state (related to orbital angular momentum?)
- Collins: Transversity distribution function & spin-dependent fragmentation function
- Qiu and Sterman (initial-state) / Koike (final-state) twist-3 pQCD calculations

- p_T dependence?
- x_F<0?
- A_N with mid-rapidity correlation?
- Spin dependence in jet?
- Heavy flavors??
- Related to "gluon saturation"???



Trigger by forward π^0

- $E_p > 25$ GeV
- $\langle \eta_\pi \rangle = 4$

STAR

Large rapidity gap π^0+$h^\pm$ correlation data

Midrapidity $h^\pm$ tracks in TPC

- $-0.75 < \eta < +0.75$

Large rapidity gap π^0 +h correlation data are suppressed in d+Au relative to p+p at small <xF> and <pT,p> Consistent with CGC picture



Coverage $1<\eta<3$

Jets direction and energy

photons

π^0's

electrons

detached vertices for charm and bottom

Forward Sillicon Tracking detectors



Central arm upgrades

Aerogel Counter-PID
RPC – for TOF
Barrel Sillicon tracking
Hadron Blind Detector
–Datliz rejection

Nosecone Calorimeters

PHENIX Forward Upgrade Components

Endcap Vertex Tracker

» silicon pixel detectors

Nosecone Calorimeter

» W-silicon (only 20cm long)

» shower max (Si)

Muon trigger

» U-tracker (RPC or MuTr)

» D-tracker (RPC's)





W and quarkonium:
improved μ-trigger rejection

K. Barish

Space is tight …



Challenges:

-only 40 cm apart from collision point …

-only 20 cm of space is available

Design thoughts

PHENIX preliminary design:

14 cm of W absorber

+

6 cm Si readout

Calorimeter: ~	**40 L_{rad} / 1.6 L_{abs}**
Lateral seg.	**1.5 x 1.5 cm^2**
EM section:	**10 Lrad**
γ/π^0 identifier:	**at a depth ~ 5Lrad (two layers of ~2x60 mm^2 silicon strips)**

- Good γ/π^0 separation
- Good position resolution
- e/h separation
- Reasonable jet measurements
- Triggering capability

Notes:.

- AuAu is challenging
- Shower max layers required
- Need upstream tracking to recover J/Ψ resolution in AuAu

This is a huge challenge, but not completely crazy!!

Coverage of NCC at 200 GeV

0.5 pb^{-1} pAu (run 12)

$\Delta\log(x_2)$ ΔQ^2=0.1x1 GeV^2

Direct photon



Level-1 triggering scheme



Summary



charged hadrons
A_N

γ, π^0, jets, charm/beauty
A_N, A_L, A_{LL}

γ, π^0, jets, μ, $\mu\mu$, charm/beauty, eμ

A_N, A_L, A_{LL}

Extra slides …

Semi-inclusive DIS



Hermes: Phys.Rev.Lett 92 (2004) 012005

- In addition to the scattered electron, one measures in addition, the hadronic final state(s) – leading K's and π's
- Access to flavor separated parton distributions



Comments:

- Treated only at LO so far
- Q^2 of 2.5 GeV^2 is a concern
- Higher twist effects not included
- u-quark "dominance" limits precision of d and s-quark (and anti-quark) polarized distributions.
- "Hadron taging" technique relies on knowledge of fragmentation functions
- **Measurements at the highest Q^2 (RHIC) are important.**
- **W production (at RHIC) is an independent and eloquent technique**

Sensitivity goal: flavor decomposition



$A_L(W^+) \rightarrow \Delta u/u(x_1)$ @$x_1 \gg x_2$

$A_L(W^+) \rightarrow -\Delta d/d(x_1)$ @$x_2 \gg x_1$

(W^+ and W^- have a different acceptance)

- The measurement of inclusive single spin muon asymmetries is the least biased way to probe $\Delta q/q$, $\Delta\bar{q}/\bar{q}$.
  - Complete theoretical treatment from first principles by Nadolsky and Yuan using re-summation and NLO techniques [NuclPhysB 666(2003) 31].
  - PHENIX is in the process of re-evaluating their sensitivities using their NLO code.
- Measurement requires a trigger with rejection ~10,000.
  - Present muon trigger has a rejection of 250 (run 3).
  - Upgrade needed … requires momentum determination at level-1 to reject background due to hadron-decay's into μ's and beam related backgrounds.
- Nose-cone calorimeter important to help reject hadron jets off-line